Aberystwyth University

*Microsatellite genotyping of brown crab *Cancer pagurus* reveals fine scale selection and 'non-chaotic' genetic patchiness within a high gene flow system*

McKeown, Niall; Hauser, Lorenz; Shaw, Paul

Published in:
Marine Ecology Progress Series

DOI:
[10.3354/meps12044](https://doi.org/10.3354/meps12044)

Publication date:
2017

Citation for published version (APA):
McKeown, N., Hauser, L., & Shaw, P. (2017). Microsatellite genotyping of brown crab *Cancer pagurus* reveals fine scale selection and 'non-chaotic' genetic patchiness within a high gene flow system. *Marine Ecology Progress Series*, 566, 91-103. <https://doi.org/10.3354/meps12044>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Title: Microsatellite genotyping of Brown crab (*Cancer pagurus*) reveals fine scale selection and “non- chaotic” genetic patchiness within a high gene flow system.

Running page head: brown crab biocomplexity

Authors: Niall J McKeown^{1*} – Lorenz Hauser² – Paul W Shaw¹

¹ Institute of Biological, Environmental and Rural Sciences (IBERS), Aberystwyth University, UK, SY23 3DA.

² School of Aquatic and Fishery Sciences, University of Washington, Washington 98195-5020, USA.

*Corresponding author email: njm2@aber.ac.uk

ABSTRACT: Brown crab, *Cancer pagurus*, supports one of the most important European fisheries, however, spatial patterns of connectivity and adaptation are largely unknown and difficult to identify due to the species’ life history which entails distinct dispersal characteristics during larval and adult life stages. To address this limitation spatial-temporal genetic structure was assessed, using 8 microsatellite loci, across the majority of the species’ NE Atlantic distribution. Neutral genetic structuring revealed a background of high gene flow throughout the region, with a superimposed pattern of chaotic genetic patchiness (CGP) linked to stochastic recruitment variability. The CGP was geographically patterned, being prevalent among English Channel samples and absent among North Sea samples, suggesting specific biological (e.g. reproductive ecology) and environmental (seascape) drivers. Such recruitment variability may compromise stock resilience and must be considered within spatial management strategies. Another prominent feature was the pronounced differentiation of males sampled within a fjord (Gulmarsfjord) from all other samples, at a single locus exhibiting the effects of divergent selection. Gulmarsfjord females were genetically similar to all other ‘non-fjord’ samples, and exhibited a comparative level of differentiation at the outlier locus from the Gulmarsfjord males. Due to known dispersal differences between the sexes, the pattern within Gulmarsfjord can be explained by the intermingling of allochthonous females with resident, locally adapted, males and demonstrates the occurrence of fine-scale local adaptation in this species. Collectively, the study highlights how considerable intraspecific eco-evolutionary diversification can occur despite high levels of dispersal / gene flow.

Key words: adaptation – gene flow – dispersal – sweepstakes recruitment – conservation - sustainability

35

36

INTRODUCTION

37 Genetic studies have yielded many insights into marine intraspecific biodiversity with
38 important findings including the detection of significant genetic population structuring (Shaw
39 et al. 1999, Knutsen et al. 2011, McKeown et al. 2015) and adaptation (Hemmer-Hansen et
40 al. 2007a, Poulsen et al. 2011, Therkildsen et al. 2013a) in systems where high gene flow
41 would be expected to prevent such differentiation (Palumbi 1994, Waples 1998). Population
42 genetic structure and adaptation, as components of intraspecific biocomplexity, are thought to
43 be significant factors underpinning species / population sustainability and evolutionary
44 potential (Iles & Sinclair 1982, Ryman et al. 1995, Ruzzante et al. 2006, Therkildsen et al.
45 2013b). The ongoing depletion of marine populations through fishing, and demographic
46 changes associated with predicted future climate change, are adding to the impetus to resolve:
47 (i) spatial / temporal patterns of neutral and adaptive genetic structure (Reiss et al. 2009); (ii)
48 historical and contemporary drivers of structuring (Hemmer-Hansen et al. 2007b); and (iii)
49 the significance of such diversity on ecological and evolutionary timescales (Stepien et al.
50 2009).

51 While there has been considerable research on fishes with mobile larval and non-
52 larval stages, genetic studies of crustaceans have typically focused on taxa with sedentary
53 adults (Jorde et al. 2015). In this context the brown crab, *Cancer pagurus* (L.), occurring
54 continuously in shallow shelf waters of the NE Atlantic from the Lofoten Islands (Norway) to
55 Morocco (Bennett 1995) and supporting one of the most important commercial European
56 fisheries, represents an interesting candidate for investigation as both larval and adult stages
57 have potential for substantial dispersal. Adults are described as benthic and mobile, but there
58 are pronounced dispersal differences between the sexes: males being largely resident, making

short random movements within small territories, while females migrate significantly longer distances, and more frequently, than males (Edwards 1979, Bennett & Brown 1983, Latrouite & Le Foll 1989, Ungfors et al. 2007). In the English Channel female migrations of up to 200 nautical miles have been reported with some crabs achieving a mean speed of 1.07-1.62 nautical miles per day (Pawson 1995). The pelagic larval stage lasts for approximately three months (Eaton et al. 2003, Weiss et al. 2009, Hunter et al. 2013) and while little is known about the ecology of juveniles they are rarely caught in offshore waters, suggesting that adult crabs only move to deeper water as they grow and reach maturity.

Tagging studies have revealed that adult female migrations are consistently against prevailing currents (Hunter et al. 2013, Ungfors et al. 2007). As the larvae are poor swimmers likely to passively drift while entrained in currents, it has been suggested that contranatal female migrations are spawning behaviours aimed at facilitating return to areas of maternal origin. Even in the absence of additional extrinsic factors, the seemingly counter active dispersal of females and larvae is expected to limit 'life-time dispersal' and may thus influence spatial patterns of recruitment and structuring of reproductive populations. Tagging, fishery landings data and sex-specific growth rates variously suggest some demographic independence between the areas of brown crab abundance in the Celtic Sea, English Channel, North Sea and Bay of Biscay (Pawson 1995). Within the North Sea, the seasonal jet-like circulation associated with the Flamborough front is predicted to prevent exchange of larvae between areas north and south of the front during spawning time (Eaton et al. 2003). In the English Channel, larval surveys have reported distinct western and eastern centres of larval abundance separated by a central area of low / no larval occurrence, and hydrodynamic modelling has indicated insufficient larval transport rates to connect these spawning areas (D. Eaton, unpublished data).

Population genetic structure of Brown crab has to date been studied only in Scandinavian waters, where Ungfors *et al.* (2009) reported no significant genetic differentiation among samples spanning 1300km of waterway distance within the Norwegian Sea, Skagerrak and Kattegat. However, genetic structuring may vary throughout a species' range and failure to identify local populations may lead to local overfishing and ultimately severe declines. While the females are highly fecund (0.5 - 2.9 million eggs per brood; Edwards 1979, Ungfors 2007), paternity analysis suggests single paternity of broods (McKeown & Shaw 2008b). Such a reproductive ecology, alongside the selective harvesting of females (Bennett 1995), which are currently regarded as overexploited, may enhance the susceptibility of brown crab to genetic erosion (McKeown & Shaw 2008b).

The objective of the present study was to test the general hypothesis of genetic panmixia in brown crab throughout a considerable portion of the species' range, with a specific focus on the English Channel and North Sea. Some genetic studies of crustaceans have reported macro-geographical homogeneity with structuring apparent only at regional scales (e.g. Domingues *et al.* 2010), while other studies have reported fine scale spatial / temporal genetic structuring (Selkoe *et al.* 2010). To encapsulate such potential complexity broad- and fine-scale spatial-temporal patterns were assessed. Furthermore, comparative analyses of males and females were performed to identify differences that may be associated with sex-specific ontogenetic movements. The sampling strategy also encompassed distinct seascape features (e.g. samples collected within semi-enclosed water bodies such as bays and fjords) to examine the effect of local hydrodynamic environments. This sampling design permitted interpretation of the mechanistic underpinnings and eco-evolutionary significance of complex patterns of genetic diversity which included evidence of broad-scale genetic connectivity, fine-scale adaptive divergence of a fjord sample, and regional variation in genetic patchiness.

108

109

MATERIALS AND METHODS

110

111

112

113

114

115

116

117

Sample collection and molecular analyses. Spatial/temporal sampling of adults throughout the NE Atlantic was performed using both research (CEFAS) and commercial vessels (See Table 1 and Fig. 1 for sample information). For each sample crabs were captured using multiple baited pots within a localised area (maximum distance among pots ~200 metres) over a single day with tissue biopsies preserved in ethanol. Although adult crabs cannot be reliably aged, samples were considered to consist of multiple age cohorts. For the samples collected on board CEFAS vessels, the majority of individuals were identified as male or female, which permitted downstream separation in statistical analysis.

118

119

120

121

Total DNA was extracted using a standard CTAB-chloroform/isoamylalcohol method (Winnepeenninckx *et al.* 1993). All individuals were typed at eight microsatellite loci (Cpag15, Cpag1b9, Cpag2a5-2, Cpag3a2, Cpag3d7, Cpag4, Cpag5d8, Cpag6c4b) following McKeown & Shaw (2008a).

122

123

124

125

126

127

128

129

130

131

Statistical analysis. Genetic variation within samples was characterised using number of alleles (N_A), allelic richness (A_R ; El Mousadik & Petit 1996), observed heterozygosity (H_O), and expected heterozygosity (H_E) (Nei 1978), all calculated using GENALEX 6.2 (Peakall & Smouse 2006). Genotype frequency conformance to Hardy-Weinberg equilibrium (HWE) expectations and genotypic linkage equilibrium between pairs of loci were tested using exact tests (10,000 batches, 5000 iterations) in GENEPOP 3.3 (Raymond & Rousset 1995). Deviations from HWE were measured using F_{IS} , calculated according to Weir & Cockerham (1984) and tested for significance by 10,000 permutations in FSTAT 2.9.3. (Goudet, 1995). Mean pairwise relatedness within samples was calculated using the relatedness estimator, r_{qg} , of Queller & Goodnight (1989) in GENALEX with associated 95% confidence intervals

determined by 1000 bootstraps. Permutation of genotypes among all samples (999 times) was used to calculate the upper and lower 95% confidence intervals for the expected range of r_{qg} under a panmictic model.

Genetic differentiation was quantified by global and pairwise F_{ST} values, with associated significance evaluated by 10,000 permutations (Goudet et al. 1996), using FSTAT. Hierarchical analysis of molecular variance (AMOVA; Excoffier et al. 1992) was performed in ARLEQUIN to partition genetic variance among groups of samples (F_{CT}) and among samples within groups (F_{SC}) with significance levels of F_{CT} and F_{SC} tested using 1000 permutations. To help visualise F_{ST} results principal coordinates analysis (PCoA) was performed on pairwise matrices. Mantel tests, as implemented in the IBDWS software (Jensen et al. 2005) were used to test for correlation between pairwise linearised F_{ST} [$F_{ST}/(1-F_{ST})$] (Rousset 1997) and shortest sea distances between sample sites (i.e. isolation by distance - IBD). IBD tests were based on 10,000 randomisations and performed on combinations of untransformed and log transformed genetic and geographical distances for various pooled and partitioned arrangements of temporal, male and female samples. Differentiation between samples was tested with global and pairwise exact G -tests in GENEPOP (10,000 batches, 5000 iterations). The simulation method implemented in POWSIM (Ryman & Palm 2006) was used to estimate the sample size dependent Type I and Type II error probabilities of the exact G -tests. Genetic structuring was also investigated using the Bayesian clustering method in STRUCTURE (Pritchard et al. 2000) both with and without prior population information and with multiple parameter sets (i.e. with and without admixture, and with and without correlated allele frequencies). Randomisation procedures in FSTAT were used to detect significant differences in heterozygosity, A_R , F_{IS} , F_{ST} and relatedness among user defined groups of samples following 10,000 permutations.

The assumption of selective neutrality of the microsatellite loci was assessed using the FDIST outlier identification test (Beaumont & Nichols 1996) implemented in LOSITAN (Antao et al. 2008) performed (i) globally (i.e. across groups of samples) and (ii) between pairs of samples. Simulations were run for 10,000 replications, 95% confidence intervals estimated using the options for neutral and forced mean F_{ST} . No differences were detected between analyses assuming infinite allele model (IAM) and stepwise mutation models and so only IAM results are presented.

RESULTS

Intrasample genetic variability and data power

A total of 2,777 individuals were assayed (mean sample size = 81.7), with an average of 17.4 alleles detected per locus (range 5-34). Loci Cpag4 and Cpag5d8 had considerably more alleles ($n = 34$ in both cases) than the other loci, with the next highest allele number reported for Cpag1b9 (19 alleles). Each locus was polymorphic in all samples with very similar levels of variation across all samples. No significant linkage disequilibrium between loci was detected, either across all samples (data pooled) or in any single sample. Single locus tests for conformity to Hardy Weinberg equilibrium (HWE) expectations for each of the initial 34 samples (Table 1) revealed the largest number of significant deviations (at critical P value = 0.05) for Cpag4 and Cpag3A2 which exhibited 19 and 12 significant test results, respectively. No other locus exhibited more than 5 deviations out of 34 tests (at critical $P = 0.05$). Multilocus tests of HWE and associated F_{IS} values were non-significant in most samples (Table 1). All single- and multi-locus deviations from HWE were due to heterozygote deficits. Application of MICRO-CHECKER (Van Oosterhout et al. 2006) algorithms to adjust for potential null alleles in cases of single locus heterozygote deficits

resulted in no change to the magnitude and pattern of genetic differentiation revealed in subsequent tests and so results for unedited data are reported. Mean intra sample relatedness conformed to predictions of a panmictic model for all but two samples (Table 1).

POWSIM analysis indicated both considerable statistical power for G tests to detect population structure and low type I error rates, for various sample size permutations relative to this study (Table 2).

Detection of divergent selection effects

Significant differentiation was detected between males and females collected in the Gulmarsfjord ($F_{ST} = 0.027$, $P < 0.0001$; exact G test $P < 0.0001$). This differentiation was driven by a single locus (Cpag6c4b) which yielded a pairwise F_{ST} of 0.160 ($P = 0.0001$; Exact G test $P = < 0.0001$). Differentiation between the sexes was not significant upon exclusion of this locus ($F_{ST} = 0.0048$; $P = 0.20$; Exact G test $P = 0.06$). Genotype proportions at Cpag6c4b conformed to HWE among both Gulmarsfjord males ($P = 0.5$) and females ($P = 0.8$), with both groups exhibiting nearly identical levels of variability at this locus, as well as at other loci. The locus-specific differentiation between the sexes was effected by a clear shift in respective allele frequency distributions (Fig. 2). The simulation based test for signals of selection within male and female samples identified Cpag6c4b as a positive outlier (Supplementary Fig 1; simulated F_{ST} smaller than Cpag6C4B F_{ST} , $P = 0.9979$) likely to be influenced by divergent selection.

Comparison of the Gulmarsfjord samples with all other samples revealed: (i) the distinctiveness of the Gulmarsfjord males; and (ii) relative similarity of the Gulmarsfjord females to all other samples (Table 3). The pronounced differentiation of the Gulmarsfjord

males from all other samples was also driven by locus Cpag6c4b (mean pairwise F_{ST} for Cpag6c4b = 0.169 (SD = 0.013); mean pairwise F_{ST} excluding Cpag6c4b = 0.003 (SD=0.003)). Outlier identification tests identified Cpag6c4b as a positive outlier in all pairwise and global tests which included the Gulmarsfjord males (Supplementary Fig. 2). Cpag6c4b allele frequency distributions among Gulmarsfjord females were similar to other samples (Fig 2) and pairwise comparisons including this sample yielded a mean pairwise F_{ST} - Cpag6c4b of 0.008 (SD = 0.013), which was similar to values based on the other seven loci (mean pairwise F_{ST} - excluding Cpag6c4b = 0.002 (SD = 0.004)). All outlier tests (pairwise and global) excluding the Gulmarsfjord males reported no significant outliers (Supplementary Fig. 3). Collectively these results indicate that, among the analysed samples, potential divergent selection effects at locus Cpag6c4b were only detectable in comparisons involving the Gulmarsfjord males.

Neutral Genetic structuring

Spatial / Temporal homogeneity among North Sea samples

Among the North Sea samples, excluding the Gulmarsfjord males, all intra-sample pairwise tests of differentiation between sexes were non-significant. Additionally, for those sites with temporally replicated samples, all intra-site comparisons were non-significant regardless of the arrangement of samples (i.e. whether tests were performed on samples pooled or segregated according to sex / time). All pairwise tests between sites yielded non-significant results, regardless of intra-site pooling / partitioning strategies. Upon pooling samples according to site all pairwise tests were non-significant (Table 3), as was global F_{ST} ($F_{ST} = 0.001$; $P = 0.057$). The corresponding multilocus global G test was significant ($P =$

0.02), however this was due to a significant value at only one locus (Cpag3d7, $P = 0.025$), omission of which resulted in a non-significant global G ($P = 0.066$).

Genetic structuring within English Channel and Celtic / Irish Sea samples

Among the English Channel samples no significant pairwise differentiation was detected between sexes in any samples. Significant temporal differentiation was reported between the Hastings 2000 and 2006 samples (Table 3), with this temporal differentiation also evident in pairwise comparisons between the relevant sex-segregated samples (Hast-00 females vs. Hast-06 males $F_{ST} = 0.0055$, $P = 0.045$, Exact G $P = 0.038$; Hast-00 females vs Hast-06 females $F_{ST} = 0.007$, $P = 0.031$, Exact G $P < 0.01$). Significant differentiation between temporal replicates was reported for both within-bay samples from this region (Newlyn and Brittany), as well as for Jersey. Pairwise differentiation between temporal replicate samples within sites in many cases exceeded that between sites and contributed to an overall pattern of low but significant global structuring within the English Channel (global $F_{ST} = 0.004$, $P = 0.001$; global G $P < 0.0001$) which did not show any consistent geographical or temporal pattern (see Table 3). Similar numerically small, yet significant, genetic structuring was also reported among Celtic / Irish Sea samples (global $F_{ST} = 0.004$, $P = 0.001$; global G $P < 0.0001$), driven by the differentiation of, and among, the more southern samples in the region (Table 3).

Inter-regional genetic structure

The two samples from the west of Ireland (NWIRE and GalBay) were not significantly differentiated from each other but exhibited a high proportion of significant pairwise test

results against samples from other regions (Table 3). This differentiation was apparent in the PCoA (Fig.3), which also highlighted the differentiation of the Saint Ives Bay sample (Celtic Sea) with 27 out of 31 significant pairwise tests (Table 3). Examining pairwise test results revealed that differentiation between samples was consistently small but in many cases significant, and did not follow a coherent spatial / temporal pattern but rather similar to the spatial / temporal patchiness reported in the English Channel and Celtic / Irish Seas. This patchiness was not apparent in the STRUCTURE analysis which reported unanimous support for a model of $K = 1$ in all analyses excluding the Gulmarsfjord males. Concordant with the lack of spatial / temporal patterning revealed by pairwise tests no significant IBD was detected (all test results $P > 0.1$) and AMOVA reported greater variation among samples within regions than between regions (Table 4). AMOVA revealed similar patterns among partitioned males and females, and randomisation tests indicated no significant differences between the sexes for a number of indices (Table 5).

DISCUSSION

The present study represents the most geographically extensive investigation of brown crab population genetic structure to date. The research employed fine- and regional-scale spatial / temporal sampling, along with combined gene flow and kinship based analyses and marker neutrality tests to elucidate the mechanistic underpinnings and eco-evolutionary significance of patterns of genetic diversity (following recommendations by Waples 1998, Nielsen et al. 2009, Iacchei et al. 2013). A striking feature of the results was the differentiation of the Gulmarsfjord males from all other samples, including females collected at the same site. This differentiation was driven by a single locus (Cpag6c4b) which was found to have a significantly higher F_{ST} than expected under neutrality in all pairwise

comparisons involving this sample, suggesting divergent selection effects. Excluding the Gulmarsfjord males, all loci (including Cpag6c4b) conformed to neutral expectations and revealed numerically small but statistically significant differentiation among samples across the NE Atlantic. This global genetic structuring did not fit to an IBD model or an obvious hierarchical geographic pattern. Pairwise tests of differentiation (F_{ST} and exact G) revealed that the majority of comparisons were non-significant, including comparisons between geographically distant sites, but that a substantial number of comparisons exhibited significant differentiation which conformed to a model of chaotic genetic patchiness (CGP) in the sense that temporal and / or fine scale differentiation often exceeded that at larger spatial scales (Johnson & Black 1984, Hedgecock 1994, Selkoe et al 2006, Banks et al. 2007). Particular samples (discussed below) were associated with a high proportion of significant pairwise tests indicating geographically local or sample-specific effects. Overall, patterns of neutral genetic variation in brown crab indicate local and unstructured genetic differentiation occurring against a background of high gene flow throughout the studied region.

The positive outlier status of locus Cpag6c4b in all comparisons involving the Gulmarsfjord males suggests that this locus, or a linked genomic region, is subject to divergent selection effects. This result adds to a number of studies reporting selection effects apparent at microsatellite loci that were at some stage assumed to be neutral (Larsson et al. 2007, Skarstein et al. 2007, Westgaard & Fevolden 2007, Nielsen et al. 2009, Gaggiotti et al. 2009, White et al. 2010). Excluding locus Cpag6c4b, the Gulmarsfjord male sample was not significantly differentiated from the Gulmarsfjord female sample, or from most other samples. There are a number of potential explanations for such a pattern of locus-specific genetic differentiation. For example, the pattern could be generated without reproductive isolation through selection on individuals during early life stages followed by random mating each generation (i.e. differential genotype selection within a panmictic gene pool). At the

298 other end of the spectrum the pattern may reflect temporally stable reproductive isolation that
299 is not detectable at neutral loci that lack the statistical power and / or are not at migration-
300 drift equilibrium (Nielsen et al. 2009). Morphological, biochemical and genetic studies have
301 demonstrated population differences among fjords, and between fjords and coastal areas, for
302 a number of taxa (e.g. Jorstad & Naevdal 1989, Suneetha & Naevdal 2001, Oresland &
303 Andre 2008, Teacher et al., 2013). While the allele frequency differences between brown
304 crab sexes within the Gulmarsfjord may indicate gender-specific selection, mechanical
305 mixing of individuals from differently adapted populations could also explain the differences.
306 Evidence of female reproductive migration and lack of return migrations (Ungfors et al.
307 2007) suggests that the sampled Gulmarsfjord females may be allochthonous, while the male
308 sample is composed of local (at least post settlement) individuals. The differentiation
309 between the sexes may therefore reflect allele frequency differences, and by extension
310 adaptive differences, in their respective parental populations and not necessarily differential
311 selection between sexes *per se*. In this sense the pattern could be considered similar to the
312 mechanical mixing of differently adapted migratory north-east Arctic cod (NEAC) and
313 sedentary Norwegian coastal cod (NCC) populations within fjords (Sarvas & Fevolden 2005,
314 Fevolden et al. 2012). The Gulmarsfjord crab data are consistent with other studies indicating
315 that features of fjords may drive local adaptation (Dick et al. 2014) with both salinity and
316 depth highlighted as candidate features by Fevolden et al. (2012), and identified as drivers of
317 selection in other systems (salinity – Nielsen et al 2009; depth – White et al. 2010). Future
318 analysis of larval recruits would provide a means to investigate the relative roles of pre- and /
319 or post settlement selection and dispersal in shaping the observed pattern while identification
320 of underlying functional genetic differences and detection of other samples with similar
321 adaptive fingerprints may help elucidate the environmental drivers.

The seemingly paradoxical pattern of CGP within broad-scale genetic homogeneity reported here has also been documented in a variety of marine species (limpets - Johnson & Black 1984; fish - Planes & Lenfant 2002, Selkoe et al. 2006; barnacles - Veliz et al. 2006). Cautious interpretation of such patterns is recommended as when differentiation is low multiple sources of artificial variance such as unrepresentative sampling (e.g. family / kin sampling - Hansen et al. 1997, Waples 1998, Waples & Gaggiotti 2006) and statistical noise (Waples 1998, Hedrick 1999, 2005) can be important and lead to false conclusions. Kin aggregation is generally assumed to be a transient phenomenon limited to newly settled recruits with little detectable signal in adult populations (Flowers et al. 2002, Planes et al, 2002, Selkoe et al. 2006; but see Iacchei et al. 2013), and as the analysed samples consisted of mixed cohorts of adults kin aggregation would be an unlikely source of error. Furthermore, mean kinship values provided no strong evidence of large proportions of closely related individuals within samples. POWSIM analysis also indicated that the sample sizes conferred low probability of Type I errors. Therefore, while the genetic differentiation only amounts to slight differences in allele frequencies that may not have substantial evolutionary effects (Waples 1998), they nonetheless signal changes in the composition that may be a useful tool for better understanding recruitment dynamics and connectivity in this species (Selkoe et al. 2006, Knutsen et al. 2011).

Fine scale genetic patchiness against a background of high gene flow has been variously attributed to three phenomena that may act in concert: large variances in individual reproductive success (sweepstakes recruitment), limited mixing of larvae from genetically different sources (larval cohesion) and local selection (Larson & Julian 1999). Sweepstakes recruitment has been reported for a number of highly fecund (Type III) marine taxa, such as brown crab, and may generate temporal / spatial genetic differentiation despite gene flow when recruitment is variable. Even in the absence of genetically isolated source populations,

as might be the case here, larval cohesion (Selkoe et al. 2006) may enhance (Waples 2002), and be effectively indistinguishable from sweepstake effects (Turner et al. 2007). The divergent selection effect suggested for Gulmarsfjord males highlights the potential for fine-scale selection in brown crab. However, outlier tests excluding this sample showed no evidence of selection effects for any other locus / sample combination. Furthermore, the temporal differentiation at a number of sites supports more prominent roles for processes like sweepstakes recruitment or larval cohesion as components of recruitment variability, rather than consistent selection effects. For example, the genetic patterns reported for the Hastings samples, the most eastern site sampled in the English Channel, are readily compatible with the proposed relationship between recruitment variability and genetic patchiness. This area was identified *a priori* as a potential hotspot of recruitment variability (Derek Eaton, unpublished study on larval abundance and modelling). As the signatures of such processes are predicted to be diminished by postlarval dispersal (Planes & Lenfant 2002), the genetic patchiness observed here must be considered a conservative reflection of the extent of recruitment heterogeneity.

The CGP in brown crab was unusual in exhibiting a geographic pattern. In contrast to substantial numbers of significant differences among samples within the Irish / Celtic Seas and the English Channel, excluding the Gulmarsfjord males there were no significant differences among North Sea samples (see Table 3), which together with the spatial / temporal genetic homogeneity among Scandinavian samples described by Ungfors et al. (2009) indicates an absence, or lower level, of genetic patchiness in the North Sea compared to other regions studied. While such structuring likely reflects complex interactions between life history and environmental variables, the geographic pattern permits identification of specific factors that may be involved. North Sea brown crabs are significantly smaller than English Channel crabs (Pawson 1995), and as brown crab fecundity is linked to female size

(Edwards 1979; Ungfors 2007) lower fecundity of North Sea crab may reduce the potential extent of reproductive skew compared to those in the English Channel. For example, Palero et al. (2011) suggested that the selected harvesting of large females reduced variance in reproductive success in *Panulirus elephas*. McKeown & Shaw (2008b) posited the genetic monogamy of female brown crab as another life history feature that may increase variance in reproductive success among individuals, but they only analysed samples from the English Channel (which exhibited CGP). Multiple paternity has been reported in a number of closely related species (e.g. Jensen & Bentzen 2012) and may occur in brown crab from other areas, wherein it could serve to reduce variance in reproductive success among males. Seascape factors may directly influence, or interact with the genetic signatures of, variance in reproductive success (Banks et al. 2007). The English Channel exhibits a higher degree of fine scale oceanographic complexity and coastal heterogeneity, in comparison to the North Sea (no CGP), both factors that have been linked to localised sweepstakes recruitment in sea urchins (Banks et al. 2007). Likewise, the high proportion of significant pairwise tests reported for samples collected within semi-enclosed bays (Newlyn, Brittany, Galway, St. Ives), suggests an association between genetic differentiation and habitat structure as an additional component of fine-scale seascape structuring (Selkoe et al. 2010). Comparative studies among taxa with common and contrasting life history strategies will be necessary to elucidate the specific drivers of genetic variation (e.g. Selkoe et al. 2010); however, the brown crab data highlight the fact that genetic structuring may be driven by factors other than dispersal.

Pairwise tests reported a general pattern wherein the West of Ireland samples were differentiated from all samples except those collected in the northern North Sea. This may reflect geographically coherent connectivity. Sotelo et al. (2008) reported spider crab from the west of Ireland to be genetically distinct from more southern samples. However, in

general the genetic patterns for brown crab cannot readily be interpreted in the context of population connectivity/isolation. While the described factors driving CPG may lead to an underestimation of migration rate (m), changes in dispersal behaviour between different life stages may result in broad scale genetic homogeneity masking of spatial gene flow restrictions (e.g. Berry et al. 2012), Partitioned analysis of sexes provided no evidence of greater structuring among male crabs that might be indicative of temporally stable spatial patterns of larval self-recruitment. Furthermore, the difficulties of deriving quantitative estimates of gene flow and dispersal from subtle genetic structure among large populations (Whitlock & McCauley 1999; Palsboll et al. 2007; Hellberg 2009), and discrepancy between levels of gene flow needed to limit genetic differentiation and dispersal needed to replenish stocks (Hauser & Carvalho 2008) are fundamental issues. Therefore, while the low level of genetic structure throughout the studied region is compatible with high gene flow it cannot be ruled out that there is significant isolation of stocks on timescales of interest to management. Resolution of such spatial stock structure may be beyond the level of neutral genetic markers and benefit from complementary analysis of markers under selection (Canino et al. 2005).

This study has implications for sustainable management of the brown crab fishery. The detection of adaptive diversification should enhance appreciation of local adaptation as a component of species biodiversity, and highlights a potential danger of indiscriminate harvesting of differentially adapted units on local scales. Stochastic recruitment variability suggested to underpin genetic patchiness may decrease resilience of local stocks to fishing and increase unpredictability in recovery (Kuparinen et al 2014), and will necessitate a tailoring of the spatial scale of management (spatial bet hedging) according to biological and physical drivers of such recruitment variability. This study provides a baseline for future genetic studies of brown crab, needed to understand recent events such as expansions in census population size within the English channel (Molfese et al. 2014), and further

demonstrates how intraspecific biodiversity and population viability is influenced by complex species-environment interactions other than dispersal.

Acknowledgements: This research was funded by DEFRA. We are indebted to all the fishers and CEFAS personnel who helped in collecting samples, particularly Derek Eaton and Julian Addison (both CEFAS).

LITERATURE CITED

- Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G (2008) LOSITAN: A workbench to detect molecular adaptation based on a F(st)-outlier method. *Bmc Bioinformatics* 9
- Banks SC, Piggott MP, Williamson JE, Bove U, Holbrook NJ, Beheregaray LB (2007) Oceanic variability and coastal topography shape genetic structure in a long-dispersing sea urchin. *Ecology* 88:3055-3064
- Beaumont MA, Nichols RA (1996) Evaluating loci for use in the genetic analysis of population structure. *Proceedings of the Royal Society B-Biological Sciences* 263:1619-1626
- Bennett DB (1995) Factors in the life history of the edible crab (*Cancer pagurus* L.) that influence modelling and management. *ICES Marine Science Symposia* 199:89-98
- Bennett DB, Brown CG (1983) Crab (*Cancer pagurus*) migration in the English Channel. *Journal of the Marine Biological Association of the United Kingdom* 63:371-398
- Berry O, England P, Marriott RJ, BurrIDGE CP, Newman SJ (2012) Understanding age-specific dispersal in fishes through hydrodynamic modelling, genetic simulations and microsatellite DNA analysis. *Molecular Ecology* 21:2145-2159
- Canino MF, O'Reilly PT, Hauser L, Bentzen P (2005) Genetic differentiation in walleye pollock (*Theragra chalcogramma*) in response to selection at the pantophysin (PanI) locus. *Canadian Journal of Fisheries and Aquatic Sciences* 62:2519-2529
- Dick S, Shurin JB, Taylor EB (2014) Replicate divergence between and within sounds in a marine fish: the copper rockfish (*Sebastes caurinus*). *Molecular Ecology* 23:575-590
- Domingues CP, Creer S, Taylor MI, Queiroga H, Carvalho GR (2010) Genetic structure of *Carcinus maenas* within its native range: larval dispersal and oceanographic variability. *Marine Ecology Progress Series* 410:111-123
- Eaton DR, Brown J, Addison JT, Milligan SP, Fernand LJ (2003) Edible crab (*Cancer pagurus*) larvae surveys off the east coast of England: implications for stock structure. *Fisheries Research* 65:191-199
- Edwards E (1979) The edible crab and its fishery in British waters. Fishing News Books, Surrey, England
- ElMousadik A, Petit RJ (1996) High level of genetic differentiation for allelic richness among populations of the argan tree *Argania spinosa* (L) Skeels endemic to Morocco. *Theoretical and Applied Genetics* 92:832-839
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes – application to human mitochondrial DNA restriction data. *Genetics* 131:479-491
- Fevolden SE, Westgaard JI, Pedersen T, Praebel K (2012) Settling-depth vs. genotype and size vs. genotype correlations at the Pan I locus in 0-group Atlantic cod *Gadus*

- morhua*. Marine Ecology Progress Series 468:267-278
- Flowers JM, Schroeter SC, Burton RS (2002) The recruitment sweepstakes has many winners: Genetic evidence from the sea urchin *Strongylocentrotus purpuratus*. Evolution 56:1445-1453
- Gaggiotti OE, Bekkevold D, Jorgensen HBH, Foll M, Carvalho GR, Andre C, Ruzzante DE (2009) Disentangling the effects of evolutionary, demographic, and environmental factors influencing the genetic structure of natural populations: Atlantic herring as a case study. Evolution 63:2939-2951
- Goudet J (1995) FSTAT (Version 1.2): A computer program to calculate F-statistics. Journal of Heredity 86:485-486
- Goudet J, Raymond M, deMeeus T, Rousset F (1996) Testing differentiation in diploid populations. Genetics 144:1933-1940
- Hansen MM, Nielsen EE, Mensberg KLD (1997) The problem of sampling families rather than populations: Relatedness among individuals in samples of juvenile brown trout *Salmo trutta* L. Molecular Ecology 6:469-474
- Hauser L, Carvalho GR (2008) Paradigm shifts in marine fisheries genetics: ugly hypotheses slain by beautiful facts. Fish and Fisheries 9:333-362
- Hedgecock D (1994) Does variance in reproductive success limit effective population sizes of marine organisms. Genetics and Evolution of Aquatic Organisms:122-134
- Hedrick PW (1999) Perspective: Highly variable loci and their interpretation in evolution and conservation. Evolution 53:313-318
- Hedrick PW (2005) A standardized genetic differentiation measure. Evolution 59:1633-1638
- Hellberg ME (2009) Gene Flow and Isolation among Populations of Marine Animals. Annual Review of Ecology Evolution and Systematics 40:291-310
- Hemmer-Hansen J, Nielsen EE, Frydenberg J, Loeschcke V (2007a) Adaptive divergence in a high gene flow environment: Hsc70 variation in the European flounder (*Platichthys flesus* L.). Heredity 99:592-600
- Hemmer-Hansen J, Nielsen EE, Gronkjaer P, Loeschcke V (2007b) Evolutionary mechanisms shaping the genetic population structure of marine fishes; lessons from the European flounder (*Platichthys flesus* L.). Molecular Ecology 16:3104-3118
- Hunter E, Eaton D, Stewart C, Lawler A, Smith MT (2013) Edible Crabs "Go West": Migrations and Incubation Cycle of Cancer pagurus Revealed by Electronic Tags. Plos One 8
- Iacchei M, Ben-Horin T, Selkoe KA, Bird CE, Garcia-Rodriguez FJ, Toonen RJ (2013) Combined analyses of kinship and FST suggest potential drivers of chaotic genetic patchiness in high gene-flow populations. Molecular Ecology 22:3476-3494
- Iles TD, Sinclair M (1982) Atlantic Herring stock discreteness and abundance. Science 215:627-633
- Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. BMC Genetics 6
- Jensen PC, Bentzen P (2012) A molecular dissection of the mating system of the Dungeness crab, *Metacarcinus magister* (Brachyura: Cancridae). Journal of Crustacean Biology 32:443-456
- Johnson MS, Black R (1984) Pattern beneath the chaos – the effect of recruitment on genetic patchiness in an intertidal limpet. Evolution 38:1371-1383
- Jorde PE, Sovik G, Westgaard J-I, Albretsen J, Andre C, Hvingel C, Johansen T, Sandvik AD, Kingsley M, Jorstad KE (2015) Genetically distinct populations of northern shrimp, *Pandalus borealis*, in the North Atlantic: adaptation to different temperatures as an isolation factor. Molecular Ecology 24:1742-1757
- Jorstad KE, Nævdal G (1989) Genetic variation and population structure of Cod, *Gadus morhua* L, in some fjords in northern Norway. Journal of Fish Biology 35:245-252

514 Knutsen H, Olsen EM, Jorde PE, Espeland SH, Andre C, Stenseth NC (2011) Are low but
 515 statistically significant levels of genetic differentiation in marine fishes 'biologically
 516 meaningful'? A case study of coastal Atlantic cod. *Molecular Ecology* 20:768-783
 517 Kuparinen A, Keith DM, Hutchings JA (2014) Increased environmentally driven recruitment
 518 variability decreases resilience to fishing and increases uncertainty of recovery. *ICES*
 519 *Journal of Marine Science* 71:1507-1514
 520 Larson RJ, Julian RM (1999) Spatial and temporal genetic patchiness in marine populations
 521 and their implications for fisheries management. *California Cooperative Oceanic*
 522 *Fisheries Investigations Reports* 40:94-99
 523 Larsson LC, Laikre L, Palm S, Andre C, Carvalho GR, Ryman N (2007) Concordance of
 524 allozyme and microsatellite differentiation in a marine fish, but evidence of selection
 525 at a microsatellite locus. *Molecular Ecology* 16:1135-1147
 526 Latrouite D, Le Foll D (1989) Migrations of the edible crab *Cancer pagurus* and the spider
 527 crab *Maja squinado*. *Oceanis* 15:133-142
 528 McKeown NJ, Arkhipkin AI, Shaw PW (2015a) Integrating genetic and otolith
 529 microchemistry data to understand population structure in the Patagonian Hoki
 530 (*Macruronus magellanicus*). *Fisheries Research* 164:1-7
 531 McKeown NJ, Shaw PW (2008a) Polymorphic nuclear microsatellite loci for studies of
 532 brown crab, *Cancer pagurus* L. *Molecular Ecology Resources* 8:653-655
 533 McKeown NJ, Shaw PW (2008b) Single paternity within broods of the brown crab *Cancer*
 534 *pagurus*: a highly fecund species with long-term sperm storage. *Marine Ecology*
 535 *Progress Series* 368:209-215
 536 Molfese C, Beare D, Hall-Spencer JM (2014) Overfishing and the Replacement of Demersal
 537 Finfish by Shellfish: An Example from the English Channel. *Plos One* 9
 538 Nei M (1978) Estimating of average heterozygosity and genetic distance from a small
 539 number of individuals. *Genetics* 89:583-590
 540 Nielsen EE, Wright PJ, Hemmer-Hansen J, Poulsen NA, Gibb LM, Meldrup D (2009) Micro
 541 geographical population structure of cod *Gadus morhua* in the North Sea and west of
 542 Scotland: the role of sampling loci and individuals. *Marine Ecology Progress Series*
 543 376:213-225
 544 Oresland V, Andre C (2008) Larval group differentiation in Atlantic cod (*Gadus morhua*)
 545 inside and outside the Gullmar Fjord. *Fisheries Research* 90:9-16
 546 Palero F, Abello P, Macpherson E, Beaumont M, Pascual M (2011) Effect of oceanographic
 547 barriers and overfishing on the population genetic structure of the European spiny
 548 lobster (*Palinurus elephas*). *Biological Journal of the Linnean Society* 104:407-418
 549 Palsboll PJ, Berube M, Allendorf FW (2007) Identification of management units using
 550 population genetic data. *Trends in Ecology & Evolution* 22:11-16
 551 Palumbi SR (1994) Genetic divergence, reproductive isolation, and marine speciation.
 552 *Annual Review of Ecology and Systematics* 25:547-572
 553 Pawson MG (1995) Biogeographical identificatoion of English Channel fish and shellfish
 554 stocks. *Fisherie research technical report, MAFF Lowestoft* (66): 72pp.
 555 Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. *Population genetic*
 556 *software for teaching and research. Molecular Ecology Notes* 6:288-295
 557 Planes S, Lecaillon G, Lenfant P, Meekan M (2002) Genetic and demographic variation in
 558 new recruits of *Naso unicornis*. *Journal of Fish Biology* 61:1033-1049
 559 Planes S, Lenfant P (2002) Temporal change in the genetic structure between and within
 560 cohorts of a marine fish, *Diplodus sargus*, induced by a large variance in individual
 561 reproductive success. *Molecular Ecology* 11:1515-1524
 562 Poulsen NA, Hemmer-Hansen J, Loeschcke V, Carvalho GR, Nielsen EE (2011)
 563 Microgeographical population structure and adaptation in Atlantic cod *Gadus*

- morhua*: spatio-temporal insights from gene-associated DNA markers. Marine Ecology Progress Series 436:231-243
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. Genetics 155:945-959
- Queller DC, Goodnight KF (1989) Estimating relatedness using genetic-markers. Evolution 43:258-275
- Raymond M, Rousset F (1995) GENETPOP (version 1.2) – Population genetics software for exact tests and ecumenicism. Journal of Heredity 86:248-249
- Reiss H, Hoarau G, Dickey-Collas M, Wolff WJ (2009) Genetic population structure of marine fish: mismatch between biological and fisheries management units. Fish and Fisheries 10:361-395
- Rousset F (1997) Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. Genetics 145:1219-1228
- Ruzzante DE, Mariani S, Bekkevold D, Andre C, Mosegaard H, Clausen LAW, Dahlgren TG, Hutchinson WF, Hatfield EMC, Torstensen E, Brigham J, Simmonds EJ, Laikre L, Larsson LC, Stet RJM, Ryman N, Carvalho GR (2006) Biocomplexity in a highly migratory pelagic marine fish, Atlantic herring. Journal of Fish Biology 69:236-236
- Ryman N, Palm S (2006) POWSIM: a computer program for assessing statistical power when testing for genetic differentiation. Molecular Ecology Notes 6:600-602
- Ryman N, Utter F, Laikre L (1995) Protection of intraspecific biodiversity of exploited fishes. Reviews in Fish Biology and Fisheries 5:417-446
- Sarvas TH, Fevolden SE (2005) The scnDNA locus Pan I reveals concurrent presence of different populations of Atlantic cod (*Gadus morhua* L.) within a single fjord. Fisheries Research 76:307-316
- Selkoe KA, Gaines SD, Caselle JE, Warner RR (2006) Current shifts and kin aggregation explain genetic patchiness in fish recruits. Ecology 87:3082-3094
- Selkoe KA, Watson JR, White C, Ben Horin T, Iacchei M, Mitarai S, Siegel DA, Gaines SD, Toonen RJ (2010) Taking the chaos out of genetic patchiness: seascape genetics reveals ecological and oceanographic drivers of genetic patterns in three temperate reef species. Molecular Ecology 19:3708-3726
- Shaw PW, Pierce GJ, Boyle PR (1999) Subtle population structuring within a highly vagile marine invertebrate, the veined squid *Loligo forbesi*, demonstrated with microsatellite DNA markers. Molecular Ecology 8:407-417
- Skarstein TH, Westgaard JI, Fevolden SE (2007) Comparing microsatellite variation in north-east Atlantic cod (*Gadus morhua* L.) to genetic structuring as revealed by the pantophysin (Pan I) locus. Journal of Fish Biology 70:271-290
- Sotelo G, Moran P, Fernandez L, Posada D (2008) Genetic variation of the spiny spider crab *Maja brachydactyla* in the northeastern Atlantic. Marine Ecology Progress Series 362:211-223
- Stepien CA, Murphy DJ, Lohner RN, Sepulveda-Villet OJ, Haponski AE (2009) Signatures of vicariance, postglacial dispersal and spawning philopatry: population genetics of the walleye *Sander vitreus*. Molecular Ecology 18:3411-3428
- Suneetha KB, Naevdal G (2001) Genetic and morphological stock structure of the pearlside, *Maurolicus muelleri* (Pisces, Sternoptychidae), among Norwegian fjords and offshore area. Sarsia 86:191-201
- Teacher AGF, Andre C, Jonsson PR, Merila J (2013) Oceanographic connectivity and environmental correlates of genetic structuring in Atlantic herring in the Baltic Sea. Evolutionary Applications 6:549-567
- Therkildsen NO, Hemmer-Hansen J, Als TD, Swain DP, Morgan MJ, Trippel EA, Palumbi SR, Meldrup D, Nielsen EE (2013a) Microevolution in time and space: SNP analysis

- of historical DNA reveals dynamic signatures of selection in Atlantic cod. *Molecular Ecology* 22:2424-2440
- Therkildsen NO, Hemmer-Hansen J, Hedeholm RB, Wisz MS, Pampoulie C, Meldrup D, Bonanomi S, Retzel A, Olsen SM, Nielsen EE (2013b) Spatiotemporal SNP analysis reveals pronounced biocomplexity at the northern range margin of Atlantic cod *Gadus morhua*. *Evolutionary Applications* 6:690-705
- Turner TF, Dowling TE, Marsh PC, Kesner BR, Kelsen AT (2007) Effective size, census size, and genetic monitoring of the endangered razorback sucker, *Xyrauchen texanus*. *Conservation Genetics* 8:417-425
- Ungfors A (2007) Sexual maturity of the edible crab (*Cancer pagurus*) in the Skagerrak and the Kattegat, based on reproductive and morphometric characters. *Ices Journal of Marine Science* 64:318-327
- Ungfors A, Hallback H, Nilsson PG (2007) Movement of adult edible crab (*Cancer pagurus* L.) at the Swedish West Coast by mark-recapture and acoustic tracking. *Fisheries Research* 84:345-357
- Ungfors A, McKeown NJ, Shaw PW, Andre C (2009) Lack of spatial genetic variation in the edible crab (*Cancer pagurus*) in the Kattegat-Skagerrak area. *Ices Journal of Marine Science* 66:462-469
- van Oosterhout C, Weetman D, Hutchinson WF (2006) Estimation and adjustment of microsatellite null alleles in nonequilibrium populations. *Molecular Ecology Notes* 6:255-256
- Veliz D, Duchesne P, Bourget E, Bernatchez L (2006) Genetic evidence for kin aggregation in the intertidal acorn barnacle (*Semibalanus balanoides*). *Molecular Ecology* 15:4193-4202
- Waples RS (1998) Separating the wheat from the chaff: Patterns of genetic differentiation in high gene flow species. *Journal of Heredity* 89:438-450
- Waples RS (2002) Evaluating the effect of stage-specific survivorship on the N-e/N ratio. *Molecular Ecology* 11:1029-1037
- Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology* 15:1419-1439
- Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution* 38:1358-1370
- Weiss M, Thatje S, Heilmayer O, Anger K, Brey T, Keller M (2009) Influence of temperature on the larval development of the edible crab, *Cancer pagurus*. *Journal of the Marine Biological Association of the United Kingdom* 89:753-759
- Westgaard J-I, Fevolden S-E (2007) Atlantic cod (*Gadus morhua* L.) in inner and outer coastal zones of northern Norway display divergent genetic signature at non-neutral loci. *Fisheries Research* 85:306-315
- White TA, Stamford J, Hoelzel AR (2010) Local selection and population structure in a deep-sea fish, the roundnose grenadier (*Coryphaenoides rupestris*). *Molecular Ecology* 19:216-226
- Whitlock MC, McCauley DE (1999) Indirect measures of gene flow and migration: F-ST not equal 1/(4Nm+1). *Heredity* 82:117-125
- Winnepenninckx B, Backeljau T, Dewachter R (1993) Extraction of high molecular weight DNA from molluscs. *Trends in Genetics* 9:407-407

662
663
664
665
666
667
668
669
670
671
672
673
674
675
676

Table 1. Brown crab sample information, including geographical region (used in AMOVA), date of collection and sample composition (i.e. numbers of males/females where identified at time of sampling; * denotes the inclusion of sample in sex-segregated analysis). Multilocus genetic variability measures: N_A (allele number); A_R (allele richness); H_O (observed heterozygosity); H_E (expected heterozygosity); F_{IS} (standardised genetic variance within samples - * denotes significant deviations from HWE expectations); within-sample relatedness (r_{qg} - * denotes values significantly different from expectations of a panmictic model).

Region	Sample site	Sample code	Sample number	Collected	Sample Composition				Mean N_d /locus	Mean A_R	H_O	H_E	F_{IS}	r_{qg}
					Male	Female	Gender unknown	Total						
West of Ireland	Northwest Ireland	NWIre	1	Jul-07			30	30	8.00	7.867	0.601	0.6571	0.086*	-0.013
West of Ireland	Galway Bay	GalBay	2	Jul-07			46	46	8.63	7.567	0.589	0.6456	0.089*	0.029
Irish/Celtic Sea	Southeast Ireland	SEIre	3	Jul-07			31	31	8.38	8.176	0.624	0.646	0.034	0.011
Irish/Celtic Sea	Aberystwyth	Aber	4	Aug-00	8	61*		69	10.00	8.221	0.622	0.630	0.014	0.058*
Irish/Celtic Sea	Newquay	Newq	5	Jun-06	43*	51*		94	10.88	8.196	0.658	0.645	-0.02	0.010
Irish/Celtic Sea	St. Ives Bay	SI Bay	6	Sep-07			55	77	11.13	8.242	0.676	0.649	-0.042	0.008
Irish/Celtic Sea	Pendeen	Pen	7	Jun-06	51*	51*		102	10.75	7.790	0.598	0.652	0.083*	0.004
English Channel	Newlyn Bay	NewBay	8	Sep-00	1	83*		84	11.13	8.357	0.689	0.673	-0.024	-0.019
English Channel	Newlyn Bay	NewBay	8	Oct-07			81	81	10.50	7.983	0.638	0.664	0.041*	0.001
English Channel	Brittany Bay	Brit Bay	9	Sep-00			58	58	9.00	7.696	0.651	0.660	0.014	0.009
English Channel	Brittany Bay	Brit Bay	9	Jul-06		102*		102	11.38	8.366	0.621	0.646	0.042*	0.026
English Channel	Brittany-Offshore	Brit offshore	10	Oct-06	56*	58*		114	11.50	8.221	0.626	0.660	0.053*	-0.003
English Channel	Jersey	Jer	11	Sep-00	40*	32*		72	10.13	7.918	0.662	0.653	-0.014	0.020
English Channel	Jersey	Jer	11	Sep-07			84	84	10.88	8.230	0.653	0.642	-0.018	0.037*
English Channel	Guernsey	Guer	12	Sep-07			80	80	10.50	8.164	0.659	0.659	0	0.000
English Channel	PortScatho	Portscat	13	Jun-06			136	136	11.63	7.915	0.675	0.640	-0.055	0.028
English Channel	Plymouth	Ply	14	Oct-00	9	52*	2	63	10.25	8.129	0.645	0.633	-0.018	0.040
English Channel	Start Point	StartP	15	Jul-06	62*	71*		133	11.75	8.084	0.643	0.649	0.009	0.007
English Channel	Lyme Bay	Lyme	16	Jul-07			52	52	9.88	8.504	0.656	0.645	-0.016	0.027

677

678

679

680 **Table 1.** Continued
681

Region	Sample site	Sample code	Sample number	Collected	Sample Composition				Mean N_a /locus	Mean A_R	H_O	H_E	F_{IS}	r_{qg}
					Male	Female	Gender unknown	Total						
English Channel	Swanage	Swan	17	Jun-06	44*	11		55	9.50	7.97	0.645	0.672	0.041	-0.016
English Channel	Brighton	Brighton	18	Sep-07			65	65	9.63	7.78	0.625	0.648	0.035	0.022
English Channel	Hastings	Hast	19	Aug-00	5	67*		72	10.25	8.23	0.601	0.665	0.097*	-0.015
English Channel	Hastings	Hast	19	Oct-06	54*	108*		162	12.25	8.32	0.628	0.645	0.026	0.019
North Sea	Harwich	Har	20	Jun-00	47*	15	1	63	9.88	8.20	0.688	0.665	-0.033	-0.014
North Sea	Harwich	Har	20	May-05	101*	58*		159	13.13	8.47	0.644	0.656	0.019	0.001
North Sea	Norfolk	Norf	21	Jun-00	39*	39*	2	80	11.13	8.23	0.641	0.652	0.017	0.014
North Sea	Bridlington	Brid	22	Aug-01	44*	40*		84	10.75	8.22	0.629	0.652	0.036	0.010
North Sea	Bridlington	Brid	22	Jun-06	50*	56*		106	11.63	8.15	0.628	0.652	0.036	0.010
North Sea	Northumberland	North	23	Jun-00	48*	46*		94	11.25	8.39	0.659	0.653	-0.011	0.004
North Sea	Northumberland	North	23	Sep-05	56*	9		65	10.50	8.34	0.612	0.647	0.054*	0.014
North Sea	Orkney-Hoy	Ork-Hoy	24	Jun-02	43*	40*	15	98	11.25	8.26	0.649	0.643	-0.008	0.020
North Sea	Orkney-Sanday	Ork-Sand	25	Jun-02			38	38	9.25	8.58	0.686	0.682	-0.006	-0.041
North Sea	Shetland	Shet	26	Jun-07				48	9.63	8.09	0.672	0.665	-0.010	-0.017
North Sea	Gulmarsfjord	Gulm	27	Jun-02	41**	39*		80	10.00	7.82	0.627	0.651	0.037	0.014

682
683
684
685

Table 2. Estimated statistical power for detecting various true levels of population differentiation (F_{ST}) by means of Fisher's exact G tests in pairwise comparisons involving various permutations of sample sizes relative to this study ($n = 30$: minimum sample size used in pairwise tests; $n = 84$: average sample size; $n = 120$: representative of larger sample sizes employed). Power is expressed as the proportion of simulations reporting statistical significance at the 0.05 level. Bold values denote values obtained for simulated $F_{ST} = 0$ (Type I error), non-bold denote values obtained for simulated $F_{ST} = 0.0025$.

	sample $n = 30$	sample $n = 84$	sample $n = 120$
sample $n = 30$	0.049 /0.409		
sample $n = 84$	0.065 /0.699	0.047 /0.981	
sample $n = 120$	0.055 /0.784	0.048 /0.997	0.062 /1

Table 3. Pairwise F_{ST} values between all samples, with intraregional comparisons outlined and shaded. See Table 1 for sample codes. PF_{ST} and Exact G tests yielded similar patterns of significance and so only PF_{ST} are reported (italics = $P < 0.05$; underlined = $P < 0.01$; bold = $P < 0.001$).

	NWIRE	GalBay	SEIre	Aber	Newq	SI Bay	Pen	NewBay-2000	NewBay-2007	Brit Bay -2000	Brit Bay-2006	Brit offshore	Jer-2000	Jer-2007
GalBay	0.003													
SEIre	0.003	0.019												
Aber	<i>0.007</i>	0.018	0.006											
Newq	0.004	0.013	0.005	-0.005										
SI Bay	0.004	0.022	0.015	<i>0.005</i>	<u>0.005</u>									
Pen	0.013	0.024	<u>0.010</u>	<i>0.005</i>	<i>0.004</i>	<i>0.004</i>								
NewBay-2000	<u>0.010</u>	0.015	<u>0.011</u>	<u>0.007</u>	<i>0.004</i>	0.011	0.002							
NewBay-2007	<u>0.011</u>	0.017	0.021	0.013	<u>0.007</u>	0.012	<u>0.007</u>	<u>0.006</u>						
Brit Bay -2000	0.019	0.022	0.017	<u>0.008</u>	<u>0.007</u>	0.018	<u>0.008</u>	<u>0.009</u>	0.013					
Brit Bay-2006	0.003	0.014	0.002	0.001	-0.001	0.003	0.002	<u>0.005</u>	<u>0.008</u>	0.013				
Brit offshore	<i>0.008</i>	0.013	<u>0.011</u>	<i>0.004</i>	0.001	<u>0.005</u>	0.002	0.001	0.002	0.002	<i>0.003</i>			
Jer-2000	<u>0.013</u>	0.025	0.007	0	0.002	<u>0.008</u>	0.001	0.001	0.008	0.004	0.002	0.001		
Jer-2007	<i>0.008</i>	<u>0.009</u>	<u>0.012</u>	<u>0.007</u>	0.002	<u>0.008</u>	<i>0.004</i>	<u>0.006</u>	<i>0.005</i>	<u>0.009</u>	0.003	0.002	<u>0.008</u>	
Guer	<i>0.008</i>	0.011	<u>0.012</u>	0.003	0.001	<i>0.006</i>	0.001	-0.001	<i>0.005</i>	<u>0.009</u>	0.001	0.001	0.002	0.003
Portscat	<i>0.006</i>	0.012	<u>0.012</u>	0.001	-0.002	<i>0.004</i>	<i>0.003</i>	0.002	0.003	0.011	-0.001	0.001	0.001	0.003
Ply	0.023	0.019	<u>0.013</u>	<i>0.007</i>	<u>0.006</u>	0.019	<u>0.007</u>	0.001	<u>0.008</u>	<u>0.008</u>	<u>0.007</u>	<i>0.004</i>	0.004	<i>0.006</i>
StartP	<u>0.012</u>	0.018	<i>0.009</i>	0.002	0.001	<u>0.006</u>	0	0.002	<u>0.008</u>	0.002	<i>0.004</i>	-0.001	0.001	<i>0.002</i>
Lyme	0.007	0.020	<i>0.011</i>	-0.001	0.003	0.004	<i>0.007</i>	<i>0.007</i>	<u>0.014</u>	0.014	0.002	<i>0.005</i>	0.003	0.014
Swan	0.007	<i>0.008</i>	<i>0.008</i>	0.015	<i>0.007</i>	<u>0.011</u>	0.002	<i>0.006</i>	<i>0.007</i>	<u>0.009</u>	<i>0.006</i>	0.002	<u>0.009</u>	0.002
Brighton	<i>0.009</i>	<u>0.010</u>	<u>0.013</u>	0.003	0.003	<u>0.012</u>	0.005	0.001	0.003	<u>0.010</u>	<i>0.004</i>	0.002	0.003	0.004
Hast-2000	<u>0.017</u>	<u>0.017</u>	<u>0.016</u>	0.011	0.008	0.022	0.011	0.003	0.012	0.003	0.013	<i>0.005</i>	<u>0.007</u>	<u>0.007</u>
Hast-2006	<i>0.006</i>	<u>0.013</u>	<u>0.012</u>	0.002	0	0.003	0	0.003	0.006	0.008	0	0.001	0.003	0.002
Har-pooled	<i>0.006</i>	0.011	<i>0.008</i>	0.002	0	0.007	0.002	0.001	0.007	<u>0.007</u>	<i>0.002</i>	0.001	0.002	0.004
Norf	0.018	0.017	<u>0.011</u>	<i>0.009</i>	<u>0.008</u>	0.022	0.008	0.002	0.016	<i>0.005</i>	0.014	<u>0.006</u>	0.003	0.011
Brid-pooled	<u>0.011</u>	<u>0.009</u>	<u>0.014</u>	<i>0.004</i>	<u>0.005</u>	0.014	0.009	<i>0.003</i>	0.012	<u>0.007</u>	<u>0.007</u>	<i>0.003</i>	<u>0.005</u>	<u>0.008</u>
North-pooled	<u>0.011</u>	0.017	<u>0.009</u>	0.003	0.002	0.014	0.002	0.001	0.011	<i>0.004</i>	<u>0.005</u>	0.001	0.001	<u>0.005</u>
Ork-Hoy	<u>0.012</u>	0.015	<i>0.008</i>	0.003	<i>0.004</i>	0.012	0.003	0.002	0.012	<i>0.005</i>	<i>0.003</i>	<i>0.003</i>	0.001	<u>0.005</u>
Ork-Sand	0.001	0.006	0.009	0.005	0	<i>0.006</i>	0.002	-0.002	-0.001	0.013	0	-0.001	0.004	0.003
Shet	0.007	<u>0.013</u>	<i>0.008</i>	0.004	-0.007	<i>0.007</i>	0.002	-0.001	0.003	0.003	0.003	-0.003	0.002	0.003
Gulm-females	0.003	0.004	0.005	0.004	0.002	<u>0.012</u>	0.005	-0.001	<i>0.008</i>	<i>0.010</i>	0.002	0.004	<i>0.007</i>	0.002
Gulm-males	0.028	0.035	0.037	0.028	0.026	0.036	0.030	0.027	0.036	0.029	0.030	0.025	0.032	0.027

727 **Table 3** continued
728

	Guer	Portscat	Ply	StartP	Lyme	Swan	Brighton	Hast-2000	Hast-2006	Har-pooled	Norf	Brid-pooled	North-pooled
Guer													
Portscat	-0.001												
Ply	0.002	<i>0.004</i>											
StartP	0.001	<i>0.002</i>	<i>0.004</i>										
Lyme	0.003	0.001	<u>0.011</u>	<i>0.007</i>									
Swan	0.004	<u>0.008</u>	0.012	<i>0.004</i>	0.016								
Brighton	-0.001	-0.003	0	<i>0.004</i>	<i>0.007</i>	<u>0.010</u>							
Hast-2000	<i>0.004</i>	<u>0.009</u>	0.002	0.003	0.015	<u>0.009</u>	0.003						
Hast-2006	0	-0.005	0.005	0.001	0.003	0.005	0.003	0.009					
Har-pooled	-0.001	0.001	<i>0.004</i>	0.001	0.003	<i>0.004</i>	0.002	0.007	0				
Norf	0.003	<u>0.008</u>	0	<u>0.005</u>	<u>0.009</u>	0.011	<i>0.006</i>	0.004	0.008	0.004			
Brid-pooled	0.002	<u>0.003</u>	<u>0.005</u>	<u>0.003</u>	<i>0.005</i>	<u>0.009</u>	0.003	<i>0.004</i>	0.005	0.001	0.003		
North-pooled	0.001	<i>0.003</i>	0.003	0	<i>0.005</i>	<u>0.005</u>	0.003	<u>0.005</u>	0.001	-0.001	0.003	0	
Ork-Hoy	0.001	<i>0.003</i>	0.001	0.001	<i>0.006</i>	<i>0.005</i>	0.003	<i>0.005</i>	0.002	0	0.001	0.001	-0.002
Ork-Sand	-0.002	-0.002	0.005	0.002	0.004	0.001	-0.002	0.005	-0.001	-0.001	0.004	0.002	0.001
Shet	-0.002	0	0.001	-0.001	0.005	0.004	-0.002	-0.002	0.002	-0.001	0.004	0	0
Gulm-females	-0.001	0.001	0.001	0.002	<i>0.009</i>	0.004	-0.004	0	0.002	0	0.005	0.001	0.002
Gulm-males	0.024	0.028	0.035	0.025	0.032	0.032	0.025	0.029	0.026	0.027	0.030	0.026	0.026

Table 3. Continued

	Ork-Hoy	Ork-Sand	Shet	Gulm-females
Ork-Hoy				
Ork-Sand	0.003			
Shet	0.002	-0.001		
Gulm-females	0.001	-0.005	-0.003	
Gulm-males	0.026	0.027	0.024	0.027

Table 4. Analysis of molecular variance (AMOVA) across the three main sampling regions (Celtic/Irish Sea; English Channel; North Sea), using either all data or sex segregated data (see Table 1), and temporal samples within sites pooled or unpooled. F_{CT} = genetic variance among regions; F_{SC} = genetic variance among samples within regions.

		F_{CT}	F_{SC}
All individuals	Temporal samples unpooled	0.001 ($P < 0.001$)	0.003 ($P < 0.001$)
	Temporal samples pooled	0.001 ($P < 0.001$)	0.002 ($P < 0.001$)
Males only	Temporal samples unpooled	0.002 ($P = 0.010$)	0.002 ($P = 0.040$)
	Temporal samples pooled	0.002 ($P < 0.001$)	0.002 ($P = 0.019$)
Females only	Temporal samples unpooled	0.001 ($P = 0.059$)	0.003 ($P < 0.001$)
	Temporal samples pooled	0.001 ($P = 0.018$)	0.0018 ($P = 0.012$)

Table 5. Comparative analysis of genetic diversity indices for brown crab females and males, and corresponding P values from 2-tailed tests.

	Female	Male	Two tailed P
A_R	7.709	7.755	0.674
H_o	0.643	0.632	0.330
F_{IS}	0.012	0.032	0.186
F_{ST}	0.003	0.003	0.638
R_{gg}	0.007	0.005	0.611

Figure 1. Brown crab sample sites (see Table 1 for site details). Red discs denote sites with temporal replicates, while underlined numbers highlight sample sites within semi-enclosed bays and fjords.

755

756

757 **Figure 2.** Allele frequencies at microsatellite locus Cpag6c4B for Gulmarsfjord males,
758 Gulmarsfjord females, and for all other ‘non-fjord’ samples pooled.

759

760

761

762

763

764

765

766

767

768

769

770

771

772

Figure 3. Principal coordinates analysis of multi-locus pairwise F_{ST} .