Aberystwyth University

Abrupt or gradual?

Trauth, Martin; Foerster, Verena; Junginger, Annett; Asrat, Asfawossen; Lamb, Henry; Schaebitz, Frank

Published in:
Quaternary Research

DOI:
[10.1017/qua.2018.30](https://doi.org/10.1017/qua.2018.30)

Publication date:
2018

Citation for published version (APA):

Trauth, M., Foerster, V., Junginger, A., Asrat, A., Lamb, H., & Schaebitz, F. (2018). Abrupt or gradual? Change point analysis of the late Pleistocene–Holocene climate record from Chew Bahir, southern Ethiopia. *Quaternary Research*, 90(2), 321-330. <https://doi.org/10.1017/qua.2018.30>

Document License CC BY-NC-ND

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

1 Abrupt or Gradual? Change Point Analysis of the Late Pleistocene- 2 Holocene Chew Bahir Record from Southern Ethiopia

3 Martin H. Trauth^{*1}, Verena Foerster², Annett Junginger³, Asfawossen Asrat⁴, Henry F. Lamb⁵, Frank
4 Schaebitz²

5 ¹Institute of Earth and Environmental Science, University of Potsdam, Karl-Liebknecht-Str. 24-25, 14476
6 Potsdam, Germany

7 ²Institute of Geography Education, University of Cologne, Gronewaldstraße 2, 50931 Köln, Germany

8 ³Senckenberg Center for Human Evolution and Palaeoenvironment (HEP), Department of Geosciences,
9 University of Tübingen, Hölderlinstrasse 12, 72074 Tübingen, Germany

10 ⁴Addis Ababa University, School of Earth Sciences; P. O. Box 1176, Addis Ababa, Ethiopia

11 ⁵Aberystwyth University, Department of Geography and Earth Sciences, Aberystwyth SY23 3DB, UK

12 *Corresponding author at Institute of Earth and Environmental Science, University of Potsdam, Karl-
13 Liebknecht-Str. 24-25, 14476 Potsdam, Germany. E-mail address: trauth@uni-potsdam.de (M.H. Trauth).
14 Telephone: +49 331 977 5810.

15

16 **ABSTRACT**

17 We have used a change point analysis on a Late Pleistocene-Holocene lake-sediment record from the Chew
18 Bahir basin in the southern Ethiopian Rift to determine the amplitude and duration of past climate transitions,
19 and then assessed their possible influence on the development of early modern human cultures. The most
20 dramatic changes occurred over 240 years (from ~15,700 to 15,460) during the onset of the African Humid
21 Period (AHP), and over 990 years (from ~4,875 to 3,885 a BP) during its protracted termination. The AHP
22 was interrupted by a distinct dry period coinciding with the high-latitude Younger Dryas (YD) stadial, which
23 had an abrupt onset (less than ~100 years) at ~13,260 a BP and lasted until ~11,730 a BP. Wet-dry-wet
24 transitions prior to the AHP may reflect the high-latitude Dansgaard-Oeschger cycles, as indicated by cross-
25 correlation of the K record with the NorthGRIP ice core record between ~45–20 ka BP. These findings
26 provide a valuable contribution to the debate regarding the amplitude and duration of past climate transitions
27 and their possible influence on the development of early modern human cultures.

28 **Keywords:** Late Pleistocene; Holocene; Change point analysis; Principal component analysis;
29 Paleoclimatology; Southern Ethiopian Rift; African Humid Period; Younger Dryas; Dansgaard-Oeschger
30 cycles

31

INTRODUCTION

Determining the nature and pace of changes in the environment of early modern humans is crucial to understanding the factors that influenced technological, behavioural and cognitive innovation: different modes of climate variability would have caused different modes of climatic stress and environmental boundaries (Hildebrand and Grillo, 2012; Vogelsang and Keding, 2013; Foerster et al., 2015, 2016). The timing and synchronicity of the dry-wet-dry oscillations that have significantly modulated the climate across eastern Africa are of particular interest, with important implications for human adaptation and survival in (e.g. highland) refugia (Ambrose et al., 1998; Hildebrand et al., 2010; Foerster et al., 2015) and for human dispersal, both within Africa and beyond (Carto et al., 2009; Castañeda et al., 2009).

Testing hypotheses concerning the relationship between climate and human cultural change is complicated by the complex interplays between climate, environment, and human behaviour, and relies largely on archaeological datasets and modelling (e.g., Widlok et al., 2012). The reconstruction of the climatic component on which we have focused can be compromised by uncertainties in the interpretation of climate proxies, the dating of climate archives, and the validation of modelling results. For example, while consensus prevails on the causes and nature of the African Humid Period (~14,800–5,500 a BP; deMenocal et al., 2000), the precise timing of this interval and the relative abruptness of its onset and termination in different regions remain a matter of debate (e.g., Kröpelin et al., 2008a, b; Brovkin and Claussen, 2008; Junginger and Trauth, 2013; Tierney and deMenocal, 2013; Shanahan et al., 2015; Tierney et al., 2017). Furthermore, evaluating the response of the tropical and subtropical African climate to North Atlantic cold events such as the Heinrich Events and the Younger Dryas stadial has proved controversial, with a general acceptance that the Younger Dryas had a drying effect on the African climate but with different interpretations of its timing (Roberts et al., 1993; Garcin et al., 2007; Tierney and deMenocal, 2013; Berke et al., 2014). The impacts of the Dansgaard-Oeschger (D-O) cycles and the Heinrich events of the northern hemisphere high latitudes on East African moisture availability are also subject to debate (e.g. Brown et al., 2007; Lamb et al., 2007; Garcin, 2008; Tierney and deMenocal, 2013).

57 In this paper we present the results of a change point analysis on high-resolution environmental records from
58 the Chew Bahir, which we used to determine the abruptness of climatic transitions. We then discuss the
59 possible impact of such transitions on human cultural evolution. The Chew Bahir is a 30 x 70 km saline
60 mudflat within a tectonically bounded basin, located in a transition zone between the Main Ethiopian Rift and
61 the Omo-Turkana basin (Fig. 1). In the Omo-Turkana basin the fossils of one of the oldest (~196 ka,
62 McDougall et al., 2005) known anatomically modern humans have been found, although recent discoveries
63 in Jebel Irhoud, Morocco, suggest that our species actually appears much earlier (~315 ka, Richter et al.,
64 2017; Hublin et al., 2017). The south-western Ethiopian Highlands, which are within the 32,400 km² Chew
65 Bahir catchment, may have been a refugium for groups of *H. sapiens* during times of climatic stress
66 (Ambrose et al., 1998; Brandt and Hildebrand, 2005) as well as a possible hotspot of cultural innovation
67 (Brandt et al., 2012). The several km thick deposits of the Chew Bahir basin therefore present a promising
68 archive of past environmental changes, containing clues to their possible influence on human expansion and
69 cultural innovation (Campisano et al., 2017).

70 Our environmental record is based on a comprehensive multi-proxy analysis of six sediment cores up to 18.8
71 m long, collected along an approximately 20 km long NW-SE transect across the western half of the basin
72 (Fig. 1). Three of these cores (CB-01, CB-03 and CB-05), which were collected in a pilot study for the
73 Hominin Sites and Paleolakes Drilling Project (HSPDP; HSPDP-CHB deep coring site in Fig.1) (Cohen et al.,
74 2016; Campisano et al., 2017) and the Collaborative Research Center CRC-806, have been described in
75 previous publications (Foerster et al., 2012, 2015; Trauth et al., 2015). By incorporating new data from CB-
76 02, CB-04 and CB-06 we can make a definitive contribution to the discussion on the timing of the onset and
77 termination of the AHP, the YD, and earlier dry-wet-dry cycles, as well as the rates of change involved. As in
78 our previous publications, we use potassium concentrations (standardized counts of XRF scans) as a proxy
79 for aridity in the Chew Bahir (Foerster et al., 2012, 2015; Trauth et al., 2015). The potassium record spans
80 the last 45,280 years, including the last two orbitally-driven dry-wet-dry climate alternations in north-east
81 Africa. A dynamic time warping algorithm was used to compare and correlate proxy time series from the six
82 sediment cores, and a principal component analysis to unmix the records and help us to separate regional
83 from local influences. Since this continuous high-resolution record from the 32,400 km² large Chew Bahir

84 catchment can be considered to be representative of the whole of the southern Ethiopian Plateau region, it is
85 a promising dataset against which to test hypotheses concerning human responses to climatic changes in
86 north-east Africa over the last ~45 ka.

87 REGIONAL SETTING

88 Chew Bahir (4.1–6.3°N, 36.5–38.1°E; WGS 84; [Fig. 1](#)) is a closed basin, separated from the Turkana Basin
89 to the west by the Hammar Range. It therefore forms a terminal sink for weathering products from its 32,400
90 km² catchment. The perennial Weyto and Segen rivers drain from the north/north-west and north-east,
91 respectively, to feed the playa lake. Large alluvial fans reach into the Chew Bahir basin from the flanks of the
92 southern Ethiopian Rift and are activated during arid intervals, when the vegetation cover is sparse ([Foerster
93 et al., 2012](#)). The Hammar Range to the west and the highlands to the north and north-east consist of Late
94 Proterozoic granitic and mafic gneisses with minor occurrences of meta-sedimentary rocks, whereas the
95 eastern part of the catchment is dominated by Miocene basaltic lava flows with subordinate rhyolite-trachyte
96 and felsic tuff intercalations. Oligocene basalt flows with subordinate rhyolites, trachytes, tuffs and
97 ignimbrites cover the Precambrian basement units in the distal north-eastern, northern, and north-western
98 parts of the catchment ([Moore and Davidson, 1978; Davidson, 1983; Fig. 1](#)).

99 The present-day climate in east and northeast Africa is influenced by a number of major air streams and
100 convergence zones, with their effects superimposed on regional influences associated with topography,
101 large lakes, and the oceans ([Nicholson, 2017](#)). According to the classic explanation of the seasonal cycle
102 rainfall in the Chew Bahir catchment is associated with the passage of the ITCZ, resulting in a strongly
103 bimodal annual cycle. The two resulting rainy seasons follow the latitudinal position of the overhead sun with
104 a time lag of about 4-6 weeks; the "long rains" occur between March and May and the "short rains" between
105 October and November. Short-term (annual to decadal) fluctuations in the intensity of precipitation relate to
106 E-W adjustments in the zonal Walker circulation associated with the El Niño-Southern Oscillation (ENSO)
107 and the Indian Ocean Dipole (IOD), possibly as a direct response to sea-surface temperature (SST)
108 variations in the Indian and Atlantic Oceans, which are in turn affected by the ENSO and the IOD. Changes

109 in the relative strength of these Pacific and Indian Ocean anomalies may also explain the continuous decline
110 of the intensity of the long rains in recent decades (Nicholson, 2017).

111 MATERIALS AND METHODS

112 The Late Pleistocene-Holocene dry-wet-dry oscillations in the Chew Bahir basin have been reconstructed
113 from the six sediment cores (CB01–06) collected along a ~20 km long NW–SE transect across part of the
114 basin (Foerster et al., 2012, 2015) (Table 1, Figs. 1 and 2). The CB01 and CB02 cores were retrieved from
115 the distal margins of an alluvial fan on the western edge of the basin, an area that is predominantly
116 influenced by the fans from the Hammar Range and the Weyto River, while the CB04 to CB06 cores were
117 collected from towards the center of the basin, in an area predominantly controlled by fluvio-lacustrine
118 processes and the Segen River. The sediments in the CB03 core, which was from an intermediate location
119 along the transect, were of mixed alluvial, lacustrine and fluvial origins with the latter relating to the Weyto
120 and Segen rivers.

121 The climate proxy record from the Chew Bahir basin discussed herein is based on the abundance of
122 potassium (K) in the sediment, as determined by micro X-ray fluorescence (μ XRF); potassium has previously
123 been shown to be a reliable proxy for aridity in the Chew Bahir cores (Foerster et al., 2012, 2014; Trauth et
124 al., 2015). We used dynamic time warping (DTW) to automatically align the K records from cores CB02 to
125 CB06 with the K record from core CB01 (Figs. 2 and 3). Dynamic time warping, included in MATLAB function
126 as the `dtw` function (Mathworks, 2017a), stretches two time series onto a common set of points in time such
127 that the sum of the Euclidean distances between corresponding points is as small as possible (Sakoe and
128 Chiba, 1978; Paliwal et al., 1982). The core depths of the aligned cores CB01 to CB06 were converted into
129 ages using the age model of Trauth et al. (2015) (Fig. 3)

130 We used a principal component analysis (PCA), included in the MATLAB function `pca` (Mathworks, 2017b),
131 to unmix (or separate) the mixed regional and local environmental signals in cores CB01 to CB06 (Pearson,
132 1901; Hotelling, 1931) (Fig. 4). More than 94% of the total variance of the data set is contained in the first
133 principal component (PC1), which we interpret to represent the regional climate signal. However, complete

134 unmixing of the regional and site-specific signal components (including noise) was not possible since the K
135 values from the CB01 to CB06 cores are not perfectly Gaussian distributed (Fig. 4). The temporal resolution
136 of the climate proxy record in CB01 has a mean of approximately 16 years, ranging from ~4 years up to
137 almost 2,000 years between about 41 and 42 ka BP. The K record after DTW alignment and PCA-based
138 unmixing was interpolated to an evenly spaced time axis running from 0 to 45,350 a BP at 15 year intervals,
139 which is close to the mean interval in the original data (~16 years).

140 In order to test the synchronicity of the African Humid Period across tropical and subtropical Africa and the
141 relative abruptness of its onset and termination, we applied a change point search algorithm from Killick et al.
142 (2012) to the standardized K record, i.e. the mean K values were subtracted from the individual K values and
143 then divided by the standard deviation. The algorithm, which has recently been included in MATLAB as the
144 `findchangepts` function (MathWorks, 2017a), detects change points by minimizing a cost function over all
145 possible numbers and locations of change points. The `findchangepts` function yields the number of
146 significant changes in the mean, the standard deviation, and the trend of a time series (not exceeding a
147 maximum number of permissible changes defined by the user) that minimize the sum of the residual error
148 and an internal fixed penalty for each change.

149 In order to test the Chew Bahir record for any influence from the pre-AHP centennial to millennial climate
150 oscillations of D-O cycles and Heinrich events we calculated the correlation coefficient between the K record
151 and the NorthGRIP (also known as NGRIP) $\delta^{18}\text{O}$ ice core record (North Greenland Ice Core Project
152 members, 2004). We then also calculated the correlation coefficient within a sliding window of a given length
153 covering the two records. We used Pearson's linear product-moment correlation coefficient and, since this
154 coefficient is very sensitive to disturbances, we also used the Spearman and Kendall correlation coefficients,
155 all implemented in the MATLAB function `corr` (MathWorks, 2017b).

156 RESULTS

157 The micro X-ray fluorescence (μXRF) scans provided data on the elemental content of six cores CB01 to
158 CB06 (for 26 to 47 elements, depending on the setting of the core scanner) (Figs. 2 to 4). Of all the elements

159 analyzed we only made statistical analyses of the K records, as these have been shown to reflect past
160 climate variability on millennial to centennial time scales (Foerster et al., 2012, 2015; Trauth et al., 2015)
161 (Figs. 2 to 4). After DTW alignment and PCA-based unmixing the application of the change point search
162 algorithm to the data means yielded about 15 significant transitions during the last ~45 ka (Fig. 5a).
163 Increasing the maximum number of changes permitted in `findchangepts` yielded additional transitions
164 bounding narrow minima (i.e. short wet episodes, based on interpretation of the K record) to the series of
165 change points. Reducing the maximum number of changes to 10, however, resulted in the gradual increase
166 in K concentrations at the end of the AHP being defined by just a single change point (at 4,860 a BP).
167 Adjusting the maximum number of change points to an intermediate value placed the termination of the AHP
168 between ~4,875 to 3,885 a BP, as suggested by two significant change points, one at the beginning of this
169 climate transition and one at the end. The transition from wet to dry at the end of the AHP therefore appears
170 to have lasted ~990 years in the Chew Bahir basin.

171 A second set of important transitions is the one defining the episode of higher K concentrations that
172 coincides with the Younger Dryas stade. According to our analysis the onset of this ~1,500 year long dry
173 episode in the Chew Bahir basin was at 13,260 a BP and it lasted until 11,730 a BP. Prior to the YD there
174 was a transition from dry to wet evident in the CB01–CB04 and CB06 cores, but not in the central core
175 (CB05). This very abrupt transition is dated at 15,640 a BP, most likely marking the onset of the AHP in the
176 Chew Bahir basin. Earlier transitions from an even drier to a wetter climate (as indicated by changes from
177 high to low K concentrations) occur at 19,860 and 20,520 a BP. In the older parts (i.e. prior to ~20 ka BP) of
178 the cores, there are numerous rapid shifts from dry (high K) to wet (low K) and back within less than a
179 thousand years; the pattern of these rapid transitions resembles the D-O cycles of high latitudes, as
180 discussed below.

181 Many of these transitions in the mean K concentration are also associated with changes in the variability of K
182 concentrations, as indicated by the change point search in the standard deviation (Fig. 5b). For instance, the
183 change point algorithm finds a step-wise decrease in variability at 4,875 a BP, and again at 3,870 a BP, most
184 likely caused by an episode of relative instability that is characterized by a relative wet climate between

185 ~7.15 and ~5 ka BP interrupted by at least fourteen distinct dry events, each lasting between ~20 and 80
186 years (Figs. 2 to 4); this episode is followed by a relatively dry and stable climate after ~5 ka BP (Fig. 3a in
187 Trauth et al., 2015). The Younger Dryas dry period is also bounded by two transitions in the standard
188 deviation (from wet to dry and then back again), at 13,260 and 11,730 a BP, which are the same dates as
189 the transitions in the mean. The onset of the AHP, as defined by a change in the mean K concentration at
190 15,640 a BP, is also associated with a change point in the standard deviation at about the same date.
191 Similarly, the older step-wise dry-wet transitions in the mean K concentration are associated with changes in
192 the standard deviation at more or less the same times.

193 The change point search algorithm also allows us to analyse changes in the trend of the K concentration
194 (Fig. 5c). It is interesting to note that the termination of the AHP took much longer if one considers the trend,
195 extending from 5,835 to 3,840 a BP and therefore including a longer period of the episode of instability
196 detected from the mean K concentrations, which lasted from ~7.15 to 3.8 ka BP. The trend analysis
197 suggests that the wet-dry-wet episode corresponding to the Younger Dryas started at 13,260 a BP and
198 ended at 11,730 a BP, which is very close to the dates obtained from analysis of the mean K concentrations.
199 According to the change point in the trend the onset of the African Humid Period was at 15,640 a BP, which
200 is exactly the same as the date obtained from the mean and standard deviation analyses. The search for
201 change points in the trend of the older part of the K record reveals numerous sawtooth-type intervals,
202 terminated by significant change points, with the earlier ones defining short periods with lower K
203 concentrations (i.e. wetter conditions).

204 Correlating the entire K record from CB01 with the $\delta^{18}\text{O}$ record from the NorthGRIP ice core (North
205 Greenland Ice Core Project members, 2004) yields a Pearson correlation coefficient of -0.61 , which is
206 similar to the Spearman correlation coefficient (-0.60) but differs from the Kendall correlation coefficient ($-$
207 0.42), i.e. high negative $\delta^{18}\text{O}$ values correlate with high positive K values (Fig. 6). These results suggest a
208 weak but consistently negative correlation, i.e., cold climates in the high northern latitudes correlate with (but
209 do not necessarily cause) dry climates in the Chew Bahir basin. Piecewise cross correlation using the
210 Spearman coefficient with a window size of 100 data points (corresponding to $100 \times 15 \text{ a} = 1,500 \text{ a}$) reveals

211 that this negative correlation is not stationary. In fact periods of negative and positive correlation alternate
212 with each other, interrupted at times by periods with no correlation. It is interesting to note that episodes with
213 a correlation below -0.5 occur prior to ~ 25 ka BP, briefly at about 20 ka BP, and between 15 and 10 ka BP,
214 i.e. at the same times as D-O cycles and the YD stadial in Greenland, whereas the correlation is positive or
215 absent (values of $0-0.5$) during relatively wet climatic episodes. Qualitatively similar results were obtained
216 from correlation analyses using Pearson's and Kendall's correlation coefficients instead of the Spearman's
217 coefficient.

218 **DISCUSSION AND CONCLUSIONS**

219 *Climatic transitions in the Chew Bahir record*

220 The potassium record from the Chew Bahir basin provides one of the most complete and detailed records of
221 eastern African climatic variations, covering two full precessional cycles (the last ~ 45 ka) (Figs. 5 and 6). The
222 six cores (CB01–06) generally recorded the same climate signals but with different intensities (Figs. 2 to 4).
223 The CB01–03 cores were collected at increasing distances from the Hammar Range, which is the main
224 sediment source, while the CB04–06 cores were collected towards the centre of the basin. The centre of the
225 Chew Bahir basin is influenced to some extent by the Segen River, whereas the more western sites are
226 increasingly influenced by runoff from the Hammar Range via the extensive alluvial fans, and episodically by
227 the Weyto River. As a result there are significant differences in the type of sediment, the rate of
228 sedimentation and, most importantly, in the potassium concentrations in the sediments as well as the
229 variability of these potassium concentrations. These differences can be used to separate influences that are
230 specific to particular coring locations from regional influences with features that are clearly common to all
231 areas. The AHP and the dry period during the YD stadial can, for example, be identified in all of the
232 investigated cores and the dry-wet-dry alternations during the D-O events in most of them, while some of the
233 less distinct dry-wet transitions during glacial times can only be observed in the CB01, CB04 and CB06
234 cores, and to a lesser extent also in CB03.

235 The high-resolution K record from the Chew Bahir basin is a promising dataset for testing currently debated
236 hypotheses on the abruptness of the onset of the AHP and the gradual character of its termination (e.g.,
237 deMenocal et al., 2000; Kröpelin et al., 2008a, b; Brovkin and Claussen, 2008; Junginger and Trauth, 2013;
238 Tierney and deMenocal, 2013; Costa, et al., 2014; Shanahan et al., 2015). The onset and termination of the
239 AHP, and also of the YD, are represented by the most dramatic changes in most environmental records from
240 northern and equatorial Africa, for example in the terrestrial dust records from off the West African coast that
241 are used as a proxy for aridity (deMenocal et al., 2000), in the $\delta^{15}\text{N}$ data from the Gulf of Oman (18°N) used
242 as a proxy for denitrification and productivity (Altabet et al., 2002), and in the δD leaf wax data from Lake
243 Tanganyika used as a record of precipitation (Tierney et al., 2008); they are also represented by the most
244 dramatic changes in the K record from lake sediments in the Chew Bahir basin (Figs. 5 and 6).

245 There is a general consensus that the AHP was a result of increased insolation, higher sea-surface
246 temperatures in the Indian Ocean, and increased atmospheric greenhouse gas concentrations (e.g.,
247 Kutzbach and Street-Perrott, 1985; Gasse, 2000; Otto-Bliesner et al., 2014). The results of our statistical
248 analysis leave no doubt that the AHP, as recorded in the sediments of the Chew Bahir basin in southern
249 Ethiopia, started abruptly at ~15,640 a BP. Given that our age model is sufficiently accurate, this transition
250 would have taken less than 250 years (Figs. 5 and 6). In contrast, the termination of the AHP was rather
251 gradual, as recorded by an 990 year transition in the mean K concentration (between 4,875 to 3,885 a BP),
252 or an even longer transition (~1,995 years) in the trend of the K concentration (between 5,835 to 3,840 a
253 BP). Both estimates for the duration of the wet-dry transition at the end of the AHP suggest that the climate
254 changed more rapidly than the orbitally-induced reduction in insolation, which in turn suggests a nonlinear
255 response of the Chew Bahir climate to the resultant forcing (Claussen et al., 1999) (Figs. 5 and 6). The mid-
256 point of the termination of the AHP at ~4,860 a BP, as determined by our change point analysis with a small
257 number of possible transitions, is in agreement with the findings of Kuper and Kröpelin (2006) and Shanahan
258 et al. (2015), suggesting a time-transgressive termination of the AHP during the course of the gradual
259 southward migration of the tropical rain belt over approximately the last 7 ka. Holocene speleothem records
260 from the south-eastern Ethiopian highlands, with annual to decadal resolutions, also show a gradual
261 decrease in rainfall between ~6.5 and 4.5 ka BP (Asrat et al., 2007; Baker et al., 2010).

One of the remarkable features of the K record from the Chew Bahir basin is the significant wet-dry-wet oscillation between 13,260 and 11,730 a BP (Foerster et al., 2012) (Figs. 5 and 6) in which the transition into and out of the dry episode is synchronous (within the errors of the age model) with the high northern latitude Younger Dryas stade (12,900–11,700 a BP; Alley, 2000), suggesting a strong climate teleconnection between high and low latitudes over millennial time scales. This teleconnection is also expressed in the TEX₈₆ temperature and dD leaf wax record from Lake Albert (Berke et al., 2014), which show dramatic variations in temperature and moisture within the YD chronozone. The transition from wet to dry at the onset of the dry episode in the Chew Bahir record occurred within ~45 years (i.e. three data points of the K record), whereas the dry-to-wet transition lasted ~250 years (i.e. 17 data points of the K record). The abruptness and magnitude of change from wet to dry and back again within decades or centuries enables us to discern the impact of this dramatic global (or hemispheric) climate change on the region.

The rapid fluctuations between ~46 and 20 ka BP are reminiscent of the high-latitude D-O cycles, whose expression in the tropics is also a matter of intense discussion (e.g. Brown et al., 2007; Garcin, 2008; Brown et al., 2008; Tierney and deMenocal, 2013). Our analysis clearly demonstrates that D-O type patterns are evident in the Chew Bahir record, as indicated by a high negative correlation between the K concentrations and the $\delta^{18}\text{O}$ record from the NorthGRIP ice core (North Greenland Ice Core Project members, 2004). It is interesting to note that a similar negative correlation also exists between ~15–10 ka BP, including the dry episode during the YD stadial, whereas there is either a positive correlation or no correlation (correlation coefficient values of 0–0.5) during those episodes with wetter climates.

Implications of different climatic transitions for humans

Understanding how the east and northeast African climate switched from dry to wet and back to dry is essential for deciphering the effect that environmental instability may have had on human behaviour (e.g., Bradtmöller et al., 2012; Widlok et al., 2012; Foerster et al., 2015). The K record in Chew Bahir lake sediments provides detailed information on how the environment responded to the gradual changes in insolation that occurred in a nonlinear, saddle-node bifurcation type of transition at the end of the AHP (Trauth et al., 2015). This high-resolution (~15 a) record provides an important basis for discussions on the

288 nature of, and reasons for, the human cultural transition in this region from fishing to herding (Ambrose,
289 1998; Marshall and Hildebrand, 2002; Scheinfeldt et al., 2010). The record clearly shows that the climate
290 evolved from rather stable wet conditions prior to ~7.15 ka BP towards rather stable dry conditions after ~3.8
291 ka BP.

292 Most of the intermediate interval between 7.15 and 3.8 ka BP is characterized by rather wet conditions but
293 these are interrupted by at least 14 dry events that recurred every 160 ± 40 years and lasted between 20 and
294 80 years (Figs. 5 and 6), which may have had a major influence on the livelihoods and life decisions of the
295 human population. The final transition to predominantly dry conditions, and hence the actual termination of
296 the AHP in the Chew Bahir basin, occurred over a relatively long time period of about 990 years. The impact
297 that this prolonged transition from wet to dry conditions had on the livelihood of humans compared to that of
298 a series of precursor droughts that occurred prior to the termination of the AHP is a matter of considerable
299 interest for future investigation. In contrast, the transition from wet to dry within ~45 years (at ~13,260 a BP,
300 within the errors of the age model) at the onset of the dry episode coinciding with the YD stadial must have
301 certainly affected humans living in the area. Our record suggests that this change in climate was likely to
302 have had a much more dramatic effect on human behaviour than the drought events that occurred just
303 before the end of the AHP.

304 It would be interesting to examine the archaeological evidence in order to determine if there was any
305 corresponding change in the behaviour of humans with respect to food gathering, population size, or
306 migration. The end of this dry episode at ~11,730 a BP would have resulted in a return to more favourable
307 conditions for humans. This transition from dry to wet conditions occurred over about 250 years which is
308 considerably longer than a human lifespan and probably did not result in any catastrophic flooding of
309 previously dry lakes (such as the Chew Bahir); it may therefore not have forced any change in human
310 habitats. Analysis of archaeological data from the Chew Bahir region related to these rapid climate changes
311 caused by the high latitude YD teleconnection will improve our understanding of the influence that such
312 changes had on human behaviour. Similarly, archaeological data for time periods characterized by D-O

313 triggered dry-wet-dry fluctuations are likely to provide additional insights into early human behaviour during
314 periods of rapid climate change.

315 Human-environment interactions are difficult to understand and even more difficult to reconstruct from
316 paleorecords. The inter-relationship is complex and although causality is often inferred purely on the basis of
317 time correlation between climatic change and human reaction (evolution, migration, adaption), it may not be
318 as simple a cause-and-effect relationship as often portrayed by natural scientists (Widlöck et al., 2012;
319 Bradtmöller, 2012). High-resolution paleoclimate records such as that of Chew Bahir help us to critically
320 evaluate established hypotheses and to design alternatives, thus leading to a better understanding of
321 human-environment interactions.

322 **ACKNOWLEDGEMENTS**

323 This project was funded by German Research Foundation (DFG) grants to M.H.T. and F.S. through the
324 Priority Program SPP 1006 ICDP and the CRC 806 Research Project at the University of Cologne. We thank
325 C. Ohlendorf and S. Stahl from the Geomorphology and Polar Research Group at the University of Bremen
326 and for the XRF scans. The previously published XRF scan results for the CB01, CB03 and CB05 cores
327 were supported by V. Wennrich at the University of Cologne. This research is part of a pilot study for the
328 Hominin Sites and Paleolakes Drilling Project (HSPDP), within the framework of the International Continental
329 Scientific Drilling Program (ICDP) and the DFG Priority Program SPP 1006 ICDP. This paper is publication
330 number ### of the Hominin Sites and Paleolakes Drilling Project (HSPDP).

331 **REFERENCES CITED**

332 Alley, R., 2000. The Younger Dryas cold interval as viewed from central Greenland. Quaternary Science
333 Reviews, 19, 213-226.
334 Altabet, M.A., Higginson, M.J., Murray, D.W., 2002. The effect of millennial-scale changes in Arabian Sea
335 denitrification on atmospheric CO₂. Nature, 415, 159-162.

336 Ambrose, S.H., 1998. Late Pleistocene human population bottlenecks, volcanic winter, and the differentiation
337 of modern humans. *Journal of Human Evolution*, 35, 115-118.

338 Asrat, A., Baker, A., Umer, M., Moss, J., Leng, M., Van Calstren, P., Smith, C., 2007. A high-resolution multi-
339 proxy stalagmite record from Mechara, Southeastern Ethiopia: Paleohydrological implications for
340 speleothem paleoclimate reconstruction. *Journal of Quaternary Science*, 22, 53-63.

341 Baker, A., Asrat, A., Leng, M., Thomas, L., Fairchild, I.J., Widmann, M., Dong, B., Van Calsteren, P., Bryant,
342 C., 2010. Decadal-scale Holocene Climate Variability Recorded in an Ethiopian Stalagmite. *The*
343 *Holocene*, 20, 827-836.

344 Berger, A., Loutre, M.-F., 1991. Insolation values for the climate of the last 10 million yr. *Quaternary Science*
345 *Reviews*, 10, 297-317.

346 Björck, S., Wohlfarth, B., 2001. ^{14}C chronostratigraphic techniques in paleolimnology, in: *Tracking*
347 *Environmental Change Using Lake Sediments Volume 1: Basin Analysis, Coring, and Chronological*
348 *Techniques*. Last, W. and Smol, J. (eds.), Springer, Dordrecht, Netherlands, 205-246.

349 Blaauw, M., Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive gamma
350 process. *Bayesian Analysis*, 6, 457-474.

351 Bradtmöller, M., Pastoors, A., Weninger, B., Weniger, G.-C., 2012. The repeated replacement model e Rapid
352 climate change and population dynamics in Late Pleistocene Europe. *Quaternary International*, 247, 38-
353 49.

354 Brandt, S.A., Fisher, E.C., Hildebrand, E.A., Vogelsang, R., Ambrose, S.H., Lesur, J. and Wang, H., 2012.
355 Early MIS 3 occupation of Mochena Borago Rockshelter, Southwest Ethiopian Highlands: Implications for
356 Late Pleistocene archaeology, paleoenvironments and modern human dispersals. *Quaternary*
357 *International*, 274, 38-54.

358 Brandt, S.A., Hildebrand, E.A., 2005. Southwest Ethiopia as an Upper Pleistocene Refugium. Paper
359 presented at the Workshop on the Middle Stone Age of Eastern Africa. Nairobi, Kenya and Addis Ababa,
360 Ethiopia, July 2005.

361 Bronk Ramsey, C., 1995. Radiocarbon calibration and analysis of stratigraphy: the OxCal program.
362 *Radiocarbon* 36, 425-430.

363 Bronk Ramsey, C., 2009a. Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51, 337-360.

364 Bronk Ramsey, C., 2009b. Dealing with outliers and offsets in radiocarbon dating. *Radiocarbon*, 51, 1023-
365 1045.

366 Brovkin, V., Claussen, M., 2008. Comment on "Climate-Driven Ecosystem Succession in the Sahara: The
367 Past 6000 Yr". *Science*, 322, 1326b-c.

368 Brown, F.H., Fuller, C.R., 2008. Stratigraphy and tephra of the Kibish Formation, southwestern Ethiopia.
369 *Journal of Human Evolution*, 55, 366-403.

370 Brown, E.T., Johnson T.C, Scholz C.A, Cohen A.S, King J.W., 2007. Abrupt change in tropical African
371 climate linked to the bipolar seesaw over the past 55,000 yr. *Geophysical Research Letters*, 34, L20702.

372 Brown, E.T., Johnson T.C, Scholz C.A, Cohen A.S, King J.W., 2008. Reply to comment by Yannick Garcin
373 on "Abrupt change in tropical African climate linked to the bipolar seesaw over the past 55,000 years".
374 *Geophysical Research Letters*, 35, L04702.

375 Campisano, C., Cohen A..S., Arrowsmith, J.R., Asrat, A., Behrensmeyer, A.K., Brown, E.T., Deino, A.L.,
376 Feibel, C.S., Kingston, J.D., Lamb, H.F., Lowenstein, T.K., Noren, A., Olago, D.O., Owen, B., Pelletier,
377 J.D., Potts, R., Reed, K.E., Renaut, R., Russel, J.M., Russel, J.L., Schäbitz, F., Stone, J.R., Trauth, M.H.,
378 Wynn, J.G., 2017. The Hominin Sites and Paleolakes Drilling Project: High-Resolution Paleoclimate
379 Records from the East African Rift System and Their Implications for Understanding the Environmental
380 Context of Hominin Evolution. *PaleoAnthropology*, 2017, 1-43.

381 Carto, S.L., Weaver, A.J., Hetherington, R., Lam, Y., Wiebe, E.C., 2009. Out of Africa and into an ice age: on
382 the role of global climate change in the late Pleistocene expansion of early modern humans out of Africa.
383 *Journal of Human Evolution*, 56, 139-151.

384 Castañeda, I.S., Mulitza, S., Schefuß, E., Santos, R.A.L., Damste, J.S.S., Schouten, S., 2009. Wet phases in
385 the Sahara/Sahel region and human expansion patterns in North Africa. *Proceedings of the National*
386 *Academy of Sciences*, 106, 1-5.

387 Claussen, M., Kubatzki, C., Brovkin, V., Ganopolski, A., Hoelzmann, P., Pachur, H.J., 1999. Simulation of an
388 abrupt change in Saharan vegetation in the Mid-Holocene. *Geophysical Research Letters*, 26, 2037-
389 2040.

390 Cohen, A., Campisano, C., Arrowsmith, R., Asrat, A., Behrensmeyer, A.K., Deino, A., Feibel, C., Hill, A.,
391 Johnson, R., Kingson, J., Lamb, H., Lowenstein, T., Noren, A., Olago, D., Owen, R.B., Potts, R., Reed,

392 K., Renaut, R., Schäbitz, F., Tiercelin, J.-J., Trauth, M.H., Wynn, J., Ivory, S., Brady, K., O'Grady, R.,
 393 Rodysill, J., Githiri, J., Russell, J., Foerster, V., Dommain, R., Rucina, S., Deocampo, D., Russell, J.,
 394 Billingsley, A., Beck, C., Dorenbeck, G., Dullo, L., Feary, D., Garelo, D., Gromig, R., Johnson, T.,
 395 Junginger, A., Karanja, M., Kimburi, E., Mbuthia, A., McCartney, T., McNulty, E., Muiruri, V., Nambiro, E.,
 396 Negash, E.W., Njagi, D., Wilson, J.N., Rabideaux, N., Raub, T., Sier, M.J., Smith, P., Urban, J., Warren,
 397 M., Yadeta, M., Yost, C., Zinaye, B., 2016. The Hominin Sites and Paleolakes Drilling Project: Inferring
 398 the Environmental Context of Human Evolution from Eastern African Rift Lake Deposits. *Scientific*
 399 *Drilling*, 21, 1-16.

400 Costa, K., Russell, J., Konecky, B., Lamb, H., 2014. Isotopic reconstruction of the African Humid Period and
 401 Congo Air Boundary migration at Lake Tana, Ethiopia. *Quaternary Science Reviews*, 83, 58-67.

402 Davidson, A., 1983. The Omo River project: reconnaissance geology and geochemistry of parts of Ilubabor,
 403 Kefa, Gemu Gofa and Sidamo. *Ethiopian Institute of Geological Surveys Bulletin*, 2, 1-89.

404 deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., Yarusinsky, M., 2000. Abrupt
 405 onset and termination of the African Humid Period: rapid climate responses to gradual insolation forcing.
 406 *Quaternary Science Reviews*, 19, 347-361.

407 Foerster, V., Junginger, A., Langkamp, O., Gebru, T., Asrat, A., Umer, M., Lamb, H., Wennrich, V.,
 408 Rethemeyer, J., Nowaczyk, N., Trauth, M.H., Schäbitz, F., 2012. Climatic change recorded in the
 409 sediments of the Chew Bahir basin, southern Ethiopia, during the last 45,000 yr. *Quaternary International*,
 410 274, 25-37.

411 Foerster, V., Vogelsang, R., Junginger, A., Asrat, A., Lamb, H.F., Schaebitz, F., Trauth,
 412 M.H., 2015. Environmental Change and Human Occupation of Southern Ethiopia and Northern Kenya
 413 during the last 20,000 yr. *Quaternary Science Reviews*, 129, 333-340.

414 Foerster, V., Vogelsang, R., Junginger, A., Asrat, A., Lamb, H.F., Schaebitz, F., Trauth, M.H., 2016. Reply to
 415 comment on "Environmental Change and Human Occupation of Southern Ethiopia and Northern Kenya
 416 during the last 20,000 years. *Quaternary Science Reviews* 129:333-340". *Quaternary Science Reviews*,
 417 141, 130-133.

418 Fritsch, F.N., Carlson, R.E., 1980. Monotone Piecewise Cubic Interpolation. *SIAM J. Numerical Analysis*, 17,
 419 238-246.

420 Garcin, Y., Vincens, A., Williamson, D., Buchet, G., Guiot, J., 2007. Abrupt resumption of the African
 421 Monsoon at the Younger Dryas-Holocene climatic transition. *Quaternary Science Reviews*, 26, 690-704.

422 Garcin, Y., 2008. Comment on "Abrupt change in tropical African climate linked to the bipolar seesaw over
 423 the past 55,000 yr" by Brown, E. T., Johnson, T. C., Scholz, C. A., Cohen, A. S., and King, J. W. (2007).
 424 *Geophysical Research Letters*, 35, L04701.

425 Garcin, Y., Melnick, D., Strecker, M.R., Olago, D., Tiercelin, J.-J., 2012. East African mid-Holocene wet-dry
 426 transition recorded in palaeo-shorelines of Lake Turkana, northern Kenya Rift. *Earth and Planetary
 427 Science Letters*, 331-332, 322-334.

428 Gasse, F., 2000. Hydrological changes in the African tropics since the Last Glacial Maximum. *Quaternary
 429 Science Reviews* 19, 189-211.

430 Hailemeskel, A., Fekadu, H., 2004. Geological map of Yabello. Geological Survey of Ethiopia, Addis Ababa
 431 ISBN 0000 0001 0674 8528.

432 Hassen, N., Yemane, T., Genzebu, W., 1997. Geology of the Agere Maryam Area. Geological Survey of
 433 Ethiopia, Addis Ababa.

434 Hildebrand, E., Brandt, S., Lesur-Gebremariam, J., 2010. The Holocene archaeology of Southwest Ethiopia:
 435 new insights from the Kafa archaeological project. *African Archaeological Review*, 27, 255-289.

436 Hildebrand, E., Grillo, K., 2012. Early herders and monumental sites in eastern Africa: New radiocarbon
 437 dates. *Antiquity*, 86, 338-352.

438 Hotelling, H., 1931. Analysis of a Complex of Statistical Variables with Principal Components, *Journal of
 439 Educational Psychology*, 24, 417-441.

440 Hublin, J.-J. et al., 2017. New fossils from Jebel Irhoud, Morocco and the pan-African origin of *Homo
 441 sapiens*. *Nature Geoscience*, 547, 289-292.

442 Johnson, T.C., Halfman, J.D., Showers, W.J., 1991. Paleoclimate of the past 4000 yr at Lake Turkana,
 443 Kenya, based on the isotopic composition of authigenic calcite. *Palaeogeography, Palaeoclimatology,
 444 Palaeoecology*, 85, 189-198.

445 Junginger, A., Trauth, M.H., 2013. Hydrological constraints of paleo-Lake Suguta in the Northern Kenya Rift
 446 during the African Humid Period (15 - 5 ka BP). *Global and Planetary Change*, 111, 174-188.

447 Junginger, A., Roller, S., Trauth, M.H., 2014. The effect of solar irradiation changes on water levels in the
 448 paleo-Lake Suguta, Northern Kenya Rift, during the late Pleistocene African Humid Period (15-5 ka
 449 BP). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 396, 1-16.

450 Key, R.M., 1988. Geology of the Sabarei area: Degree sheets 3 and 4, with coloured 1:250 000 geological
 451 map and results of geochemical exploration (Report). Ministry of Environment and Natural Resources,
 452 Mines and Geology Dept., Nairobi, Kenya.

453 Killick, R., Fearnhead, P., Eckley, I.A., 2012. Optimal detection of change points with a linear computational
 454 cost. *Journal of the American Statistical Association*, 107, 1590-1598.

455 Kröpelin, S., Verschuren, D., Lézine, A.-M., Eggermont, H., Cocquyt, C., Francus, P., Cazet, J.-P., Fagot, M.,
 456 Rumes, B., Russell, J.M., Darius, F., Conley, D.J., Schuster, M., von Suchodoletz, H., Engstrom, D.R.,
 457 2008a. Climate-Driven Ecosystem Succession in the Sahara: The Past 6000 Yr. *Science* 320, 765-768.

458 Kröpelin, S., Verschuren, D., Lézine, A.-M., 2008b. Response to Comment by Brovkin and Claussen on
 459 "Climate-Driven Ecosystem Succession in the Sahara: The Past 6000 Yr". *Science*, 322, 1326.

460 Kuper, R., Kröpelin, S., 2006. Climate-Controlled Holocene Occupation in the Sahara: Motor of Africa's
 461 Evolution. *Science*, 313, 803-307.

462 Kutzbach, J.E., Street-Perrott, F.A., 1985. Milankovitch forcing of fluctuations in the level of tropical lakes
 463 from 18 to 0 ka BP. *Nature*, 317, 130-134.

464 Lamb, H.F., Bates, C.R., Coombes, P.V., Marshall, M.H., Umer, M., Davies, S.J., Dejen, E., 2007. Late
 465 Pleistocene desiccation of Lake Tana, source of the Blue Nile. *Quaternary Science Reviews*, 26, 287-
 466 299.

467 Levin, N.E., Zipser, E.J., Cerling, T.E., 2009. Isotopic composition of waters from Ethiopia and Kenya:
 468 Insights into moisture sources for eastern Africa. *Journal of Geophysical Research*, 114, D23306.

469 Martínez, I., Keigwin, L., Barrows, T.T., Yokoyama, Y., Southon, J., 2003. La Niña-like conditions in the
 470 eastern equatorial Pacific and a stronger Choco jet in the northern Andes during the last glaciation.
 471 *Paleoceanography*, 18, 1033-1033.

472 Marshall, F., Hildebrand, E., 2002. Cattle before crops: the beginnings of food production in Africa. *Journal of*
 473 *World Prehistory*, 16, 99-143.

474 MathWorks, 2017a. MATLAB Signal Processing Toolbox – User's Guide. The MathWorks, Inc., Natick, MA.

MathWorks, 2017b. MATLAB Statistics Toolbox – User’s Guide. The MathWorks, Inc., Natick, MA.

McDougall, I., Brown, F.H., Fleagle, J.G., 2005. Stratigraphic placement and age of modern humans from Kibish, Ethiopia. *Nature*, 433, 733-736.

Moore, J.M., Davidson, A., 1978. Rift structure in southern Ethiopia. *Tectonophysics*, 46, 159-173.

Nicholson, S.E., 2017. Climate and climatic variability of rainfall over eastern Africa. *Reviews of Geophysics*, 55, 590-635.

North Greenland Ice Core Project members, 2004. High-resolution record of northern hemisphere climate extending into the last interglacial period. *Nature*, 431, 147-151.

Otto-Bliesner, B.L., Russell, J.M., Clark, P.U., Liu, Z., Overpeck, J.T., Konecky, B., deMenocal, P., Nicholson, S.E., He, F., Lu, Z., 2014. Coherent changes of southeastern equatorial and northern African rainfall during the last deglaciation. *Science*, 346, 1223-1227.

Paliwal, K.K., Agarwal, A., Sinha, S.S., 1982. A Modification over Sakoe and Chiba’s Dynamic Time Warping Algorithm for Isolated Word Recognition. *Signal Processing*, 4, 329-333.

Pearson, K., 1901. On lines and planes of closest fit to a system of points in space. *Philosophical Magazine and Journal of Science*, 6, 559-572.

Reimer, P.J. et al., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Yr cal BP. *Radiocarbon*, 55, 1869-1887.

Richter, D. et al., 2017. The age of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age. *Nature Geoscience*, 546, 293-296.

Roberts, N., Taieb, M., Barker, P., Damnati, B., Icole, M., Williamson, D., 1993. Timing of the Younger Dryas event in east Africa from lake-level changes. *Nature*, 366, 146-148.

Sakoe, H., Chiba, S., 1978. Dynamic Programming Algorithm Optimization for Spoken Word Recognition. *IEEE Transactions on Acoustics, Speech, and Signal Processing*, ASSP-26, 43-49.

Scheinfeldt, L.B., Soi, S., Tishkoff, S.A., 2010. Working toward a synthesis of archaeological, linguistic, and genetic data for inferring African population history. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 8931-8938.

501 Shanahan, T.M., McKay, N.P., Hughen, K.A., Overpeck, J.T., Otto-Bliesner, B., Heil, C.W., King, J., Scholz,
502 C.A., Peck, J., 2015. The time-transgressive termination of the African Humid Period. *Nature Geoscience*,
503 8, 140-144.

504 Tierney, J.E., Russell, J.M., Sinninghe Damsté, J.S., Huang, Y., Verschuren, D., 2011. Late Quaternary
505 behavior of the east African monsoon and the importance of the Congo Air Boundary. *Quaternary*
506 *Science Reviews*, 30, 798-807.

507 Tierney, J.E., deMenocal, P.B., 2013. Abrupt Shifts in Horn of Africa. *Hydroclimate Since the Last Glacial*
508 *Maximum*. *Science*, 342, 843-846.

509 Tierney, J.E., deMenocal, P.B., Zander, P.D., 2017. A climatic context for the out-of-Africa migration.
510 *Geology*, 45, 1023-1026.

511 Trauth, M.H., 2014. A new probabilistic technique to determine the best age model for complex stratigraphic
512 sequences. *Quaternary Geochronology*, 22, 65-71.

513 Trauth, M.H., Bergner, A.G.N., Foerster, V., Junginger, A., Maslin, M.A., Schaebitz, F., 2015. Episodes of
514 Environmental Stability and Instability in Late Cenozoic Lake Records of Eastern Africa. *Journal of*
515 *Human Evolution*, 87, 21-31.

516 Vogelsang, R., Keding, B., 2013. Climate, culture, and change: From hunters to herders in northeastern and
517 southwestern Africa, in: *Comparative Archaeology and Paleoclimatology – Socio-cultural responses to a*
518 *changing world*, Baldia, M.O., Perttula, T.K., Frink, D.S. (eds.), BAR International Series, 2456,
519 Archaeopress, Oxford, UK, 43-62.

520 Widlok, T., Aufgebauer, A., Bradtmöller, M., Dikau, R., Hoffmann, T., Kretschmer, I., Panagiotopoulos,
521 K., Pastoors, A., Peters, R., Schäbitz, F., Schlummer, M., Solich, M., Wagner, B., Weniger, G.-C.,
522 Zimmermann, A., 2012. Towards a theoretical framework for analyzing integrated socio-environmental
523 systems. *Quaternary International*, 274, 259-272.

524 LIST OF TABLES

525 Table 1 | Cored locations within the Chew Bahir basin (longitudes and latitudes), core lengths, total depth of
526 cored material below the lake floor, date of core collection, and core recovery.

527 LIST OF FIGURES

528 Figure 1 | (A) Geologic map of the Chew Bahir basin, showing the three generalized rock types: Cenozoic rift
529 sediments, Cenozoic rift volcanics, and Proterozoic basement. Compilation based on Omo River Project
530 Map ([Davidson, 1983](#)), Geology of the Sabarei Area ([Key, 1987](#)), Geology of the Yabello Area ([Hailemeskel
531 and Fekadu, 2004](#)), and Geology of the Agere Maryam Area ([Hassen et al., 1997](#)). (B) Topographic map of
532 the Chew Bahir basin, showing the outline of the catchment, the drainage network, the location of the short
533 cores in the pilot study (2009, 2010), and the 2014 HSPDP-CHB drill site.

534 Figure 2 | Variation in the K concentration of the CB01–06 cores (in standardized counts per second, cps)
535 plotted against the core depth (in cm). For comparison with CB01, which was measured using a
536 molybdenum tube, all records were standardized (mean=0, standard deviation=1). Please note reverse scale
537 of potassium axis.

538 Figure 3 | Variation in the K content (in standardized counts per second, cps) of the Chew Bahir plotted
539 against age (in ka BP). We used dynamic time warping (DTW) to automatically align the K records from
540 cores CB02 to CB06 with that from core CB01. The core depths of the aligned cores were converted into
541 ages using the age model of [Trauth et al. \(2015\)](#). Please note reverse scale of potassium axis. We have
542 offset the individual time series by 2 standardized cps in each case to allow an overlap-free display of the K
543 records.

544 Figure 4 | Results of a principal component analysis to unmix (or separate) the regional environmental signal
545 from the local signal in cores CB01 to CB06. More than 94% of the total variance of the data set is contained
546 in the first principal component (PC1), which we interpret to represent the regional climate signal. We have
547 offset the individual time series by 2 standardized cps in each case to allow an overlap-free display of the K
548 records.

549 Figure 5 | Results of change point analyses on the K record after DTW alignment and PCA-based unmixing:
550 (a) change points in the mean; (b) change points in the standard deviation; (c) change points in the trend.

551 The change point algorithm yields about 15 significant transitions in the mean, standard deviation and trend,
552 the most marked of which occur at the onset of, and during, the termination of the African Humid Period
553 (AHP), interrupted by the Younger Dryas (YD) dry episode. Note reversed scale of potassium axis.

554 Figure 6 | Comparing the K record after DTW alignment and PCA-based unmixing with the $\delta^{18}\text{O}$ data from
555 the NorthGRIP ice core ([North Greenland Ice Core Project members, 2004](#)) suggests a weak negative
556 correlation (Spearman coefficient less than -0.5) between environmental fluctuations in the Chew Bahir
557 basin and high-latitude northern hemisphere millennial-scale climate fluctuations that occurred between 46
558 and 25 ka BP (during the Dansgaard-Oeschger cycles), and between 15 and 5 ka BP (including the Younger
559 Dryas stade). Please note reverse scale of potassium axis.