Aberystwyth University

Resting-state global functional connectivity as a biomarker of cognitive reserve in mild cognitive impairment

Franzmeier, Nicolai; Araque Caballero, Miguel A.; Taylor, Alexander; Simon-Vermot, Lee; Beurger, Katharina; Ertl-Wagner, Birgit; Mueller, Claudia; Catak, Cihan; Janowitz, Daniel; Baykara, Ebru; Gesierich, Benno; Duering, Marco; Ewers, Michael

Published in:
Brain Imaging and Behavior

DOI:
[10.1007/s11682-016-9599-1](https://doi.org/10.1007/s11682-016-9599-1)

Publication date:
2017

Citation for published version (APA):
Franzmeier, N., Araque Caballero, M. A., Taylor, A., Simon-Vermot, L., Beurger, K., Ertl-Wagner, B., Mueller, C., Catak, C., Janowitz, D., Baykara, E., Gesierich, B., Duering, M., & Ewers, M. (2017). Resting-state global functional connectivity as a biomarker of cognitive reserve in mild cognitive impairment. *Brain Imaging and Behavior*, 11(2), 368-382. <https://doi.org/10.1007/s11682-016-9599-1>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Brain Imaging and Behavior

Resting-state global functional connectivity as a biomarker of cognitive reserve in mild cognitive impairment --Manuscript Draft--

Manuscript Number:		
Full Title:	Resting-state global functional connectivity as a biomarker of cognitive reserve in mild cognitive impairment	
Article Type:	SI: Resilience/Reserve in AD	
Keywords:	cognitive reserve; biomarker; mild cognitive impairment; Alzheimer's disease; global functional connectivity; resting-state fMRI	
Corresponding Author:	Nicolai Franzmeier, M.Sc. Institute for Stroke and Dementia Research Munich, GERMANY	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Institute for Stroke and Dementia Research	
Corresponding Author's Secondary Institution:		
First Author:	Nicolai Franzmeier, M.Sc.	
First Author Secondary Information:		
Order of Authors:	Nicolai Franzmeier, M.Sc. Miguel Angel Araque Caballero, Dr. Alexander Nigel William Taylor, Dr. Lee Simon-Vermot, M.Sc. Katharina Buerger, PD Dr. Birgit Ertl-Wagner, Prof. Dr. Cihan Catak, Dr. Daniel Janowitz Claudia Mueller, Dr. Ebru Baykara, M.Sc. Benno Gesierich, Dr. Marco Duering, PD Dr. Michael Ewers, Prof. Dr.	
Order of Authors Secondary Information:		
Funding Information:	LMUexcellent European Research Council (PCIG12-GA-2012-334259) Alzheimer's Forschung Initiative (DE-15035) National Institutes of Health (U01 AG024904) U.S. Department of Defense (US) (W81XWH-12-2-0012)	Mr. Michael Ewers Mr. Michael Ewers Mr. Michael Ewers Not applicable Not applicable
Abstract:	Cognitive reserve (CR) shows protective effects in Alzheimer's-disease (AD) and reduces the risk of dementia. Despite the clinical significance of CR, a clinically useful	

diagnostic biomarker of brain changes underlying CR is not available yet. Our aim was to develop a fully-automated approach applied to fMRI to produce a biomarker associated with CR in subjects at increased risk of AD. We computed resting-state global functional connectivity (GFC), i.e. the average connectivity strength, for each voxel within the cognitive control network, which may sustain CR due to its central role in higher cognitive function. In a training sample including 43 mild cognitive impairment (MCI) subjects and 24 healthy controls (HC), we found that MCI subjects with high CR (> median of years of education, CR+) showed increased frequency of high GFC values compared to MCI CR- and HC. A summary index capturing such a surplus frequency of high GFC was computed (called GFC reserve (GFC-R) index). GFC-R discriminated MCI CR+ vs MCI CR-, with the area under the ROC = 0.84. Cross-validation in an independently recruited test sample of 23 MCI subjects showed that higher levels of the GFC-R index predicted higher years of education and an alternative questionnaire-based proxy of CR (CRIq score), controlled for memory performance, gray matter of the cognitive control network, age, and gender. In conclusion, the GFC-R index that captures GFC changes within the cognitive control network provides a biomarker candidate of functional brain changes of CR in patients at increased risk of AD.

Regular Article

Title: Resting-state global functional connectivity as a biomarker of cognitive reserve in mild cognitive impairment.

Authors: N. Franzmeier,¹ M. Á. Araque Caballero,¹ A. Taylor,¹ L. Simon-Vermot,¹ K. Buerger, B.,^{1,2} B. Ertl-Wagner,¹ C. Mueller,¹ C. Catak,¹ D. Janowitz,¹ E. Baykara,¹ B. Gesierich,¹ M. Duering,¹ M. Ewers¹ for the Alzheimer's Disease Neuroimaging Initiative*

1 Institute for Stroke and Dementia Research, Klinikum der Universität München, Ludwig-Maximilians-Universität LMU, Feodor-Lynen Straße 17, 81377 Munich, Germany.

2 German Center for Neurodegenerative Diseases (DZNE, Munich), Feodor-Lynen Straße 17, 81377 Munich, Germany.

* Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Corresponding author: Nicolai Franzmeier, M.Sc.

Corresponding author's address: Klinikum der Universität München, Institut für Schlaganfall- und Demenzforschung (ISD), Feodor-Lynenstraße 17, D-81377 Munich.

Corresponding author's phone and fax: +49 (0) 89440046221, ++49 (0) 89 4400-46162

Corresponding author's e-mail address: nicolai.franzmeier@med.uni-muenchen.de

Number of characters in title: 96

Number of characters in running head:

Number of words in abstract: 248

Number of words in main text: 5911

Number of words in introduction: 611

Number of words in discussion: 1627

Number of color figures: 5

Number of tables: 2

ABSTRACT:

Cognitive reserve (CR) shows protective effects in Alzheimer's-disease (AD) and reduces the risk of dementia. Despite the clinical significance of CR, a clinically useful diagnostic biomarker of brain changes underlying CR is not available yet. Our aim was to develop a fully-automated approach applied to fMRI to produce a biomarker associated with CR in subjects at increased risk of AD. We computed resting-state global functional connectivity (GFC), i.e. the average connectivity strength, for each voxel within the cognitive control network, which may sustain CR due to its central role in higher cognitive function. In a training sample including 43 mild cognitive impairment (MCI) subjects and 24 healthy controls (HC), we found that MCI subjects with high CR (> median of years of education, CR+) showed increased frequency of high GFC values compared to MCI CR- and HC. A summary index capturing such a surplus frequency of high GFC was computed (called GFC reserve (GFC-R) index). GFC-R discriminated MCI CR+ vs MCI CR-, with the area under the ROC = 0.84. Cross-validation in an independently recruited test sample of 23 MCI subjects showed that higher levels of the GFC-R index predicted higher years of education and an alternative questionnaire-based proxy of CR (CRIq score), controlled for memory performance, gray matter of the cognitive control network, age, and gender. In conclusion, the GFC-R index that captures GFC changes within the cognitive control network provides a biomarker candidate of functional brain changes of CR in patients at increased risk of AD.

INTRODUCTION:

Cognitive reserve (CR) refers to the ability to cognitively perform relatively well in the presence of brain pathology (Stern, 2002, 2009). Life-time experiences - such as education and occupational attainment – or IQ are commonly used as proxy measures of CR (Stern, 2009). In Alzheimer's disease (AD), higher levels of such CR proxies are associated with higher cognitive performance relative to the level of brain damage, such as measured by cerebral FDG-PET hypometabolism or impaired blood flow (Bastin et al., 2012; Boots et al., 2015; Ewers et al., 2014; Scarmeas et al., 2003; Stern, Alexander, Prohovnik, & Mayeux, 1992; Stern et al., 1995), grey matter atrophy (Bastin et al., 2012; Boots et al., 2015), white matter damage (Brickman et al., 2011), and primary pathologies including amyloid-beta ($A\beta$) and tau (Rentz et al., 2010; Vemuri et al., 2015; Vemuri et al., 2011). These results suggest that higher levels of CR as measured by education and other proxies are associated with a higher ability to cope with brain pathology in AD.

Compensatory functional brain changes that may underlie CR have been investigated in a number of task-related fMRI studies (Stern et al., 2005; Stern et al., 2008) or resting state fMRI studies in HC subjects (Arenaza-Urquijo et al., 2013). Task-related fMRI studies in MCI and AD revealed an association between increased CR proxies (education, occupation) and higher brain activation (Bosch et al., 2010; Solé-Padullés et al., 2009). However, task-fMRI is often difficult to perform for cognitively impaired patients, and may thus not be suitable for clinical use to assess CR in AD. From a clinical point of view, a major question is whether simple measures of basic brain function are indicative of CR, and thus could be used as a marker of CR-related brain changes in AD. The need of a biomarker of CR-related brain changes is urgent in view of a growing number of clinical trials that target protective brain mechanisms in AD, such as cognitive training or meditation (Buschert et al., 2011; Reijnders, van Heugten, & van Boxtel, 2013; Schultz et al., 2015; Wells et al., 2013).

1 The overall goal of the current study was to develop a neuroimaging-based diagnostic
2 biomarker of functional brain changes underlying CR in subjects with mild cognitive
3 impairment (MCI). We focused on resting-state global functional connectivity (GFC, also
4 known as weighted degree centrality) within the cognitive control network as a measure of
5 functional brain processes of CR. The cognitive control network includes major brain hubs
6 with high GFC (Cole, Pathak, & Schneider, 2010), where greater GFC has been previously
7 associated with higher IQ, i.e. a proxy of CR, in young subjects. In the current study, CR was
8 measured by the proxy of years of education, which is the best validated CR proxy measure to
9 date in AD (Stern, 2012).

10 Using a cross-validation approach, we compared the frequency distribution of GFC values
11 within the cognitive control network between MCI subjects with high CR (more years of
12 education) to MCI with low CR (lower years of education) and HC groups. A newly
13 developed summary index that detects GFC frequency differences between MCI subjects with
14 low and MCI subjects with high CR, henceforth called GFC reserve (GFC-R) index, was
15 tested as a predictor of years of education in an independent validation sample of MCI
16 subjects. We hypothesized firstly that MCI subjects with more years of education show an
17 increased number of relatively high GFC values within the cognitive control network
18 compared to MCI subjects with less years of education. Secondly, we hypothesized that
19 higher levels of the GFC-R index are predictive of more years of education and a second
20 questionnaire based CR proxy in the validation sample of MCI subjects.

METHODS:

Subjects:

We included two independent samples each of amnesic MCI and HC subjects to cross-validate our findings. The training sample included 24 amyloid-PET negative ($A\beta^-$) HC subjects and 43 Amyloid-PET positive ($A\beta^+$) patients with amnesic MCI. Amyloid PET status was defined based on pre-established cut-off values of global [^{18}F] AV-45 PET standardized uptake value ratio (for $A\beta^-$ = global AV-45 PET SUVR < 1.11) (Landau et al., 2013). All data were downloaded from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database, freely accessible for researchers (<http://adni.loni.usc.edu/>). ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial MRI, PET, other biological markers, and clinical and neuropsychological assessment can be combined to measure and predict the progression of MCI and early AD (www.adni-info.org).

The test sample comprised 32 HC subjects as well as 23 subjects with amnesic MCI, recruited between 2014 and 2015 at the memory clinic of the Institute for Stroke and Dementia Research (ISD) at the Klinikum der Universitaet Muenchen in Germany.

For the ISD study, the inclusion criteria were defined as follows: 1) age > 60 years, 2) no signs of depression, 3) no presence or history of neurological or psychiatric disorders (except for MCI), 4) no presence or history of alcohol or drug abuse, 5) no diabetes mellitus, 6) no MRI contraindications. All subjects underwent structural MRI, resting-state fMRI and cognitive testing using the CERAD-Plus test battery (Luck, 2009). A subject was defined as HC, when reporting no subjective memory complaints and scoring within 1.5 standard deviations (SD) of the age, gender and education adjusted norms in all subtests of the CERAD-Plus battery (Luck, 2009). MCI was diagnosed according to the Petersen criteria (Petersen, 2004), when scoring 1.5 SD below the age, gender and education adjusted norms in at least one of the learning or recall subtests of the CERAD-Plus battery.

For the training sample (ADNI), details about the inclusion can be found online (<https://adni.loni.usc.edu/wp-content/uploads/2008/07/adni2-procedures-manual.pdf>). Similar to the diagnosis of MCI in the ISD sample, MCI was diagnosed in ADNI according to the Petersen criteria (Petersen, 2004).

Cognitive reserve and neuropsychological assessment:

The number of years of formal education was used as a proxy for CR in both samples. In the test sample (ISD) we additionally used the cognitive reserve index questionnaire (Nucci, Mapelli, & Mondini, 2012) as a second proxy of CR. The CRIq is a standardized questionnaire based measure for the assessment of CR that combines information about education, working activity and leisure time. For neuropsychological assessments, memory performance was assessed using memory tests that were comparable between the ADNI and the ISD sample. As a measure of episodic memory performance, the total score of the CERAD word list learning tests was assessed in the ISD sample (Luck, 2009), and the total score of the Rey Auditory Verbal Learning Test (RAVLT) in the ADNI sample (Schoenberg et al., 2006). Both tests are designed as list-learning paradigms in which the patient is read a list of words by the examiner in several trials (CERAD: 10 words in 3 trials; RAVLT: 15 words in 15 trials) and is asked to recall as many words from the list as possible after each trial. The total score reflects the number of words correctly remembered cumulated across trials.

Standard protocol approvals and patient consent:

The study at the ISD was approved by the ethics committee of the Ludwig Maximilian University of Munich and was conducted in accordance with the Declaration of Helsinki. All study participants provided written, informed consent to the study. For the ADNI-sample ethical approval was obtained by the ADNI investigators.

MRI acquisition:

Training sample (ADNI):

All MRI scans were performed on Philips 3T MRI scanners, using an 8-channel head matrix coil. High-resolution T1-weighted scans were acquired using a 3D MP-RAGE sequence, with whole brain coverage and a voxel resolution of $1 \times 1 \times 1.2$ mm. Resting state-fMRI images were acquired using a single shot T2*-weighted EPI sequence collecting 140 volumes, with a TR of 3000 ms, a flip angle of 80° and 3.3 mm isotropic voxel resolution. Prior to the resting-state scan, subjects were instructed to keep their eyes open.

Test sample (ISD):

All MRI scans were performed on a Siemens Verio 3T MRI scanner using a 32-channel head coil. For each subject a structural image was obtained using a high-resolution 3D MPRAGE T1-weighted sequence with 1 mm isotropic voxel resolution. Functional resting-state images were acquired using a T2*-weighted echo-planar imaging (EPI) pulse sequence collecting 180 volumes with a TR = 2580 ms, flip angle = 80° and 3.5 mm isotropic voxel resolution. Prior to the resting-state scan the subjects were instructed to keep their eyes closed and not to fall asleep during the scanning procedure. Using the same field of view as the functional resting-state images, field maps were acquired (TE = 7.38/4.92 ms, TR = 675 ms) to correct for susceptibility artifacts and inhomogeneity of the magnetic field during preprocessing of the resting-state data.

Spatial normalization of MRI scans:

The spatial normalization of the MRI scans was done separately for both samples, following the same protocol of image processing based on SPM 12 (Wellcome Trust Centre for Neuroimaging, University College London, United Kingdom: www.fil.ion.ucl.ac.uk/spm).

1 T1-weighted images were segmented into probabilistic maps of grey matter, white matter and
2 cerebrospinal fluid maps through the SPM's new-segment approach (Ashburner & Friston,
3 2005). Next, the spatial normalization parameters were estimated using a high-dimensional
4 diffeomorphic registration algorithm to warp each subjects' grey matter map to a group-
5 specific grey matter template that was defined in an iterative procedure, as implemented in
6 SPM's DARTEL toolbox (Ashburner, 2007). Subsequently, the group-specific template was
7 registered to the MNI template in order to estimate the affine transformation parameters.
8 Next, the non-linear (DARTEL flow-fields) and the affine transformation parameters were
9 combined and applied to the segmented grey matter maps, so that all grey matter images were
10 spatially normalized to the MNI space. The spatially-normalized grey matter maps were
11 averaged and binarized at a voxel value > 0.3 to create a group-specific grey matter mask for
12 later fMRI functional connectivity analyses. Similarly, we averaged and binarized the
13 spatially-normalized white-matter (binarized at threshold > 0.9) and cerebrospinal-fluid
14 (binarized at threshold > 0.7) that were used during preprocessing of the resting-state fMRI
15 data. For later extraction of grey matter volume, we created spatially normalized grey matter
16 maps for each subject, that were smoothed with a 8-mm full-width at half-maximum
17 (FWHM) Gaussian kernel and modulated to preserve the volume of the images.

43 **Preprocessing of resting-state fMRI:**

44 The preprocessing of both samples was done separately, but following the same protocol. All
45 volumes were realigned to the first volume to correct for motion, coregistered to native-space
46 T1-weighted images and smoothed using an 8 mm FWHM Gaussian kernel. None of the
47 subjects' motion parameters exceeded 2 mm translations or 2 degrees rotations. Unique to the
48 ISD-sample, there was additional slice-timing and field map correction. Next, the DARTEL
49 flow-fields and affine registration parameters that were estimated during preprocessing of the
50 T1-weighted images were combined and applied to all resting-state fMRI volumes to spatially

1 normalize the images to MNI space. The spatially normalized fMRI images were further
2 detrended and band-pass filtered, using a frequency band of 0.01-0.08 Hz. In a second step we
3
4 regressed out the 6 motion parameters (3 translations, 3 rotations) and the BOLD signal
5
6 averaged across the white matter and cerebrospinal fluid masks that were created during
7
8 preprocessing of the T1-weighted images.
9
10

11 **Assessment of GFC:**

12
13
14 For each subject, GFC was determined based on the preprocessed and spatially normalized
15
16 resting-state fMRI scans. For each voxel in the grey matter, the GFC was determined by
17
18 computing seed-based Pearson-Moment correlations of the BOLD signal changes between the
19
20 seed voxel and each of the other voxels within the grey matter (as defined by the customized
21
22 grey matter mask). For each seed voxel, only Pearson-Moment correlation coefficients with r
23
24 > 0 were retained, Fisher z -transformed and averaged across the voxels within the grey-
25
26 matter-mask space to obtain the GFC coefficient. This resulted in a 3D brain map of GFC
27
28 coefficients for each subject. Note that we included only positive correlation coefficients for
29
30 computing the GFC, because positive and negative correlations may cancel each other out
31
32 when averaging the correlation coefficients.
33
34
35
36
37
38
39
40
41
42

43 **Spatial maps of resting-state networks:**

44
45 The cognitive control network covers the anterior cingulate cortex, dorsolateral prefrontal
46
47 cortex, anterior insular cortex, dorsal premotor cortex and posterior parietal cortex (Cole,
48
49 Bassett, Power, Braver, & Petersen, 2014; Cole et al., 2013; Cole & Schneider, 2007). For the
50
51 current study, we determined the spatial boundaries of the cognitive control network based on
52
53 an a-priori conducted meta-analysis in order to avoid a sample specific bias in the spatial
54
55 definition of the network. The meta-analysis was conducted using NeuroSynth, a web-based
56
57 tool for fully automated detection of brain activation coordinates from published task-fMRI
58
59
60
61
62
63
64
65

data (<http://www.neurosynth.org>). By entering a search term in the NeuroSynth database, brain activation associated with the search term entered is analyzed across studies, yielding a probabilistic map of brain activation related to that term (Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011). For the current study, we used “cognitive control” as a search term, yielding a z-scored probability map based on 428 task-fMRI studies (as of September 14, 2015). In order to obtain a reliable map of the cognitive control network we applied a false discovery rate corrected p-threshold of $p(\text{FDR}) < 0.01$ (see Fig. 1). The spatial map of the cognitive control network was additionally masked with the group-specific grey matter masks for each sample separately in order to restrict all further analyses to voxels that had a high likelihood of falling within the grey matter. We performed control analyses on 7 major brain networks (Yeo et al., 2011), to test whether a relationship between CR and GFC was specific to the cognitive control network. Accordingly, we downloaded the 7 network parcellations that are freely available online (ftp://surfer.nmr.mgh.harvard.edu/pub/data/Yeo_JNeurophysiol11_MNI152.zip). Again, all 7 networks were additionally masked with the group specific grey matter masks for each sample. To control for potentially confounding effects of brain atrophy, we extracted the grey matter volume within the network masks for each subject, applied to the modulated smoothed and normalized grey matter images that were created during the preprocessing of structural MRI images.

Generation of GFC index related to CR (GFC-R index):

Study design

Our aim was to develop a summary index to quantify GFC frequency changes within the cognitive control network that were associated with the CR proxy years of education in patients with MCI. In brief, the ADNI sample served as a training sample to create the GFC-R index that is related to our CR proxy *education*. Subsequently, we tested the validity of this

GFC based index as a predictor of the CR proxy years of education and the CR-questionnaire (CRIq) composite score in the ISD sample, which served as an independent test sample. A Flow diagram illustrating the individual steps to create the GFC-reserve index is shown in Fig 2.

Dichotomization of subjects according to CR

The HC and MCI groups were each dichotomized into groups of low and high CR (CR- vs CR+), split at the median of years of education within the entire sample. The groups were dichotomized separately within the ISD (CR+: > median education = 13) and the ADNI sample (CR+: > median education = 16).

Histogram Analysis of GFC

For each diagnostic group (MCI vs. HC) within the CR+ and CR- subjects, we plotted a histogram of the GFC frequencies across voxels of the cognitive control network (Figure 2A & B). Visual inspection of the histograms in the training sample (ADNI) revealed, that the GFC histogram of the MCI CR- subjects showed an overall shift to the left of the HC subjects, with a decreased frequency of relatively high GFC values, but an increase of lower GFC values compared to the HC CR- group (Figure 2B). Conversely, the GFC histogram of the MCI CR+ subjects showed a shift to the right of the HC CR+ group.

In a next step, we binned the GFC voxel values for each subject at intervals of 0.01 from $z = 0$ to $z = 0.6$ resulting in a total of 60 bins, each containing the number of voxels (i.e. the GFC frequency) falling within that bin. To quantify changes in GFC frequency in MCI with respect to the HC group, we bin-wise subtracted each MCI CR+ subject's GFC frequencies from the averaged GFC frequencies in the HC CR+ group. The analogous subtraction was done for the MCI CR-, where each MCI subject's histogram was subtracted from the average histogram of

the HC CR- group. Thus, for each MCI CR group, alterations of GFC frequencies (called GFC-Diff, Figure 2C) were obtained according to the following equation.

$$GFC-Diff_{ijk} = GFC\ frequency(MCI)_{ijk} - Mean\ GFC\ frequency(HC)_{ik} \quad (Eq. 1)$$

where, i = CR group (CR + or CR-), j = MCI subject, k = GFC bin (1-60).

In bins where a MCI subject had a higher GFC frequency than the HC group, GFC-Diff values were positive (green shaded area in Figures 2C-E). Conversely, in bins where a MCI subject showed a lower GFC frequency compared to the HC group, the GFC-Diff score was negative (red shaded area in Figures 2C-2E). To identify GFC bins where MCI CR+ and MCI CR- subjects showed different GFC frequency changes, we compared GFC-Diff scores between the CR groups for each of the 60 bins, using two-sample t-tests with the significance threshold being $\alpha = 0.05$ for each t-test (Figure 2D). We did not correct for multiple testing at this stage, since the analysis was an intermediate step, exclusively done in order to select bins where MCI CR- and MCI CR+ groups differed in terms of GFC-Diff.

The results of the t-tests showed that GFC-Diff scores were greater (i.e. more positive) in MCI CR+ compared to MCI CR- in the range from 0.34 to 0.5, suggesting that MCI CR+ had significantly increased frequencies of relatively high GFC values (henceforth referred to as $GFC-Diff_{CR+>CR-}$) relative to MCI CR- subjects. In contrast, GFC-Diff scores were increased in MCI CR- compared to MCI CR+ subjects in a range from 0.2 to 0.26, suggesting that MCI CR- subjects had a higher frequency of relatively low GFC values (henceforth referred to as $GFC-Diff_{CR+<CR-}$) compared to MCI CR+ subjects. In order to create a subject-specific summary score of GFC frequency differences indicative of CR+ status, we subtracted the sum of GFC-Diff values in the $GFC-Diff_{CR+<CR-}$ from the sum of GFC-Diff values in the $GFC-Diff_{CR+>CR-}$. Finally, this difference was divided by the total number of voxels

in the cognitive control network mask to standardize it to a range between -1 and 1 (Figure 2E & F, Eq. 2).

$$GFC-R\ index_j = \frac{\sum GFC-Diff_{CR+>CR-} - \sum GFC-Diff_{CR+<CR-}}{\text{Number of voxels in mask}} \quad (Eq. 2)$$

where j = subject

This coefficient was then used as our GFC-R index. A negative GFC-R index indicates an increased GFC frequency in $GFC-Diff_{CR+<CR-}$ and a simultaneous decrease in $GFC-Diff_{CR+>CR-}$, i.e. a MCI CR- characteristic pattern. Conversely a positive GFC-reserve indicates an increased GFC frequency in $GFC-Diff_{CR+>CR-}$ and a decreased frequency in $GFC-Diff_{CR+<CR-}$, a pattern that was typically seen in MCI CR+.

All steps described above were conducted also for the test sample (ISD). Supplementary Figure 1 is showing – equivalent to Figure 2B - the distribution of GFC voxels averaged across subjects within CR and diagnostic (MCI vs. HC) groups. When conducting the t-tests to compare the GFC-Diff values between MCI CR+ and MCI CR-, we found $GFC-Diff_{CR+<CR-}$ in a range from 0.2-0.22 (vs. 0.2-0.26 in the training sample) and the $GFC-Diff_{CR+>CR-}$ in a range from 0.29-0.41 (vs. 0.34-0.5 in the training sample). The $GFC-Diff_{CR+<CR-}$ fully overlapped between both samples, whereas the $GFC-Diff_{CR+>CR-}$ only partly overlapped. For our validation analysis, we used the $GFC-Diff_{CR+>CR-}$ and $GFC-Diff_{CR+<CR-}$ ranges derived from the training sample to compute the GFC-R index in the test sample. All above delineated steps were conducted accordingly for 7 major brain networks derived from a previous publication to control whether a relationship between CR and GFC was specific for the cognitive control network. Again, $GFC-Diff_{CR+>CR-}$ and $GFC-Diff_{CR+<CR-}$ ranges were assessed in the training sample and used to create the GFC-R index in the test sample. The histogram analysis was conducted fully-automated using in-house MATLAB scripts.

Statistical Analysis:

Demographic variables were compared between groups using t-tests for continuous variables and χ^2 -test for gender.

In order to test whether the GFC-R index differed between MCI CR+ vs. MCI CR- groups in the training sample, we conducted an ANCOVA, with group as the predictor, and age, gender, the global AV45 uptake, the grey matter volume within the cognitive control network and the learning score of the RAVLT as covariates. To evaluate how accurately the GFC-R index classified between MCI CR+ and MCI CR- subjects we performed a Receiver Operating Characteristic (ROC) Curve analysis. Prediction accuracy was quantified using the area under the curve (AUC). The 95% Confidence interval (CI) for each ROC was computed with 2000 stratified bootstrap replicates for each ROC analysis. Equivalent models were run in the test sample, with the exception of AV45 PET uptake, which was not available in the ISD test sample.

Lastly, we tested whether the GFC-R index predicted the CR proxies (years of education, CRiQ) in the MCI subjects of the test sample (pooled across CR+ and CR-). To this end we conducted a multiple regression analysis, with the GFC-R index as a predictor of years of education or the CRiQ, controlled for age, gender, the learning score of the CERAD and the total grey matter volume within the cognitive control network. For the ADNI sample, the association between the continuous AV-45 PET measure and GFC-R was tested in the MCI subjects (who were by definition of the inclusion criteria all AV-45 PET positive). We conducted a linear regression analysis, with AV45 uptake as independent variable and the GFC-R index as dependent variable, and age, gender, the RAVLT learning score and grey matter volume as covariates. Next, we tested whether our findings on the GFC-R for the prediction of years of education were specific for the cognitive control network. Thus, the regression analyses on GFC-R were repeated for each GFC-R index derived on the GFC

frequencies in one of 7 major functional brain networks (i.e. Default Mode Network (DMN), Visual Network, Somatomotor Network, Dorsal Attention Network (DAN), Ventral Attention Network (VAN), Limbic Network, Frontoparietal Network (FPAN)) (Yeo et al., 2011).

All statistical analyses were conducted using the statistical software package *R* (R Development Core Team, 2013). Linear models were computed using the `lm` command in *R*. Linear model assumptions (skewness, kurtosis, heteroscedasticity) were tested using the `gvlma` function implemented in *R*. For all models reported, no significant ($\alpha = 0.05$) violations of linear model assumptions were found.

RESULTS:

Demographics, cognitive measures and the mean GFC-reserve index values for the training and test sample are depicted in table 1.

GFC Distribution:

Figure 3 shows the spatial distribution of significant GFC values in the brain displayed in percentiles for the training (ADNI) and the test sample (ISD). We found a high spatial correspondence of significant GFC values between both samples with a correlation coefficient of $r = 0.84$, $p < 0.001$. The highest GFC values were observed predominantly within the frontal cortex, lateral parietal cortex, and areas of the medial brain surface. Those brain areas are known to be part of the DMN and the cognitive control network as reported previously (Cole et al., 2010).

The GFC-reserve index is decreased in MCI CR- as compared to MCI CR+:
MCI CR- showed significantly lower GFC-R index values than the MCI CR+ subjects in the training sample ($F(6,36) = 16.82$, $p = 0.0002$; see Figure 4A) and the test sample ($F(5,17) = 7.93$, $p = 0.0119$). In an exploratory regression analysis, we tested the association between the

global AV45 uptake and the GFC-R index of the cognitive control network with age, gender, grey matter volume of the cognitive control network and the RAVLT learning score as covariates. The model showed no relationship between AV45 and the GFC-R index ($t(37) = -0.1482, p = 0.54$).

ROC analysis:

Using a ROC analysis, we evaluated how accurate the GFC-reserve index discriminated between MCI CR+ and MCI CR- subjects (Figure 4B). The AUC was 0.840 with the 95% CI ranging between 0.72 and 0.95 within the training sample. Similarly, in the test sample, we found a AUC of 0.79 with the CI ranging from 0.60-0.99.

The GFC-reserve index is a predictor of CR proxies in the ISD test sample:

Using linear regression, we tested whether the GFC-R index predicted CR proxies in the test sample, when controlling for age, gender, the word list learning score of the CERAD battery and grey matter volume of the cognitive control network. For years of education, the regression model was significant ($F(5,17) = 12.63, p = 0.00003$) with an adjusted R^2 of 0.72, showing that a higher GFC-R index significantly predicted higher years of education ($t(17) = 2.232, p = 0.039$). For the CRIq score, a higher GFC-R index predicted a higher CRIq score ($t(17) = 2.638, p = 0.173$, overall model fit: $F(5,17) = 4.422, p = 0.009$, adjusted R^2 0.44). The relationship between the GFC-R index and the CR proxies is illustrated in Figure 5. When testing the Pearson-moment correlation between the GFC-R index and our CR proxies, the correlation was significant for both years of education ($r = 0.46, p = 0.026$) and the CRIq ($r = 0.6, p = 0.0024$).

Control analyses in other brain networks:

1 In order to test, whether our findings on the prediction of years of education by GFC-R index
2 were specific for the cognitive control network, we repeated the regression analysis for GFC-
3 R index derived from each of seven other major cortical networks (Yeo et al., 2011). For none
4 of the other networks, the GFC-R index predicted years of education or the CRIq ($p > 0.05$,
5 table 2). This suggests that the relationship between GFC changes and education is specific
6 for the cognitive control network.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

DISCUSSION:

The first major finding of the current study was that MCI subjects with high CR (as measured by years of education) had an increased frequency of high GFC values within the cognitive control network compared to MCI subjects with lower years of education. Secondly, a newly derived summary measure of abnormal GFC frequencies within the cognitive control network, the GFC-R index, was associated with more years of education and a higher CRIq score, a composite measure of CR, in an independent cross-validation sample of MCI patients. The predictive value of GFC-R index was independent of demographic variables including age and gender, episodic memory performance or grey matter volume of the cognitive control network. These results suggest that the GFC-R index constitutes a biomarker candidate of CR-related functional brain changes in MCI.

For our first major finding, MCI CR+ showed a right-ward shift of the GFC histogram to that in HC CR+, i.e. MCI CR+ showed an increased frequency of relatively high GFC values. In contrast, there was a left-ward shift of the GFC histogram in the MCI CR- group, i.e. an increased frequency of lower GFC values. A previous study on GFC changes in MCI reported decreased GFC in the frontal, parietal, and temporal cortices in MCI (J. Wang et al., 2013). That latter study, however, did not assess the impact of years of education on GFC differences. Our results extend those previous results showing that the levels of CR are an important modifying factor, where MCI CR- subjects show a decrease in GFC but MCI CR+ subjects show an increase in GFC within the cognitive control network.

The increase in the frequency of high GFC values in MCI CR+ may reflect either pre-existing high levels of GFC before the development of MCI or, alternatively, a compensatory increase in GFC during the development of MCI. Previous studies showed that higher IQ is associated with higher GFC within the left frontal core region of the cognitive control network in young subjects (Cole, Yarkoni, Repovs, Anticevic, & Braver, 2012). Given that years of education and IQ are correlated (Matarazzo & Hermann, 1984), it is possible that MCI CR+ subjects

1 had already higher levels of GFC before disease onset, thus possessing higher brain reserve.
2 However, the fact that MCI CR+ subjects showed abnormally increased frequency of high
3 GFC values when compared to HC CR+, i.e. at similarly high levels of education, suggests a
4 compensatory increase of GFC in MCI. Such an interpretation of compensatory increase of
5 GFC in MCI is consistent with several previous studies showing increased resting-state
6 functional connectivity in MCI and AD compared to HC (K. Wang et al., 2007), that is
7 attributable to higher levels of education (Bozzali et al., 2015). In summary, the MCI CR+
8 subjects show increased frequency of high GFC values within the cognitive control network,
9 which probably reflects compensatory changes in MCI.
10

11 For our second major finding, we could show in a cross-validation approach that higher levels
12 of the GFC-R index are predictive of higher levels of education and CRIq in MCI and could
13 well separate high vs. low education groups in MCI as shown by the ROC analysis, with an
14 AUC of 0.79. Note however that sensitivity and specificity is not of primary clinical
15 significance in the context of CR, which is likely to be continuously distributed. More
16 importantly, the GFC-R index significantly predicted the level of CR proxies in the validation
17 sample. The point prediction is difficult but clinically important as previous studies showed
18 that with each additional year of education, the onset of dementia is delayed by 0.21 years
19 (Hall et al., 2007) and the risk of AD dementia is reduced (Sando et al., 2008; Stern et al.,
20 1994). A critical test in the future will be whether the GFC-R index predicts slower cognitive
21 decline in subjects with preclinical AD or MCI as has been reported for years of education as
22 a proxy of CR (Soldan et al., 2015). The advantage of using fMRI based CR biomarkers such
23 as GFC-R in such prediction models is that GFC-R could be used as a measure to track CR
24 changes over time. CR may be reduced as the disease progresses since brain pathology may
25 eventually use up the reserve (Members et al., 2010). In contrast, proxies of CR such as
26 education or occupational attainment are time-invariant.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 We found that only the GFC-R index derived from GFC values within the cognitive control
2 network but not within any of the other major resting-state networks was predictive of years
3 of education or CRIq. These results suggest that frequency changes of GFC specifically
4 within the cognitive control network are related to CR. These results are broadly consistent
5 with previous findings showing that higher GFC of brain regions in the cognitive control
6 network but not the default mode network were predictive of higher IQ in healthy subjects
7 (Cole et al., 2012). A possible explanation includes that the cognitive control network has a
8 unique role in the brain, such that it is highly connected with the other networks and may
9 orchestrate the activation of other networks during cognitive tasks (Cole, Repovs, &
10 Anticevic, 2014; Cole et al., 2013). Brain regions with increased connectedness in the brain
11 have previously shown to be more resilient to targeted attacks as shown in graph theoretical
12 analysis of resting state fMRI (Achard, Salvador, Whitcher, Suckling, & Bullmore, 2006).
13 Higher GFC of the cognitive control network may enable to more flexibly activate different
14 networks during cognitive processing (Cole et al., 2013), which in neurodegenerative disease
15 may render a more flexible coping with local damage of specific neural networks, thus
16 increasing CR. This will need to be tested in future combined resting-state and task-related
17 fMRI studies.

18 We used years of education as our primary outcome measure, i.e. the gold standard, since
19 educational attainment has been recommended as the best validated indicator of cognitive
20 reserve (Stern, 2012). Years of education has been tested as a CR proxy in numerous studies
21 in AD (for review see (Stern, 2012)) and is robustly associated with reduced risk of AD
22 dementia across studies (Meng & D'Arcy, 2012; Valenzuela & Sachdev, 2006). Alternative
23 proxy measures of CR include assessments such as occupational attainment, premorbid IQ or
24 leisure activities. Since we used international cross-validation samples, equivalent measures
25 of such variables were not available in both samples in the current study. However, we found
26 a positive association between the GFC-R index and the CRIq (Nucci et al., 2012) an

alternative CR proxy that takes into account education, working and leisure activities, supporting criterion validity of the GFC-R index.

A promising alternative marker of CR has been recently proposed, consisting of the residual episodic memory variability after accounting for brain atrophy and demographic variables (Reed et al., 2010; Zahodne et al., 2015; Zahodne et al., 2013). Such a measure captures well CR as the discrepancy between the level of cognitive performance and brain pathology, but is non-informative about any structural or functional brain changes that may underlie CR. The current biomarker captures functional brain changes of CR and would thus be complimentary to such memory-variance based marker or any of the standard proxy measures of CR.

For the interpretation of the current results several caveats need to be taken into account. It is important to note that the GFC-R index is not a biomarker candidate of CR *per se*, rather it is a biomarker of functional brain changes that underlie CR. Ideally, the primary outcome parameter for the validation of the current biomarker constitute specific functional mechanisms that cause CR. Although several task fMRI studies have attempted to extract specific functional brain changes of CR in subjects with MCI and AD, no core mechanism, however, has yet emerged (for review see (Barulli & Stern, 2013)). Thus, more work is needed to disentangle the functional brain processes that underlie CR, which could then provide a point of reference for the validation of functional biomarkers of CR. Still, years of education has been validated in numerous studies as a marker of CR and may thus constitute the best primary outcome as a reference measures for the validation of functional biomarkers of CR at this point.

It should be also taken into account that the reliability of GFC assessment is an important factor for the utility of GFC-R as a biomarker. Previous studies showed that GFC exhibits a fair to excellent test-retest reliability and its retest reliability ranks among the highest of resting state fMRI functional connectivity measures (Liao et al., 2013; J. H. Wang et al., 2011). Multicenter variability of resting state fMRI is an active field of research and needs

1 still to be established for the various connectivity indices including that of GFC. However,
2 the current cross-validation of the GFC-R between completely different samples using
3 different scanners and acquisition protocols suggest robustness of the current findings (Feis et
4 al., 2015). Summary indices that average across large number of voxels may be more robust
5 to multicenter variability that measures focusing on small ROIs (Ewers et al., 2006). Still, the
6 test-retest and multicenter variability of the GFC-R index needs to be established in future
7 studies.
8

9 A strength of the current approach is the fully automated way to extract GFC frequency
10 changes based on resting state fMRI. Thus, functional MRI data can be assessed without
11 reliance on a task and data processing can be done without manual intervention, which
12 provides a high attractiveness to be used in clinical praxis. Possible clinical applications of the
13 GFC-R index as diagnostic biomarker candidate include the use outcome measure in clinical
14 trials such as cognitive training, physical training that target compensatory brain mechanisms
15 (Suo et al., 2016). Secondly, the GFC-R index could be used to track changes in CR during
16 the progression of the disease. Future longitudinal studies may address these next steps.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Funding

The research was funded by grants of the LMUexcellent Initiative and the European Commission (ERC, PCIG12-GA-2012-334259), Alzheimer's Forschung Initiative (AFI, DE-15035). Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

Conflict of Interest Statement:

N. Franzmeier has nothing to disclose

M. Á. Araque Caballero has nothing to disclose

A. Taylor has nothing to disclose

L. Simon-Vermot has nothing to disclose

K. Buerger received funding and honoraria from the German Center for Neurodegenerative Diseases (DZNE).

B. Ertl-Wagner has received paid speakerships and/or honoraria for manuscript preparations from Siemens, Philips, Bracco, Roche, Novartis and Bayer Vital; she has been a member of advisory boards for Philips and Bracco.

C. Mueller has nothing to disclose

C. Catak has nothing to disclose

D. Janowitz has nothing to disclose

E. Baykara has nothing to disclose

B. Gesierich has nothing to disclose

M. Duering received travel expenses for lectures from the United Leukodystrophy Foundation, the European Stroke Conference and Kenes International. he receives research support from the German Center for Neurodegenerative Diseases (DZNE).

M. Ewers received funding from LMUexcellent, European Research Council (ERC) as well as the Alzheimer Forschung Initiative (AFI), and honoraria from the Deutsche Gesellschaft für Neurologie (DGN).

The authors declare that they have no conflict of interest

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES:

- Achard, S., Salvador, R., Whitcher, B., Suckling, J., & Bullmore, E. (2006). A resilient, low-frequency, small-world human brain functional network with highly connected association cortical hubs. *J Neurosci*, 26(1), 63-72. doi:10.1523/JNEUROSCI.3874-05.2006
- Arenaza-Urquijo, E. M., Landeau, B., La Joie, R., Mevel, K., Mezenge, F., Perrotin, A., . . . Chetelat, G. (2013). Relationships between years of education and gray matter volume, metabolism and functional connectivity in healthy elders. *Neuroimage*, 83, 450-457. doi:10.1016/j.neuroimage.2013.06.053
- Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *Neuroimage*, 38(1), 95-113. doi:10.1016/j.neuroimage.2007.07.007
- Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *Neuroimage*, 26(3), 839-851. doi:10.1016/j.neuroimage.2005.02.018
- Barulli, D., & Stern, Y. (2013). Efficiency, capacity, compensation, maintenance, plasticity: emerging concepts in cognitive reserve. *Trends Cogn Sci*, 17(10), 502-509. doi:10.1016/j.tics.2013.08.012
- Bastin, C., Yakushev, I., Bahri, M. A., Fellgiebel, A., Eustache, F., Landeau, B., . . . Salmon, E. (2012). Cognitive reserve impacts on inter-individual variability in resting-state cerebral metabolism in normal aging. *NeuroImage*, 63(2), 713-722. doi:10.1016/j.neuroimage.2012.06.074
- Boots, E. A., Schultz, S. A., Almeida, R. P., Oh, J. M., Kosciak, R. L., Dowling, M. N., . . . Okonkwo, O. C. (2015). Occupational Complexity and Cognitive Reserve in a Middle-Aged Cohort at Risk for Alzheimer's Disease. *Arch Clin Neuropsychol*. doi:10.1093/arclin/acv041
- Bosch, B., Bartres-Faz, D., Rami, L., Arenaza-Urquijo, E. M., Fernandez-Espejo, D., Junque, C., . . . Molinuevo, J. L. (2010). Cognitive reserve modulates task-induced activations and deactivations in healthy elders, amnesic mild cognitive impairment and mild Alzheimer's disease. *Cortex*, 46(4), 451-461. doi:10.1016/j.cortex.2009.05.006
- Bozzali, M., Dowling, C., Serra, L., Spano, B., Torso, M., Marra, C., . . . Cercignani, M. (2015). The impact of cognitive reserve on brain functional connectivity in Alzheimer's disease. *J Alzheimers Dis*, 44(1), 243-250. doi:10.3233/JAD-141824
- Brickman, A. M., Siedlecki, K. L., Muraskin, J., Manly, J. J., Luchsinger, J. A., Yeung, L. K., . . . Stern, Y. (2011). White matter hyperintensities and cognition: testing the reserve hypothesis. *Neurobiol Aging*, 32(9), 1588-1598. doi:10.1016/j.neurobiolaging.2009.10.013
- Buschert, V. C., Frieze, U., Teipel, S. J., Schneider, P., Merensky, W., Rujescu, D., . . . Buerger, K. (2011). Effects of a newly developed cognitive intervention in amnesic mild cognitive impairment and mild Alzheimer's disease: a pilot study. *J Alzheimers Dis*, 25(4), 679-694. doi:10.3233/JAD-2011-100999
- Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and task-evoked network architectures of the human brain. *Neuron*, 83(1), 238-251. doi:10.1016/j.neuron.2014.05.014
- Cole, M. W., Pathak, S., & Schneider, W. (2010). Identifying the brain's most globally connected regions. *NeuroImage*, 49(4), 3132-3148. doi:10.1016/j.neuroimage.2009.11.001
- Cole, M. W., Repovs, G., & Anticevic, A. (2014). The frontoparietal control system: a central role in mental health. *Neuroscientist*, 20(6), 652-664. doi:10.1177/1073858414525995
- Cole, M. W., Reynolds, J. R., Power, J. D., Repovs, G., Anticevic, A., & Braver, T. S. (2013). Multi-task connectivity reveals flexible hubs for adaptive task control. *Nat Neurosci*, 16(9), 1348-1355. doi:10.1038/nn.3470

- 1 Cole, M. W., & Schneider, W. (2007). The cognitive control network: Integrated cortical
2 regions with dissociable functions. *Neuroimage*, 37(1), 343-360.
3 doi:10.1016/j.neuroimage.2007.03.071
- 4 Cole, M. W., Yarkoni, T., Repovs, G., Anticevic, A., & Braver, T. S. (2012). Global
5 connectivity of prefrontal cortex predicts cognitive control and intelligence. *J*
6 *Neurosci*, 32(26), 8988-8999. doi:10.1523/JNEUROSCI.0536-12.2012
- 7 Ewers, M., Brendel, M., Rizk-Jackson, A., Rominger, A., Bartenstein, P., Schuff, N., . . .
8 Alzheimer's Disease Neuroimaging, I. (2014). Reduced FDG-PET brain metabolism
9 and executive function predict clinical progression in elderly healthy subjects.
10 *Neuroimage Clin*, 4, 45-52. doi:10.1016/j.nicl.2013.10.018
- 11 Ewers, M., Teipel, S. J., Dietrich, O., Schonberg, S. O., Jessen, F., Heun, R., . . . Hampel, H.
12 (2006). Multicenter assessment of reliability of cranial MRI. *Neurobiol Aging*, 27(8),
13 1051-1059. doi:10.1016/j.neurobiolaging.2005.05.032
- 14 Feis, R. A., Smith, S. M., Filippini, N., Douaud, G., Dopper, E. G., Heise, V., . . . Mackay, C.
15 E. (2015). ICA-based artifact removal diminishes scan site differences in multi-center
16 resting-state fMRI. *Front Neurosci*, 9, 395. doi:10.3389/fnins.2015.00395
- 17 Hall, C. B., Derby, C., LeValley, A., Katz, M. J., Verghese, J., & Lipton, R. B. (2007).
18 Education delays accelerated decline on a memory test in persons who develop
19 dementia. *Neurology*, 69(17), 1657-1664. doi:10.1212/01.wnl.0000278163.82636.30
- 20 Landau, S. M., Breault, C., Joshi, A. D., Pontecorvo, M., Mathis, C. A., Jagust, W. J., . . .
21 Alzheimer's Disease Neuroimaging, I. (2013). Amyloid-beta imaging with Pittsburgh
22 compound B and florbetapir: comparing radiotracers and quantification methods. *J*
23 *Nucl Med*, 54(1), 70-77. doi:10.2967/jnumed.112.109009
- 24 Liao, X. H., Xia, M. R., Xu, T., Dai, Z. J., Cao, X. Y., Niu, H. J., . . . He, Y. (2013).
25 Functional brain hubs and their test-retest reliability: a multiband resting-state
26 functional MRI study. *NeuroImage*, 83, 969-982.
27 doi:10.1016/j.neuroimage.2013.07.058
- 28 Luck, T. R.-H. S. W. B. (2009). CERAD-NP-Testbatterie: Alters-, geschlechts- und
29 bildungsspezifischen Normen ausgewählter Subtests. *Z Gerontol Geriatr*, 42, 372-384.
- 30 Matarazzo, J. D., & Hermann, D. O. (1984). The relationship of education and IQ in the
31 WAIS--R standardization sample. *Journal of Consulting and Clinical Psychology*,
32 52(4), 631-634.
- 33 Members, E. C. C., Brayne, C., Ince, P. G., Keage, H. A., McKeith, I. G., Matthews, F. E., . . .
34 Sulkava, R. (2010). Education, the brain and dementia: neuroprotection or
35 compensation? *Brain*, 133(Pt 8), 2210-2216. doi:10.1093/brain/awq185
- 36 Meng, X., & D'Arcy, C. (2012). Education and dementia in the context of the cognitive
37 reserve hypothesis: a systematic review with meta-analyses and qualitative analyses.
38 *PLoS One*, 7(6), e38268. doi:10.1371/journal.pone.0038268
- 39 Nucci, M., Mapelli, D., & Mondini, S. (2012). Cognitive Reserve Index questionnaire (CRIq):
40 a new instrument for measuring cognitive reserve. *Aging Clin Exp Res*, 24(3), 218-
41 226. doi:10.3275/7800
- 42 Petersen, R. C. (2004). Mild cognitive impairment as a diagnostic entity. *J Intern Med*,
43 256(3), 183-194. doi:10.1111/j.1365-2796.2004.01388.x
- 44 R Development Core Team. (2013). R: A language and environment for statistical computing.
45 Vienna, Austria: R Foundation for Statistical Computing.
- 46 Reed, B. R., Mungas, D., Farias, S. T., Harvey, D., Beckett, L., Widaman, K., . . . DeCarli, C.
47 (2010). Measuring cognitive reserve based on the decomposition of episodic memory
48 variance. *Brain*, 133(Pt 8), 2196-2209. doi:10.1093/brain/awq154
- 49 Reijnders, J., van Heugten, C., & van Boxtel, M. (2013). Cognitive interventions in healthy
50 older adults and people with mild cognitive impairment: a systematic review. *Ageing*
51 *Res Rev*, 12(1), 263-275. doi:10.1016/j.arr.2012.07.003

- 1 Rentz, D. M., Locascio, J. J., Becker, J. A., Moran, E. K., Eng, E., Buckner, R. L., . . .
2 Johnson, K. A. (2010). Cognition, reserve, and amyloid deposition in normal aging.
3 *Ann Neurol*, 67(3), 353-364. doi:10.1002/ana.21904
- 4 Sando, S. B., Melquist, S., Cannon, A., Hutton, M., Sletvold, O., Saltvedt, I., . . . Aasly, J.
5 (2008). Risk-reducing effect of education in Alzheimer's disease. *Int J Geriatr*
6 *Psychiatry*, 23(11), 1156-1162. doi:10.1002/gps.2043
- 7 Scarmeas, N., Zarahn, E., Anderson, K. E., Habeck, C. G., Hilton, J., Flynn, J., . . . Stern, Y.
8 (2003). Association of life activities with cerebral blood flow in Alzheimer disease:
9 implications for the cognitive reserve hypothesis. *Arch Neurol*, 60(3), 359-365.
- 10 Schoenberg, M. R., Dawson, K. A., Duff, K., Patton, D., Scott, J. G., & Adams, R. L. (2006).
11 Test performance and classification statistics for the Rey Auditory Verbal Learning
12 Test in selected clinical samples. *Arch Clin Neuropsychol*, 21(7), 693-703.
13 doi:10.1016/j.acn.2006.06.010
- 14 Schultz, S. A., Larson, J., Oh, J., Kosciak, R., Dowling, M. N., Gallagher, C. L., . . . Okonkwo,
15 O. C. (2015). Participation in cognitively-stimulating activities is associated with brain
16 structure and cognitive function in preclinical Alzheimer's disease. *Brain Imaging*
17 *Behav*, 9(4), 729-736. doi:10.1007/s11682-014-9329-5
- 18 Soldan, A., Pettigrew, C., Lu, Y., Wang, M. C., Selnes, O., Albert, M., . . . Team, B. R.
19 (2015). Relationship of medial temporal lobe atrophy, APOE genotype, and cognitive
20 reserve in preclinical Alzheimer's disease. *Human brain mapping*, 36(7), 2826-2841.
21 doi:10.1002/hbm.22810
- 22 Solé-Padullés, C., Bartrés-Faz, D., Junqué, C., Vendrell, P., Rami, L., Clemente, I. C., . . .
23 Molinuevo, J. L. (2009). Brain structure and function related to cognitive reserve
24 variables in normal aging, mild cognitive impairment and Alzheimer's disease.
25 *Neurobiology of Aging*, 30(7), 1114-1124.
- 26 Stern, Y. (2002). What is cognitive reserve? Theory and research application of the reserve
27 concept. *J Int Neuropsychol Soc*, 8(3), 448-460.
- 28 Stern, Y. (2009). Cognitive reserve. *Neuropsychologia*, 47(10), 2015-2028. doi:S0028-
29 3932(09)00123-7 [pii]
30 10.1016/j.neuropsychologia.2009.03.004
- 31 Stern, Y. (2012). Cognitive reserve in ageing and Alzheimer's disease. *The Lancet Neurology*,
32 11(11), 1006-1012. doi:10.1016/s1474-4422(12)70191-6
- 33 Stern, Y., Alexander, G. E., Prohovnik, I., & Mayeux, R. (1992). Inverse relationship between
34 education and parietotemporal perfusion deficit in Alzheimer's disease. *Ann Neurol*,
35 32(3), 371-375. doi:10.1002/ana.410320311
- 36 Stern, Y., Alexander, G. E., Prohovnik, I., Stricks, L., Link, B., Lennon, M. C., & Mayeux, R.
37 (1995). Relationship between lifetime occupation and parietal flow: implications for a
38 reserve against Alzheimer's disease pathology. *Neurology*, 45(1), 55-60.
- 39 Stern, Y., Gurland, B., Tatemichi, T. K., Tang, M. X., Wilder, D., & Mayeux, R. (1994).
40 Influence of education and occupation on the incidence of Alzheimer's disease. *Jama*,
41 271(13), 1004-1010.
- 42 Stern, Y., Habeck, C., Moeller, J., Scarmeas, N., Anderson, K. E., Hilton, H. J., . . . van
43 Heertum, R. (2005). Brain Networks Associated with Cognitive Reserve in Healthy
44 Young and Old Adults. *Cereb. Cortex*, 15(4), 394-402. doi:10.1093/cercor/bhh142
- 45 Stern, Y., Zarahn, E., Habeck, C., Holtzer, R., Rakitin, B. C., Kumar, A., . . . Brown, T.
46 (2008). A Common Neural Network for Cognitive Reserve in Verbal and Object
47 Working Memory in Young but not Old. *Cerebral Cortex*, 18(4), 959-967.
48 doi:10.1093/cercor/bhm134
- 49 Suo, C., Singh, M. F., Gates, N., Wen, W., Sachdev, P., Brodaty, H., . . . Valenzuela, M. J.
50 (2016). Therapeutically relevant structural and functional mechanisms triggered by
51 physical and cognitive exercise. *Mol Psychiatry*. doi:10.1038/mp.2016.19

- 1 Valenzuela, M. J., & Sachdev, P. (2006). Brain reserve and dementia: a systematic review.
2 *Psychol Med*, 36(4), 441-454. doi:10.1017/S0033291705006264
- 3 Vemuri, P., Lesnick, T. G., Przybelski, S. A., Knopman, D. S., Preboske, G. M., Kantarci, K.,
4 . . . Jack, C. R., Jr. (2015). Vascular and amyloid pathologies are independent
5 predictors of cognitive decline in normal elderly. *Brain*, 138(Pt 3), 761-771.
6 doi:10.1093/brain/awu393
- 7 Vemuri, P., Weigand, S. D., Przybelski, S. A., Knopman, D. S., Smith, G. E., Trojanowski, J.
8 Q., . . . Alzheimer's Disease Neuroimaging, I. (2011). Cognitive reserve and
9 Alzheimer's disease biomarkers are independent determinants of cognition. *Brain*,
10 134(Pt 5), 1479-1492. doi:10.1093/brain/awr049
- 11 Wang, J., Zuo, X., Dai, Z., Xia, M., Zhao, Z., Zhao, X., . . . He, Y. (2013). Disrupted
12 functional brain connectome in individuals at risk for Alzheimer's disease. *Biol*
13 *Psychiatry*, 73(5), 472-481. doi:10.1016/j.biopsych.2012.03.026
- 14 Wang, J. H., Zuo, X. N., Gohel, S., Milham, M. P., Biswal, B. B., & He, Y. (2011). Graph
15 theoretical analysis of functional brain networks: test-retest evaluation on short- and
16 long-term resting-state functional MRI data. *PLoS One*, 6(7), e21976.
17 doi:10.1371/journal.pone.0021976
- 18 Wang, K., Liang, M., Wang, L., Tian, L., Zhang, X., Li, K., & Jiang, T. (2007). Altered
19 functional connectivity in early Alzheimer's disease: A resting-state fMRI study. *Hum*
20 *Brain Mapp*, 28(10), 967-978. doi:10.1002/hbm.20324
- 21 Wells, R. E., Yeh, G. Y., Kerr, C. E., Wolkin, J., Davis, R. B., Tan, Y., . . . Kong, J. (2013).
22 Meditation's impact on default mode network and hippocampus in mild cognitive
23 impairment: a pilot study. *Neurosci Lett*, 556, 15-19. doi:10.1016/j.neulet.2013.10.001
- 24 Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-
25 scale automated synthesis of human functional neuroimaging data. *Nat Methods*, 8(8),
26 665-670. doi:10.1038/nmeth.1635
- 27 Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., . . .
28 Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by
29 intrinsic functional connectivity. *J Neurophysiol*, 106(3), 1125-1165.
30 doi:10.1152/jn.00338.2011
- 31 Zahodne, L. B., Manly, J. J., Brickman, A. M., Narkhede, A., Griffith, E. Y., Guzman, V. A.,
32 . . . Stern, Y. (2015). Is residual memory variance a valid method for quantifying
33 cognitive reserve? A longitudinal application. *Neuropsychologia*, 77, 260-266.
34 doi:10.1016/j.neuropsychologia.2015.09.009
- 35 Zahodne, L. B., Manly, J. J., Brickman, A. M., Siedlecki, K. L., Decarli, C., & Stern, Y.
36 (2013). Quantifying cognitive reserve in older adults by decomposing episodic
37 memory variance: replication and extension. *J Int Neuropsychol Soc*, 19(8), 854-862.
38 doi:10.1017/S1355617713000738

Figure legends:

Fig. 1: Meta-analytical activation map across 428 task-fMRI studies that were associated with search term “cognitive control” (FDR-corrected at $p < 0.01$) in NeuroSynth, projected on a brain surface. Colors indicate z-scores.

Fig. 2: Analysis flow diagram, illustrating the steps of GFC-R index computation. (A) Voxel-wise GFC is computed based on preprocessed resting-state fMRI for each subject and masked with the binarized cognitive control network map. (B) The GFC frequency distribution within the cognitive control network is plotted for groups split by diagnosis (HC & MCI) and CR status (CR- & CR+). (C) GFC within the cognitive control network is binned in intervals of 0.01 for each subject. Within each CR group, the difference in GFC differences (GFC-Diff) between each MCI subject and the average GFC within the HCs group is computed. Colored areas indicate whether MCI subjects showed lower (red) or higher (green) GFC frequency than the HC subjects. (D) GFC-Diff scores are compared between MCI CR+ and MCI CR- groups via bin-wise two-sample t-tests. (E) GFC-Diff scores are summed up across the selected bins for each MCI subject. In order to create a subject-specific summary score of GFC frequency differences indicative of CR+ status, the sum of GFC-Diff values in the GFC-Diff_{CR+<CR-} was subtracted from the sum of GFC-Diff values in the GFC-Diff_{CR+>CR-}. (F) This difference was divided by the total number of voxels in the cognitive control network mask to standardize it to a range between -1 and 1 to derive the GFC-R index for each MCI subject.

Fig. 3: Distribution of significant GFC values in the brain. T-values of voxel wise one-sample t-tests of the GFC among the pooled HC and MCI subjects (FWE corrected at the voxel level at $\alpha = 0.001$) were converted to percentiles to facilitate visual group comparison between both samples.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 4: (A) Boxplots of the GFC-R index split by CR group for the training and the test sample. MCI CR- subjects show significantly lower GFC-CR values as MCI CR+ subjects in both samples. (B) shows the ROC curves with the specificity on the x- and the sensitivity on the y-axis. AUC = Area under the curve, * = $p < 0.05$, ** = $p < 0.001$

Fig. 5: Scatterplot for the relationship between the GFC-R index and the CR proxies (years of education & CRIq) in the test sample.

Supplementary Fig. 1: Shown is the distribution of GFC voxels averaged across subjects within CR (CR+ vs. CR-) and diagnostic (MCI vs. HC) groups for the test sample. The graphs are equivalent to Figure 2B of the main manuscript.

Tables:

Table 1: Demographics and neuropsychological characteristics of the study samples subjects split by Diagnosis and CR group

Training sample (ADNI)				
	HC CR- (n = 13)	HC CR+ (n = 11)	MCI CR- (n = 24)	MCI CR+ (n = 19)
Age (years) ²	75.12 ± 5.85	74.30 ± 7.56	74.90 ± 5.87	69.10 ± 6.16
Gender (female/male)	3/10	5/6	14/10	12/7
Education ^{1,3}	15.15 ± 1.41	18.64 ± 1.12	14.17 ± 1.58	18.58 ± 1.02
Global AV45	0.99 ± 0.45	0.98 ± 0.04	1.4 ± 0.18	1.37 ± 0.15
Uptake				
MMSE ^{2,3}	29.12 ± 0.91	27.91 ± 1.45	26.71 ± 1.63	28.16 ± 1.34
RAVLT	45 ± 13.46	43.74 ± 7.81	31.71 ± 9.40	38.70 ± 8.91
Learning ¹				
Test sample (ISD)				
	HC CR- (n = 17)	HC CR+ (n = 15)	MCI CR- (n = 13)	MCI CR+ (n = 10)
Age (years)	70.17 ± 3.94	72.52 ± 6.33	77.02 ± 3.63	73.87 ± 4.23
Gender ⁴ (female/male)	13/4	5/10	11/2	8/2
Education ^{1,3}	11.59 ± 1.33	16.6 ± 2.1	10.92 ± 1.98	17.1 ± 2.08
MMSE ¹	29.53 ± 0.87	29.33 ± 0.72	25.15 ± 1.52	27.9 ± 2.33
CERAD Word	23 2.6	24.07 3.24	13.3 2.84	18.9 3.14
List Learning ¹				
1) MCI CR+ > MCI CR-				
2) MCI CR+ < MCI CR-				
3) HC CR+ > HC CR-				
4) HC CR+ < HC CR-				

Table 2: Control analyses of the GFC-R index as a predictor of CR proxies in major brain networks

Training sample (ADNI)			Test sample (ISD)			
Functional Network	<i>GFC-Diff</i>		<i>GFC-CR as a predictor of education in MCI CR+ and CR- pooled¹</i>		<i>GFC-CR as a predictor of CRIq in MCI CR+ and CR- pooled¹</i>	
	CR+<CR-	CR+>CR-	T	p	T	p
Cognitive Control	0.2-0.26	0.34-0.5	2.232	0.039	2.638	0.017
Default Mode	0.2-0.27	0.37-0.52	0.663	0.516	1.365	0.190
Dorsal Attention	0.22-0.28	0.41-0.51	0.981	0.341	0.948	0.356
Ventral Attention	0.19-0.23	0.3-0.5	0.927	0.367	1.820	0.087
Frontoparietal	0.2-0.27	0.35-0.52	1.258	0.225	1.656	0.116
Limbic	0.25	0.36-0.46	0.566	0.579	0.562	0.582
Visual	0.22-0.28	0.43-0.53	0.256	0.801	1.571	0.135
Somatomotor	0.21-0.28	0.32-0.46	1.2526	0.227	0.849	0.408

1) Models controlled for age, gender, grey matter volume of the tested network, CERAD Word list learning score

Meta-analysis: fMRI activation - cognitive control

Analysis flow diagram (training sample)

Training sample (ADNI)

Test sample (ISD)

Test sample (ISD)

Test sample (ISD)