Aberystwyth University

Recognising the potential role of native ponies in conservation management

Fraser, Mariecia; Stanley, Christina; Hegarty, Matthew

Published in:
Biological Conservation

DOI:
[10.1016/j.biocon.2019.04.014](https://doi.org/10.1016/j.biocon.2019.04.014)

Publication date:
2019

Citation for published version (APA):

Fraser, M., Stanley, C., & Hegarty, M. (2019). Recognising the potential role of native ponies in conservation management. *Biological Conservation*, 235, 112-118. <https://doi.org/10.1016/j.biocon.2019.04.014>

Document License CC BY-NC-ND

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Recognising the potential role of native ponies in conservation management

M.D. Fraser¹, C.R. Stanley² and M.J. Hegarty³

¹Pwllpeiran Upland Research Centre, Aberystwyth University, Cwmystwyth, Aberystwyth
SY23 4AB, UK

²Department of Biological Sciences, University of Chester, Parkgate Road, Chester, CH1 4BJ,
UK

³Institute of Biological, Environmental and Rural Sciences, Aberystwyth University,
Gogerddan, Aberystwyth SY23 3EB, UK

Running title: Native ponies and conservation

Abstract

Population control of feral horses has been the subject of public debate in many parts of the world in recent years due to wide-reaching ecological and societal impacts. However, the feral populations in these high-profile cases are not ‘native’ but are instead descended from animals which escaped from or were released by settlers. This paper considers i) the potential role of indigenous equids as conservation grazers within native ecosystems currently in poor condition, and ii) the value of supporting semi-wild native ponies specifically. We argue that the high ecological overlap between ponies and cattle reported in a range of studies means that they should be considered as alternative tools for conservation management, particularly in scenarios where there is a need to reduce the dominance of plant species avoided by more-selective small ruminants such as sheep. Semi-wild ponies could be particularly suited to conservation grazing because their genomes have been predominately shaped by natural and not artificial selection, meaning they may have adaptations no longer present in domesticated equids. With agricultural and environmental policy in the EU and UK under major review, it is anticipated that the wider delivery of public goods, rather than primary production, will be prioritised under future subsidy payment schemes. Recognising the value of native ponies as conservation grazers would broaden the range of routes by which land managers could achieve biodiversity gain, while simultaneously supporting at-risk equine genotypes.

Keywords: horses; equids; semi-wild; conservation grazing; agri-environment; genetic adaptation

1. Background

Grazing continues to be a major driver of land use change worldwide and can be both beneficial and detrimental to wildlife habitat. In recent years a link between feral horse (*Equus ferus caballus*) overpopulation and environmental damage has become a contentious issue due to public protests at proposals to cull large numbers (Driscoll, 2018; Scasta et al., 2018). However, the populations in these high-profile cases are not ‘native’, but are instead for the most part descended from animals which escaped from, or were released by, European settlers in the 16th century (National Research Council, 2013), and are thus categorised as ‘feral’ (defined as living in a wild state after escape from captivity or domestication). Ecologically, they are alien species, and there is mounting evidence that they are putting ecosystems at risk through trampling and vegetation community change (Davies et al., 2014; Nimmo, 2018; Rogers, 1991).

What, then, is the role of indigenous equids within their native ecosystems? In such situations, can targeted grazing by horses (taller than approx. 148 cm at the withers) and ponies (less than approx. 148 cm at the withers) achieve conservation gains? These primitive horses or ponies have lived as free-ranging populations for thousands of years with little human intervention. Such herds are classified by the EU as ‘semi-wild’ and have related derogations which exempt them from legislation relating to e.g. animal identification and treatment. Whilst overall population increase is commonly a concern with feral equids, semi-wild horse and pony populations are more generally under threat. Within the UK very few pony herds that can be designated as semi-wild remain. Two examples of semi-wild native ponies thought to have been *in situ* since the Bronze Age are Dartmoor Hill Ponies and Carneddau Mountain Ponies (Fig 1). In both cases ponies have

61 been removed to make way for domestic livestock in response to economic pressures on
62 farmers, and there is now a serious risk that they may die out. Census data from the 1960s
63 onwards estimate the Dartmoor Hill Pony population has declined from a maximum of
64 12,250 to 1,200 (*J. Hibbs, personal communication*); while there are around 300 Carneddau
65 Ponies found on Snowdonia's Carneddau Mountains today (*H. Kehoe, personal*
66 *communication*). The UK government criterion which defines semi-wild ponies (i.e. that
67 they remain outside of human control for their survival and reproduction) also means they
68 are not protected by societies or registers. As a result, they fall outside of initiatives on the
69 sustainable use of farm animal genetic resources incentivising the conservation of rare or 'at
70 risk' breeds (including breeds of horses and ponies), and the genetic variation within them.
71 Rather short-sightedly the conservation of animal genetic resources has been considered
72 primarily in relation to their potential contribution to agricultural productivity and
73 sustainability (Hall & Bradley, 1995; Rege & Gibson, 2003), with little or no regard given to
74 ecological resilience.

76 Agricultural abandonment of grasslands and heathlands across the EU due to changes in
77 farming practice has been identified as a specific threat to related habitats and species
78 (Hermoso et al., 2018; Keenleyside & Tucker, 2010), particularly in areas that are marginal
79 for agricultural production due to environmental challenges. Left unmanaged, many of
80 these vegetation communities have become dominated by plant species rejected by stock
81 (e.g. *Juncus* spp., *Molinia caerulea*, *Deschampsia cespitosa*, *Pteridium* spp, *Ulex* spp.). This
82 degradation of native plant communities limits the value of these areas for achieving
83 biodiversity conservation objectives. Consequently, government agencies and conservation
84 charities are resorting to mechanical cutting of vegetation (Talle et al., 2018; Valasiuk et al.,
85 2018), as a substitute for grazing, to ensure the long-term survival of Europe's most valuable
86 and threatened habitats and species (as listed under both the EU Birds Directive
87 2009/147/EC and the EU Habitats Directive 92/43/EEC; i.e. sites with NATURA 2000 status).
88 This is however costly, does not create the same degree of structural heterogeneity as
89 grazing, and over time leads to nutrient depletion.

91 Farming systems in areas designated by the EU as less favoured have for many years been
92 supported by specific agricultural subsidies. It is anticipated that under future support
93 schemes the wider delivery of public goods, rather than primary production, will be
94 prioritised. Crucially this could allow the grazing outcomes of stock types other than sheep
95 and cattle to be eligible for support payments. The biodiversity benefits of mixed-low-
96 intensity grazing systems have been well documented (Critchley et al., 2008; Fraser et al.,
97 2014; Liu et al., 2015; Lopez et al., 2017b) (Table 1), yet in countries including the UK semi-
98 natural grassland and heathland communities continue to be managed predominately under
99 sheep-only systems. A policy change which recognised the role of native ponies could have
100 multiple benefits; improved biodiversity, diversification of income streams, and
101 conservation of threatened genotypes of grazer. However, to date there have been little
102 robust data collected regarding the comparative impacts of equid grazing, largely because of
103 proportionately low levels of research funding compared to those with domesticated stock.
104 Unless addressed, this will continue to compromise our ability to optimise deployment of
105 conservation grazing tools, undermining our ability to meet conservation targets.

106
107 To explore these issues further we evaluated i) the potential role of native ponies as

conservation grazers, and ii) the value of supporting semi-wild ponies specifically. Firstly, we examined the comparative foraging strategies and dietary preferences of equids, and identified situations where grazing by ponies could be particularly beneficial. Secondly, we considered the behavioural ecology of ponies in the context of conservation grazing schemes and free-living populations, highlighting both the merits of using ponies as conservation grazers and behavioural factors that must be considered when using ponies in these contexts. Finally, we explored the population genetics of UK native ponies and associated evidence of adaptation.

2. Dietary preferences of equids

2.1 Physiological factors influencing foraging

A variety of factors influence the foraging choices and grazing behaviour exhibited by large herbivores. One of the key determinants of foraging behaviours is body size, since energy requirements scale to 0.75 rather than 1 (Demment & Van Soest, 1985). Consequently, larger animals are generally less selective grazers than their smaller counterparts (Sensenig & Demment, 2010), and tend to prioritise maintaining their intake rate rather than the nutritional value of what is consumed when resources become limited. These differences are evident when the dietary preferences of cattle and sheep are compared (Critchley et al., 2008; Cuchillo-Hilario et al., 2018; Fraser et al., 2009; Grant et al., 1985). Gut morphology and function also have a role to play in influencing diet composition. While ruminants are fore-gut fermenters (with fermentation taking place in the reticulorumen), equids are hind-gut fermenters (fermentation occurs in the caecum and colon). A mainly post-gastric site of fermentation (i.e. after the stomach) means they can digest and absorb available soluble carbohydrate and protein directly, without the potential inefficiencies associated with the synthesis of microbial protein (Santos et al., 2011). Evidence suggests that equids achieve higher nutrient extraction rates than bovids on all forages, whether housed or at pasture (Duncan et al., 1990; Illius & Gordon, 1992; Santos et al., 2011). Without selective retention of large particles in the rumen, digesta passes relatively quickly through the equine fermentation zone. This faster throughput is an advantage which outweighs their lower digestive efficiency, particularly on poor quality forages, allowing them to ingest large amounts of fibre-rich forage.

2.2 Comparative foraging strategies

Studies of natural grazing systems in Africa were among the first to identify differences in foraging strategies of large grazers, and categorised the equid present (the zebra) along with ruminants of a similar size as generalist (rather than specialist) feeders (Jarman & Sinclair, 1979). There is consensus from a range of ecosystems that equids prefer graminoids to browse species (Celaya et al., 2011; Ferreira et al., 2013; Gordon, 1989; Lopez et al., 2017a; Menard et al., 2002; Pratt et al., 1986; Scasta et al., 2016). However, as generalist feeders, they will switch to alternative plant species and plant parts when preferred items become depleted. Thus, like cattle, equids will incorporate woody vegetation into their diets as high-quality grassland availability declines (Putman et al., 1987; Scasta et al., 2016). However, equids have been found to be more reluctant to browse on *Calluna* spp. (a plant species that heathland grazing prescriptions are frequently designed to protect) than are cattle (Celaya et al., 2011; Ferreira et al., 2013) or sheep (Ferreira et al., 2013). Instead, ponies prefer to consume *Ulex* spp. (Putman et al., 1987) (a plant species that grazing prescriptions are often aiming to control), with dietary inclusion rates higher than for cattle

or sheep (Ferreira et al., 2013).

Studies in areas with mixed habitats found that the summer diet of ponies, like that of cattle, consisted primarily of grasses (80-90%) (Putman et al., 1987). However, *M. caerulea*, a species scarcely eaten by cattle, contributed to 20% of their diet at this time. In the UK, increased *M. caerulea* abundance has frequently been at the expense of more diverse upland heath and mire habitats (Yeo & Blackstock, 2002); a situation exacerbated by sheep-only grazing since sheep strongly avoid consuming this species. To counteract this, agri-environment schemes have offered incentives for cattle grazing, since as less selective feeders they are more likely to switch to consumption of *M. caerulea* as preferred resources decline (Critchley et al., 2008; Fraser et al., 2011). However, loss of cattle from many less favoured areas in response to economic and social challenges (poor returns, aging farmer populations, lack of labour) has meant that there in many regions there are insufficient cattle numbers to deliver recommended grazing prescriptions. Despite the potential for targeted pony grazing to also reduce *M. caerulea* dominance through increased utilisation this has not been fully explored or exploited. Indeed, the high dietary overlap between ponies and cattle shown by a range of studies (Celaya et al., 2011; Gordon, 1989; Menard et al., 2002; Pratt et al., 1986; Scasta et al., 2016) suggests that these animals could be considered broadly as alternative tools for conservation management, eligible for similar agri-environmental scheme payments based on the outcomes of their grazing.

Given that between-species comparisons of dietary preferences and grazing behaviour involving ponies are rare, it is not surprising that there is a complete lack of evidence as to similarities and differences in the diet composition of different pony breeds or types. Studies with cattle have found few differences between the diets selected by contrasting traditional and modern breed types (Fraser et al., 2009; Fraser et al., 2013), with utilisation of *M. caerulea* similar among breeds. However, results from cattle breed comparisons also highlight that adaptations to adverse environments as well behavioural responses to topographical/climatic conditions must also to be considered. Since such factors affect energy requirements and nutrient use efficiencies they can potentially influence both grazing patterns and welfare (Fraser et al., 2009; Ricci et al., 2014).

3. Behavioural ecology

3.1 Social organisation

Ponies show a relatively high level of social complexity compared with grazers such as sheep or cows, which can have certain implications for conservation grazing schemes. Free-living horses live mostly in either family bands or bachelor bands. Family bands are comprised of between two and 35 mares, along with one to two stallions and all immature offspring (Boyd & Keiper, 2005). These often show patterns of fission and fusion, where groups split up and reform frequently (CS, *personal observation*). Bands also show seasonal fluctuations in the relative level of social cohesion, most likely due to a combination of food availability and stallion herding frequency (Stanley et al., 2017). Whilst foals can be nutritionally independent of their mothers at the age of around six months old, youngsters of both sexes only naturally disperse from the group in which they are born between the ages of two and five years old (Boyd & Keiper, 2005). These social factors must be taken into account, in addition to grazing capacity, when considering stocking densities and sex ratios of ponies in conservation grazing schemes; for example if there are no neighbouring groups to which

youngsters can disperse, they will need to be removed from the group to avoid them being the targets of aggression (Stanley & Shultz, 2012). Strong social bonds exist between females and their band stallion, but social bonds between females are also important; mares are known to remain together after the death of their band stallion (Keiper, 1985). Existing social bonds and opportunities for their maintenance should therefore be a key consideration for both the selection of individuals and their management in both conservation grazing schemes and free-living populations.

Horses are polygynous, meaning they show high levels of reproductive skew (Rubenstein & Nunez, 2009). Since not all males can hold harems at any one time, yet the sex ratio is most commonly 1:1 in free-living populations (Ransom & Kaczensky, 2016), surplus males commonly form bachelor bands. These are known to number up to 16 individuals (McCort, 1984) and are thought to be mostly unstable in terms of membership (Boyd & Keiper, 2005). If the number of breeding stallions is controlled in feral populations, conservation grazing schemes elsewhere could provide a potential sink for these excess bachelor males.

3.2 Home ranges

Horse bands do not generally defend discrete territories (Boyd & Keiper, 2005); instead, they occupy overlapping but well-defined home ranges that can persist for a number of years (McCort, 1984) and vary significantly in size, both within and between populations (Boyd & Keiper, 2005). Such site fidelity could make ponies more suited to targeting specific locations within a larger habitat for conservation grazing. Whilst hefting (selective breeding to instil a specific home range) has traditionally been used to achieve this in sheep, reduced flock sizes in recent years has reduced the effectiveness of this strategy. A herd (a group of horse bands living in same geographical area) can show significant social structure, where bands follow similar movements and have an inter-band dominance hierarchy (Miller, 1979). Home range size can vary seasonally in feral horse populations, although there seems to be no consistent trend as local conditions seem to have a significant influence (Boyd & Keiper, 2005). Habitat utilisation does, however, appear to vary seasonally in most populations; in mountainous regions, vertical migrations may occur to benefit from changing vegetation quality and abundance (Berger, 1986; Linklater et al., 2000), whilst water availability (Berger, 1977) and tabanid fly abundance (Keiper & Berger, 1982) can also influence home range use. It is important to note that whilst the factors affecting home range use have been studied across a variety of feral populations, our understanding of the influence of herd composition on grazing behaviour is lacking.

3.3 Influence of contraception

Contraceptive vaccines are being increasingly used to control female fertility and therefore limit population sizes in feral horses. Since natural predators are frequently absent from habitats in which feral horses thrive, management interventions are often required to control population sizes (Ransom et al., 2016; Saltz, 2002). However, the use of contraception can influence both behaviour and range use (Nunez et al., 2009; Ransom et al., 2010), and this could have implications both for the welfare of free-living populations and range use in conservation grazing schemes. Gelding excess colts is another approach to population control, but again the potential impacts on subsequent resource utilisation are poorly understood. Of course, such approaches to meeting sustained external pressures to control numbers also pose additional risks to already rare populations such as the

Carneddau and Dartmoor Hill Ponies, including a reduction in genetic diversity and the introduction of artificial selection to these populations. Interventions such as removal of stallions, gelding of stallions, and mare contraception all result in management decisions being imposed upon the herds, threatening their semi-wild status and potentially impacting upon the selection of specific genes and traits correlated with adaptation to the harsh environments they have evolved in.

Males used in conservation grazing schemes are frequently castrated (gelded) to avoid breeding. Gelding is also commonly carried out to facilitate handling and management (McDonnell, 2005). However, groups of geldings likely differ in home range use to the more well-studied family bands, or even compared with non-castrated bachelor bands. This could have implications for the management of gelding groups, specifically in terms of predicting their seasonal movements.

4. Population genetics

4.1 How genetically distinct are breeds?

Science has only recently begun to address the question of how distinct free-living, unmanaged populations of horses and ponies are from their registered counterparts. Factors such as adaptation, artificial selection (in the registered breeds) and inbreeding must be considered. Genetically speaking, there is no universally accepted definition of a 'breed', which is an artificial human concept based on a closed (or restricted) breeding pool of individuals which share a common phenotype (typically purely morphological i.e. height, coat colour). In cattle, for example, the primary distinction between Red and Black Angus is due to a mutation in a single coat colour gene (Matukumalli et al., 2009). In the case of semi-wild pony populations then, the question of how distinct they are from their registered relatives is perhaps not as important as the reason for any differences (i.e. are they signatures of adaptation?).

4.2 How do native pony breeds differ?

In the first studies of their kind, several studies of population diversity and relationships in both registered and semi-wild UK pony populations were undertaken (Winton et al. 2013; McMahon et al. 2015; Hegarty et al. 2017). Genotype data of semi-wild populations were generated using 15 simple sequence repeat markers (SSR) and 162 single nucleotide polymorphism (SNP) markers on samples from 16 herds of Welsh Mountain Hill Pony (McMahon et al., 2015) and 19 herds of Dartmoor Hill Ponies (Hegarty et al., 2017). These data were compared to existing genotypes (Winton et al., 2013) of Section A Welsh Ponies and Section D Welsh Cobs (breed society registered), the Carneddau Pony (semi-wild), Connemara Ponies (breed society registered), Irish Draught horse (breed society registered) and Dartmoor Ponies (breed society registered) (Fig 2).

These analyses demonstrated that the Carneddau Pony is genetically distinct from the Section A Welsh Pony (Winton et al., 2013), despite the Carneddau Ponies generally being thought of as being either an offshoot or an ancestor of the Welsh Section A. Population structure analysis showed that the two types are related, but distinct, and mitochondrial DNA analysis identified several rare variants present at high frequency in the Carneddau Ponies but not any of the other types. High levels of deviation from Hardy-Weinberg equilibrium were detected which are indicative of selective pressure, and it was argued that

conservation or further study of the Carneddau Ponies was important to avoid losing useful genetic adaptations.

Similar results were observed in studies of the Welsh Mountain Hill Pony (McMahon et al., 2015) and Dartmoor Hill Ponies (Hegarty et al., 2017). In both cases, the populations can be distinguished from their registered counterparts. Thus, whilst both semi-wild populations are related to their registered cousins, there are key genetic signatures which are common across animals from the same group and can be used to distinguish them. The largest genetic difference observed was that between the Carneddau and the Section A Welsh, with a Wright's F_{st} (Wright, 1965) measure of genetic distance 0.157 when all datasets were compared. This value is comparable to the difference seen between established horse breeds in studies in France (Leroy et al., 2009) and Poland (Stachurska et al., 2014). Whilst we continue to argue that genetic degrees of difference should not be a hard limit on what humans consider a 'breed', the Carneddau Pony is clearly a distinct population far older than the Welsh Section A. Interestingly, other close relationships are apparent between the 'upland' types, suggesting the possibility of a shared 'upland' signature linked to adaptation, though other factors such as common ancestry may be in play. These studies were the first in the world to explore the comparative population genetics of different breeds and types of equine, and clearly show there is much to be learned about adaptation and related environmental impacts.

5. Conclusions

At a time when the ecological role of feral horses is under considerable scrutiny this paper presents evidence that grazing by native horse and ponies could play an important role in restoring and maintaining habitats of conservation importance. Semi-wild populations could be particularly suited to conservation grazing schemes due to both physiological and behavioural adaptations to a free-roaming lifestyle. They might also have a particular genetic value; their genomes have been shaped by natural and not artificial selection, meaning these ponies may have ecological adaptations no longer present in domesticated ponies. These populations are also highly valued on a cultural basis and are an important part of their regions' heritage. Conservation grazing schemes could provide an important sink for excess individuals from these populations, allowing satellite populations to be maintained as an 'insurance policy' against catastrophic events to founder populations. Such factors should be taken into account as new policies and prescriptions are being developed to meet revised targets for biodiversity gain.

Acknowledgements

We would like to thank the Friends of the Dartmoor Hill Ponies and PONT (Pori, Natur a Threftadaeth) for background information on the Dartmoor Hill Ponies and the Carneddau Ponies. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Berger, J. 1977. Organizational systems and dominance in feral horses in the Grand Canyon. *Behavioral Ecology and Sociobiology*, **2**(2), 131-146.
- Berger, J. 1986. *Wild horses of the Great Basin*. University of Chicago Press, Chicago.
- Boyd, L., Keiper, R.R. 2005. Behavioural ecology of feral horses. in: *The Domestic Horse*, (Eds.) D. Mills, S. McDonnell, Cambridge University Press. Cambridge, pp. 55-82.
- Celaya, R., Ferreira, L.M.M., Garcia, U., Garcia, R.R., Osoro, K. 2011. Diet selection and performance of cattle and horses grazing in heathlands. *Animal*, **5**(9), 1467-1473.
- Critchley, C.N.R., Adamson, H.F., McLean, B.M.L., Davies, O.D. 2008. Vegetation dynamics and livestock performance in system-scale studies of sheep and cattle grazing on degraded upland wet heath *Agriculture, Ecosystems and Environment*, **128**, 59-67.
- Cuchillo-Hilario, M., Wrage-Monnig, N., Isselstein, J. 2018. Forage selectivity by cattle and sheep co-grazing swards differing in plant species diversity. *Grass and Forage Science*, **73**(2), 320-329.
- Davies, K.W., Collins, G., Boyd, C.S. 2014. Effects of feral free-roaming horses on semi-arid rangeland ecosystems: an example from the sagebrush steppe. *Ecosphere*, **5**(10).
- Demment, M.W., Van Soest, P.J. 1985. A nutritional explanation for body-size patterns of ruminant and non-ruminant herbivores. *American Naturalist*, **125**, 641-675.
- Dennis, P., Skartveit, J., McCracken, D.I., Pakeman, R.J., Beaton, K., Kunaver, A., Evans, D.M. 2008. The effects of livestock grazing on foliar arthropods associated with bird diet in upland grasslands of Scotland. *Journal of Applied Ecology*, **45**, 279-287.
- Driscoll, D. 2018. Without culling, Victoria's feral horse plan looks set to fail. *The Conversation*.
- Duncan, P., Foose, T., Gordon, I.J., Gakahu, C., Lloyd, M. 1990. Comparative nutrient extraction by grazing bovids and equids: a test of the nutritional models of equid/bovid competition. *Oecologia, Berlin*, **84**, 411-418.
- Evans, D.M., Redpath, S.M., Elston, D.A., Evans, S.A., Mitchell, R.J., Dennis, P. 2006a. To graze or not to graze? Sheep, voles, forestry and nature conservation in the British uplands. *Journal of Applied Ecology*, **43**(3), 499-505.
- Evans, D.M., Redpath, S.M., Evans, S.A., Elston, D.A., Gardner, C.J., Dennis, P., Pakeman, R.J. 2006b. Low intensity, mixed livestock grazing improves the breeding abundance of a common insectivorous passerine. *Biology Letters*, **2**(4), 636-638.
- Evans, D.M., Villar, N., Littlewood, N.A., Pakeman, R.J., Evans, S.A., Dennis, P., Skartveit, J., Redpath, S.M. 2015. The cascading impacts of livestock grazing in upland ecosystems: a 10-year experiment. *Ecosphere*, **6**(3).
- Ferreira, L.M.M., Celaya, R., Benavides, R., Jauregui, B.M., Garcia, U., Santos, A.S., Garcia, R.R., Rodrigues, M.A.M., Osoro, K. 2013. Foraging behaviour of domestic herbivore species grazing on heathlands associated with improved pasture areas. *Livestock Science*, **155**(2-3), 373-383.
- Fraser, M.D., Moorby, J.M., Vale, J.E., Evans, D.M. 2014. Mixed grazing systems benefit both upland biodiversity and livestock production. *PLoS ONE*, **9**, e89054.
- Fraser, M.D., Theobald, V.J., Dhanoa, M.S., Davies, O.D. 2011. Impact on sward composition and stock performance of grazing *Molinia*-dominant grassland. *Agriculture, Ecosystems and Environment*, **144**, 102-106.

383 Fraser, M.D., Theobald, V.J., Griffiths, J.B., Morris, S.M., Moorby, J.M. 2009. Comparative diet
384 selection by cattle and sheep grazing two contrasting heathland communities. *Agriculture,*
385 *Ecosystems and Environment*, **129**(1-3), 182-192.

386 Fraser, M.D., Vale, J.E., Dhanoa, M.S. 2013. Alternative upland grazing systems: impacts on livestock
387 performance and sward characteristics. *Agriculture, Ecosystems and Environment*, **175**, 8-20.

388 Garcia, R.R., Garcia, U., Osoro, K., Celaya, R. 2011. Ground-dwelling arthropod assemblages of
389 partially improved heathlands according to the species of grazer and grazing regime.
390 *European Journal of Entomology*, **108**(1), 107-115.

391 Gordon, I.J. 1989. Vegetation community selection by ungulates on the Isle of Rhum. III.
392 Determinants of vegetation community selection. *Journal of Applied Ecology*, **26**, 65-79.

393 Grant, S.A., Suckling, D.E., Smith, H.K., Torvell, L., Forbes, T.D.A., Hodgson, J. 1985. Comparative
394 studies of diet selection by sheep and cattle: the hill grasslands. *Journal of Ecology*, **73**, 987-
395 1004.

396 Hall, S.J.G., Bradley, D.G. 1995. Conserving livestock breed biodiversity. *Trends in Ecology &*
397 *Evolution*, **10**(7), 267-270.

398 Hegarty, M., McElhinney, N., Ham, E., Potter, C., Winton, C.L., McMahon, R. 2017. An evaluation of
399 the genetic relationships between the Hill Dartmoor and the registered Dartmoor Pony
400 Breed <http://hdl.handle.net/2160/46846>.

401 Hermoso, V., Moran-Ordóñez, A., Brotons, L. 2018. Assessing the role of Natura 2000 at maintaining
402 dynamic landscapes in Europe over the last two decades: implications for conservation.
403 *Landscape Ecology*, **33**(8), 1447-1460.

404 Illius, A.W., Gordon, I.J. 1992. Modelling the nutritional ecology of ungulate herbivores: evolution of
405 body size and competitive interactions. *Oecologia, Berlin*, **89**, 428-434.

406 Jarman, P.J., Sinclair, A.R.E. 1979. Feeding strategy and the pattern of resource partitioning in
407 ungulates. in: *Serengeti, Dynamics of an Ecosystem*, (Eds.) A.R.E. Sinclair, M. Norton-Griffiths,
408 University of Chicago Press. Chicago, pp. 130-163.

409 Keenleyside, C., Tucker, G.M. 2010. Farmland Abandonment in the EU: an Assessment of Trends and
410 Prospects.

411 Keiper, R. 1985. *The Assateague Ponies*. Tidewater Press, Cambridge, MD.

412 Keiper, R.R., Berger, J. 1982. Refuge-seeking and pest avoidance by feral horses in desert and island
413 environments. *Applied Animal Ethology*, **9**(2), 111-120.

414 Leroy, G., Callède, L., Verrier, E., Mériaux, J., Ricard, A., Danchin-Burge, C., Rognon, X. 2009. Genetic
415 diversity of a large set of horse breeds raised in France assessed by microsatellite
416 polymorphism. *Genetics Selection Evolution*, **41**, 5.

417 Linklater, W.L., Cameron, E.Z., Stafford, K.J., Veltman, C.J. 2000. Social and spatial structure and
418 range use by Kaimanawa wild horses (*Equus caballus*: Equidae). *New Zealand Journal of*
419 *Ecology*, **24**, 139-152.

420 Liu, J., Feng, C., Wang, D.L., Wang, L., Wilsey, B.J., Zhong, Z.W. 2015. Impacts of grazing by different
421 large herbivores in grassland depend on plant species diversity. *Journal of Applied Ecology*,
422 **52**(4), 1053-1062.

423 Lopez, C.L., Ferreira, L.M.M., Garcia, U., Moreno-Gonzalo, J., Rodrigues, M.A.M., Osoro, K., Ferre, I.,
424 Celaya, R. 2017a. Diet selection and performance of horses grazing on different heathland
425 types. *Animal*, **11**(10), 1708-1717.

426 Lopez, C.L., Garcia, R.R., Ferreira, L.M.M., Garcia, U., Osoro, K., Celaya, R. 2017b. Impacts of horse
 427 grazing on botanical composition and diversity in different types of heathland. *Rangeland*
 428 *Journal*, **39**(4), 375-385.

429 Matukumalli, L.K., Lawley, C.T., Schnabel, R.D., Taylor, J.F., Allan, M.F., Heaton, M.P., O'Connell, J.,
 430 Moore, S.S., Smith, T.P.L., Sonstegard, T.S., Van Tassell, C.P. 2009. Development and
 431 characterization of a high density SNP genotyping assay for cattle. *PLoS One*, **4**, e5350.

432 McCort, W.D. 1984. Behavior of feral horses and ponies. *Journal of Animal Science*, **58**(2), 493-499.

433 McDonnell, S. 2005. Sexual behaviour. in: *The domestic horse: the origins, development and*
 434 *management of its behaviour*, (Eds.) D. Mills, S. McDonnell, Cambridge University Press.
 435 Cambridge, pp. 110-125.

436 McMahan, R., Debbonaire, A., McEwan, N., Nash, D., Davies-Morel, M. 2015. A preliminary
 437 examination of the genetic variation within and between the Improvement Society herds of
 438 Welsh Mountain ponies <http://hdl.handle.net/2160/46844>.

439 Menard, C., Duncan, P., Fleurance, G., Georges, J.Y., Lila, M. 2002. Comparative foraging and
 440 nutrition of horses and cattle in European wetlands. *Journal of Applied Ecology*, **39**(1), 120-
 441 133.

442 Miller, R. 1979. Interband dominance in feral horses. *Ethology*, **51**(1), 41-47.

443 National Research Council. 2013. Using science to improve the BLM wild horse and burrow program:
 444 a way forward.

445 Nimmo, D.G. 2018. Feral horses in Australia: Left unchecked they can have major ecological impacts.
 446 *Austral Ecology*, **43**(6), E14-E15.

447 Nunez, C.M., Adelman, J.S., Mason, C., Rubenstein, D.I. 2009. Immunocontraception decreases group
 448 fidelity in a feral horse population during the non-breeding season. *Applied Animal*
 449 *Behaviour Science*, **117**(1), 74-83.

450 Pratt, R.M., Putman, R.J., Ekins, J.R., Edwards, P.J. 1986. Use of habitat by free-ranging cattle and
 451 ponies in the New Forest, southern England. *Journal of Applied Ecology*, **23**, 539-557.

452 Putman, R.J., Pratt, R.M., Ekins, J.R., Edwards, P.J. 1987. Food and feeding-behaviour of cattle and
 453 ponies in the New Forest, Hampshire. *Journal of Applied Ecology*, **24**(2), 369-380.

454 Ransom, J.I., Cade, B.S., Hobbs, N.T. 2010. Influences of immunocontraception on time budgets,
 455 social behavior, and body condition in feral horses. *Applied Animal Behaviour Science*,
 456 **124**(1), 51-60.

457 Ransom, J.I., Kaczensky, P. 2016. *Wild Equids: Ecology, Management, and Conservation*. JHU Press.

458 Ransom, J.I., Lagos, L., Hrabar, H., Nowzari, H., Usukhjargal, D., Spasskaya, N. 2016. *Wild and Feral*
 459 *Equid Population Dynamics*. In: *Wild Equids: Ecology, Management and Conservation*. John
 460 Hopkins University Press, Maryland.

461 Rege, J.E.O., Gibson, J.P. 2003. Animal genetic resources and economic development: issues in
 462 relation to economic valuation. *Ecological Economics*, **45**(3), 319-330.

463 Ricci, P., Umstatter, C., Holland, J.P., Waterhouse, A. 2014. Does diverse grazing behavior of suckler
 464 cows have an impact on predicted methane emissions? *Journal of Animal Science*, **92**(3),
 465 1239-1249.

466 Rogers, G.M. 1991. Kaimanawa feral horses and their environmental impacts. *New Zealand Journal*
 467 *of Ecology*, **15**, 49-64.

- 468 Rubenstein, D.I., Nunez, C.M. 2009. Sociality and reproductive skew in horses and zebras. in:
469 *Reproductive Skew in Vertebrates*, (Eds.) R. Hager, C.B. Jones, Cambridge University Press.
470 Cambridge, pp. 196-226.
- 471 Saltz, D. 2002. The dynamics of equid populations. In: Zebra, asses, and horses: an action plan for the
472 conservation of wild equids. World Conservation Union, Gland, Switzerland.
- 473 Santos, A.S., Rodrigues, M.A.M., Bessa, R.J.B., Ferreira, L.M., Martin-Rosset, W. 2011. Understanding
474 the equine cecum-colon ecosystem: current knowledge and future perspectives. *Animal*,
475 **5**(1), 48-56.
- 476 Scasta, J.D., Beck, J.L., Angwin, C.J. 2016. Meta-Analysis of Diet Composition and Potential Conflict of
477 Wild Horses with Livestock and Wild Ungulates on Western Rangelands of North America.
478 *Rangeland Ecology & Management*, **69**(4), 310-318.
- 479 Scasta, J.D., Hennig, J.D., Beck, J.L. 2018. Framing contemporary US wild horse and burro
480 management processes in a dynamic ecological, sociological, and political environment.
481 *Human-Wildlife Interactions*, **12**(1), 31-45.
- 482 Sensenig, R.L., Demment, M.W. 2010. Allometric scaling predicts preferences for burned patches in a
483 guild of East African grazers. *Ecology*, **91**, 2898-2907.
- 484 Stachurska, A., Nogaj, A., Brodacki, A., Nogaj, J., Batkowska, J. 2014. Genetic distances between
485 horse breeds in Poland estimated according to blood protein polymorphism. *Czech Journal*
486 *of Animal Science*, **59**, 257-267.
- 487 Stanley, C.R., Mettke-Hofmann, C., Hager, R., Shultz, S. 2017. Social stability in semiferall ponies:
488 networks show interannual stability alongside seasonal flexibility. *Animal Behaviour*.
- 489 Stanley, C.R., Shultz, S. 2012. Mummy's boys: sex differential maternal-offspring bonds in semi-feral
490 horses. *Behaviour*, **149**, 251-274.
- 491 Talle, M., Deak, B., Poschlod, P., Valko, O., Westerberg, L., Milberg, P. 2018. Similar effects of
492 different mowing frequencies on the conservation value of semi-natural grasslands in
493 Europe. *Biodiversity and Conservation*, **27**(10), 2451-2475.
- 494 Valasiuk, S., Giergiczny, M., Zylicz, T., Klimkowska, A., Angelstam, P. 2018. Conservation of
495 disappearing cultural landscape's biodiversity: are people in Belarus willing to pay for wet
496 grassland restoration? *Wetlands Ecology and Management*, **26**(5), 943-960.
- 497 Winton, C.L., Hegarty, M.J., McMahon, R., Slavov, G.T., McEwan, N.R., Davies Morel, M.C.G.,
498 Morgan, C.M., Powell, W., Nash, D.M. 2013. Genetic diversity and phylogenetic analysis of
499 native mountain ponies of Britain and Ireland reveal a novel rare population. *Ecology and*
500 *Evolution*, **3**, 934-947.
- 501 Wright, S. 1965. The interpretation of population structure by F-statistics with special regard to
502 systems of mating. *Evolution*, **19**, 395-420.
- 503 Yeo, M.J.M., Blackstock, T.H. 2002. A vegetation analysis of the pastoral landscapes of upland Wales,
504 UK. *Journal of Vegetation Science*, **13**(6), 803-816.

505

Group		Genus/species	Livestock species	Response to mixed grazing	Country	
Plants	Tracheophyta	Combined	Cattle + sheep	+	UK China	(Critchley et al., 2008; Evans et al., 2015) (Liu et al., 2015)
Mammals	Rodentia	Vole (<i>Microtus agrestis</i>)	Cattle + sheep	+	UK	(Evans et al., 2006a; Evans et al., 2015)
	Carnivora	Red fox (<i>Vulpes vulpes</i>)	Cattle + sheep	+	UK	(Evans et al., 2015)
Birds	Passeriformes	Meadow pipit (<i>Anthus pratensis</i>)	Cattle + sheep	+	UK	(Evans et al., 2006b; Evans et al., 2015)
Arthropods	Araneae	Combined	Cattle + sheep Cattle + goats	+	Spain	(Garcia et al., 2011)
	Opiliones	Combined	Cattle + sheep Cattle + goats	+/-	Spain	(Garcia et al., 2011)
	Coleoptera	Combined	Cattle + sheep	+	UK	(Dennis et al., 2008)

Table 1: Examples of responses of different taxa to mixed grazing of semi-natural vegetation communities; where + = positive effects from mixed grazing, +/- = no response to mixed grazing; compared to single-species grazing treatments. Study duration ranged from 3 to 10 years.

Figure 1: The distinct geographic locations where the semi-wild Carneddau Ponies and Dartmoor Hill Ponies are found. Breeding studs and animals registered with breed societies such as the Welsh Pony and Cob Society and the Dartmoor Pony Society are found throughout the British Isles (map reproduced from Ordnance Survey map data by permission of the Ordnance Survey © Crown copyright 2001).

Figure 2: Results of population assignment clustering for all pony types based on 162 single nucleotide polymorphism markers, highlighting the close relationships of the Hill Welsh, Carneddau and Dartmoor Hill ponies. Subfigures show the results of population assignment for these three types relative to Section A Welsh (for Carneddau and Hill Welsh) and Registered Dartmoor Pony (for Dartmoor Hill Pony) in order to show that deeper levels of resolution enables separation of these types.