Aberystwyth University

What is the effect of 19 years of restoration managements on soil and vegetation on formerly improved upland grassland?

Pavlu, L.; Pavlu, Vilem; Fraser, Mariecia

Published in:

Science of the Total Environment

DOI:

[10.1016/j.scitotenv.2020.142469](https://doi.org/10.1016/j.scitotenv.2020.142469)

Publication date:

2021

Citation for published version (APA):

Pavlu, L., Pavlu, V., & Fraser, M. (2021). What is the effect of 19 years of restoration managements on soil and vegetation on formerly improved upland grassland? *Science of the Total Environment*, 755(2), 142469. [142469]. <https://doi.org/10.1016/j.scitotenv.2020.142469>

Document License

CC BY-NC-ND

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400

email: is@aber.ac.uk

To what extent did 19 years of restoration managements on formerly improved upland grassland alter soil and vegetation?

Lenka Pavlu^a, Pavlu V. Vilém^{a,b,*} and Mariecia D. Fraser^c

^a Crop Research Institute, Department of Weeds and Vegetation of Agroecosystems, Grassland Research Station Liberec, CZ 460 01, Czechia

^b Czech University of Life Sciences Prague, Faculty of Environmental Sciences, Department of Ecology, CZ 165 21, Czechia

^c Pwllpeiran Upland Research Centre, Aberystwyth University, Cwmystwyth, Aberystwyth, SY23 4AD, United Kingdom

* Corresponding author at: Department of Weeds and Vegetation of Agroecosystems, Grassland Research Station Liberec, Crop Research Institute, Rolnická 6, CZ 460 01 Liberec, Czechia.
E-mail addresses: pavlu@vurv.cz, grass@volny.cz, pavluv@fzp.czu.cz (V.V. Pavlu).

ABSTRACT

Finding the best management strategies to restore grassland diversity and achieve a compromise between agricultural use and biodiversity protection is a global challenge. This paper reports novel data relating to the impacts of 19 years of restoration managements predicted to increase botanical diversity within reseeded upland temperate grassland common in less favoured areas in Europe. The treatments imposed were: continuous sheep grazing, with and without lime application; hay cutting only, with and without lime application; hay cutting followed by aftermath grazing, with and without lime application; and a control

treatment continuing the previous site management (liming, NPK application and continuous sheep grazing).

Defoliation type, irrespective of liming, was the key driver influencing plant species diversity (hay cutting followed by aftermath grazing > hay cutting > grazing). Grazing only managements supported grasses at the expense of forbs, and thus related plant species diversity significantly declined. Limed treatments had higher concentrations of Ca and Mg in the soil compared to those receiving no lime. However, no effects on species richness or plant species composition were found. Potassium was the only element whose plant-available concentration in the soil tended to decrease in response to cutting treatments with herbage removal.

Postponing the first defoliation to the middle of the growing season enables forbs to reach seed production, and this was the most effective restoration management option for upland grassland (as hay cutting only, and as hay cut followed by aftermath grazing). Although continuous low-density sheep grazing is often adopted as a means of improving floristic biodiversity, deleterious effects of this on plant diversity mean that it cannot be recommended as a means of long-term maintenance or restoration management of European temperate grasslands.

Keywords: Cutting, Grazing, Liming, Nutrients, Restoration, Species richness, Sward

1. Introduction

Grasslands are among the most important biotopes in Central Europe, providing habitats where almost two-thirds of endangered plant species occur (Jongepierová et al., 2018), and some type of defoliation is required for their management (Bakker, 1989). Both intensive farming practices (e.g. high application rates of N, P and K plus frequent offtake of

biomass) and cessation of management can cause a significant decline in related biodiversity. Finding optimal ways to restore grassland floristic diversity is the subject of much debate amongst land managers, ecologists, and conservationists. The overall aim of fertilizer and/or lime application in intensive grassland management is to increase biomass yield, but increasing nutrient availability simultaneously reduces plant species richness (Bakker et al., 2002; Hejerman et al., 2010; Storkey et al., 2015; Goulding et al., 2016; Humbert et al., 2016; Titěra et al., 2020). Cessation of fertilization will not lead to immediate restoration of species-rich grassland due to a combination of a lack of diaspore sources together with high residual levels of available nutrients and changes in soil microbial activity (Pegtel et al., 1996; Smith et al., 2008; Pavlů et al., 2011).

There are several strategies that can be applied to reverse nutrient accumulation caused by intensive grassland management. Long-term biomass removal by hay-making or grazing without fertilization are both seen as potential ways to remove soil nutrients (Hansson and Fogelfors, 2000; Van Diggelen and Marrs, 2003) and increase plant species diversity. However, reducing concentrations of excess nutrients within the soil and increasing species richness through taking two or more annual cuts can be a lengthy and difficult process (Pavlů et al., 2011), clearly depending on the period over which any particular management regime has been adopted, on soil nutrient status, and vegetation structure. Consequently, conservation management after nutrient removal should include the traditional practices that have historically contributed to the formation of the biological diversity of semi-natural grasslands (Bonari et al., 2017).

According to some authors it is possible to reduce K concentrations in the soil relatively quickly (Parr and Way, 1988; Schaffers et al., 1998; Pavlů et al., 2013). However, P reduction is reported as being much slower (Perring et al., 2009). High concentrations of plant-available P in the soil are particularly associated with low species richness and

dominance of highly productive species (Janssens et al., 1998; Hejerman et al., 2010; Klaus et al., 2011) while, in contrast, high soil K concentrations are compatible with high values of plant diversity (Janssens et al., 1998; Crawley et al., 2005). Although Janssens et al. (1998) conclude N is the main element limiting plant diversity, its availability is considered to be controlled by P as this is an important nutrient for the symbiotic fixation of atmospheric N in legumes and for the mineralization of organic matter in soils. The interdependency of P and N was highlighted by the studies of Crawley et al. (2005) and Klaus et al. (2013). Lime application to increase soil pH is another wide-spread agricultural practice undertaken to improve herbage production through enhancing nutrient availability in the soil (Holland et al. 2018) that has a long-lasting impact on grassland (Spiegelberger et al., 2006). Liming can have both negative and positive effects on plant diversity depending on the number of species with different pH optima in a species pool. However, in areas characterised by long-term acidification, liming is used as an important tool in the restoration of species-rich grassland habitats (de Graaf et al., 1998).

The restoration of species-rich grassland on previously agriculturally-improved pastures has several further specific abiotic and biotic constraints, including a degradation period. In general, re-establishment of species from the seed bank is considered to be poor if this degradation phase takes more than a few decades. However, if the lost species are still present in the locally surrounding vegetation there is a chance to restore degraded communities (van Diggelen and Marrs, 2003 and citations therein). In situations where inappropriate abiotic conditions and lack of propagules are not barriers to re-establishment of desirable vegetation, management regime can be a key driver influencing floristic diversity (Pavlů et al., 2011). As indicated previously, two basic defoliation options are hay-making and grazing, which can also be used in combination (van Diggelen and Marrs, 2003). However, the effects of grazing and hay-making on species richness and plant species

composition differ in several ways (Hansson and Fogelfors, 2000; Krahulec et. al., 2001; Mogg et al., 2002; Mládková et al. 2015). During hay-making the above-ground biomass is non-selectively cut and removed at the same time, while factors affecting vegetation under grazing management include stocking rate, selective grazing, trampling and nutrient enrichment (Wallis DeVries, 1998; Stewart and Pullin, 2008; Ludvíková et al., 2014). Sheep are more selective of forbs and legumes than cattle, with preferential grazing encouraged by low grazing pressure (Dummont et al., 2011). A combination of grazing with cutting is generally recommended to minimize conservation risks (Krahulec et al., 2001).

In this paper we describe grassland community status after 19 years of continual exposure to various alternative restoration regimes. We compare the species composition and richness, and soil chemical characteristics when managed according to one of seven regimes that represent the common and best practices in less favoured areas dominated by temperate European upland grassland. Within this context, we aimed to answer the following questions: (i) what are the effects of long-term restoration managements on species richness, plant species composition and soil chemical properties?, (ii) what is the effect of previous liming on species richness, plant species composition and soil chemical properties?, and (iii) can any regime can be recommended for restoring plant diversity of temperate upland grasslands?

2. Materials and Methods

2.1. Experimental design

The experimental plots used (the Brignant plots) were set up in 1994 to test the effectiveness of various restoration management regimes in achieving reversion of upland improved permanent pasture to semi-natural vegetation (Hayes and Tallowin, 2007). They

were established at the Pwllpeiran Upland Research Centre on permanent pasture that had been ploughed and reseeded in 1973, and which had received regular inputs of fertilizer and lime. Sown grass species still dominated the sward at the time the plots were established, particularly *Lolium perenne*, at 58% cover. The plots are located at 310 m a.s.l. (O.S. Ref: SN752757) on free-draining typical brown podzolic soils. The area receives a mean annual rainfall of approximately 1850 mm and has average minimum and maximum air temperatures of 5.2 °C and 11.9 °C respectively. The plots are arranged in a randomized block design with three blocks and a total of seven grassland management regimes imposed (see Supplementary materials Figs S1 and S2). The treatments, which have been running continuously since 1994, are: sheep grazing, with (GL+) and without (GL-) lime application; hay cutting only, with (HL+) and without (HL-) lime application; and hay cutting followed by aftermath sheep grazing, with (HGL+) and without (HGL-) lime application. Control (CO) plots continuing the previous site management (i.e. limed, fertilised and continually grazed by sheep) are also included within each block. These receive an annual application of 60 kg ha⁻¹ N and 30 kg P ha⁻¹, with K also applied as required to maintain an index of 2+ (ADAS, 1983). Soil samples are taken in spring and tested for K concentration. Additional K is then applied by hand if the concentration has fallen below that equivalent to an index of 2 (ADAS, 1983). All of the lime treatment plots received a single application of lime in 1998 with the intention of maintaining a soil pH of 6.0. Treatments are imposed on three replicate plots of 0.08 ha (hay cut only) or 0.15 ha (grazed) in size. The schematic block design of the experiments and an aerial photo are provided in Appendix A, Figs S1 and S2.

From 1994 to 2012 the plots were stocked with ewes (usually Welsh Hill Speckled Faced yearlings) with numbers adjusted to maintain a sward surface height of approximately 4 to 6 cm. Turnout occurred late April/early May when there was sufficient biomass to sustain stock. There was no spring grazing of the HGL+ and HGL- treatments, to allow colonising

forb species an opportunity to establish. The HL+, HL-, HGL+, HGL- plots had a single hay harvest taken annually after the 15th of July, as and when weather conditions allowed. Plots were subsequently restocked on the HGL+ and HGL- treatments after a short period of regrowth. All stock were removed at the end of September/early October, depending on seasonal climatic conditions and related biomass growth.

2.2. Measurements

Data and samples were collected in July 2012, 18 years after the treatments were imposed. Visual percentage cover of vascular plant species was estimated in ten randomly located (0.4 m × 0.4 m) quadrats per plot in July 2012. The mean of ten quadrats of botanical composition was used for statistical evaluation. The nomenclature of the plant species follows Kubát et al. (2002). The plant species diversity was evaluated by plant species richness, Shannon (H) species diversity index and Shannon (J) species evenness index (Begon et al., 2005).

Fifteen individual soil cores were taken to a depth of 7.5 cm from randomly located areas from within each plot. The soil cores per plot were bulked then air-dried, biomass residues and roots were removed and the samples were then ground in a mortar to pass a 2 mm sieve. All chemical analyses were performed in an accredited laboratory of the Crop Research Institute in Chomutov. Plant-available Ca, K, Mg and P were extracted by the Mehlich III method (Mehlich 1984) and concentrations were determined by inductively coupled plasma optical emission spectrometry (GBC Scientific Equipment Pty Ltd, Melbourne, Australia). Determination of pH (CaCl₂) was done using a pH meter (Sentron Welling, Leek, The Netherlands). Total N soil concentrations were determined using the Kjeldahl method and organic C concentrations by the conventional oxidation procedure

incorporating chromo-sulphuric acid and colorimetry (AOAC, 1984).

2.3. Data analysis

A linear mixed-effects model (LMM) with fixed effects of treatment and random effect of the block was used to analyse the effect of treatment on selected plant species, cover of graminoids and forbs, species richness, Shannon (H) species diversity index, Shannon (J) species evenness index, and soil chemical properties. If necessary, data was log-transformed to meet LLM assumption requirements. Benjamini-Hochberg's procedure was applied to control for false-discovery rate (FDR) (Verhoeven, Simonsen, & McIntyre 2005). To identify significant differences between individual treatments a post-hoc comparison using Tukey's range test was applied. All LMM analyses were performed in Statistica 13.1 (Dell Inc., Texas, 2016).

Redundancy analysis (RDA) in the CANOCO 5 program (ter Braak and Šmilauer, 2012) was used to evaluate multivariate vegetation and soil data. All cover, soil and herbage chemical properties data in RDA were logarithmically transformed [$y = \log(y + 1)$]. For multivariate data analyses, two approaches were taken: i) impact of liming (comparing Limed+ versus Limed- treatments, regardless of cutting management); ii) impact of defoliation management (comparing Grazing, Hay, and Hay + Grazing treatments, regardless of liming). For all analyses 999 permutations were performed, with blocks used as covariables to restrict permutations into blocks. To visualize the results of the RDA analyses standard biplot ordination diagrams were generated.

3. Results

3.1. Species richness

Significant differences in the mean numbers of all vascular plant species, the mean numbers of vascular plant species with cover $\geq 1\%$, and Shannon (H) species diversity indices were recorded between treatments (Table 1). Shannon (J) species evenness indices did not show significant differences between treatments. The mean numbers of all vascular plant species, the mean numbers of vascular plant species with cover $\geq 1\%$ and the Shannon species diversity indices were almost always significantly higher under treatments that included a cutting regime (HL+, HL-, HGL+, and HGL-) than those which included grazing management only (GL+, GL-, CO) (Fig. 1). Fertilizer applications and liming had no significant influence on species richness or diversity (Table 1).

3.2. Plant species composition

Based on RDA the effect of liming on plant species composition was negligible, and most of the variability in plant species composition was explained by defoliation regime (Table 2, Analyses 1 – 3). The first axis of the RDA of the vegetation data (Table 2, Analysis 1) displayed a gradient of defoliation management (Fig. 2). Three groups of treatments with similar plant species composition were identified on the ordination diagram: CO, GL+ and GL- treatments as the first group; HL+ and HL- treatments as the second group; and HGL+ and HGL- as the third group. The first group was species poor (number of species <12) and favoured grasses (especially *Poa pratensis*, *Agrostis capillaris*, *Festuca rubra*) and *Urtica dioica*. The second group was medium species rich (number of species about 17), and

included *Alopecurus pratensis*, *Cerastium holosteoides*, *Cirsium palustre*, *Holcus mollis*. The third group was the most species rich (number of species >19) and included the forb species *Betonica officinalis*, *Hypochoeris radicata*, *Potentilla erecta*, *Ranunculus acris*, *Taraxacum* spp.

Based on LMM results the cover of graminoids was significantly supported by grazing only management regimes (CO, GL-, GL+) regardless of fertiliser and liming inputs (Table 1). The total cover of graminoids in these treatments ranged from 112.8 ± 3.9 % (GL-) to 117.3 ± 1.6 % (GL+), whereas the cover in the treatments that incorporated cutting ranged from 46.0 ± 3.9 % (HGL) to 58.6 ± 4.0 % (HGL+). Grazing managements (CO, GL+, GL-) supported the dominant grasses *A. capillaris* and *P. pratensis*. While there was a tendency for *Anthoxanthum odoratum* to be suppressed by fertilizer applications (CO), *Festuca rubra* tended to be supported by a grazing only management.

Unlike graminoids, forbs were significantly suppressed by grazing only management (CO, GL-, GL+), regardless of fertilization and liming (Table 1). The total cover of forbs for the grazed treatments ranged from 2.5 ± 1.5 % (GL+) to 3.3 ± 1.1 % (GL-), whereas for the cut treatments the total cover ranged from 59.5 ± 4.8 % (HGL+) to 65.7 ± 3.3 % (HL+). The main species (*Plantago lanceolata*, *Ranunculus repens*, *Rumex acetosa*), as well as other less abundant species (such as *Ranunculus acris*, *C. holosteoides*, *Crepis capillaris*, *Leontodon autumnalis*, *Rhinanthus minor* and *Taraxacum* spp.), were supported by cutting management regimes, whereas *Trifolium repens* cover was not influenced by defoliation method.

3.3. Soil chemical properties

Redundancy analysis indicated that management regime explained the highest proportion of the soil nutrient content variability in the first and all axes (Table 2, Analyses 4

– 6). The first axis of RDA displayed a gradient of soil pH (Fig. 3). Soil from the plots that had received lime had a higher pH and plant available concentrations of Ca and Mg. Plant available concentrations of P, K and organic C were positively correlated with the CO treatment fertilized by N, P and K. Liming plot treatment was the second best explanatory predictor for soil chemical properties (Table 2, Analyses 4 – 6).

Results from LMM show that management regime did not influence the concentrations of N_{tot} , C_{org} , or the C:N ratio in the soil (Table 3). Concentrations of K in the soil had a tendency ($P=0.040$) to be higher in the grazing only managements. Liming influenced concentrations of Ca and Mg in the soil. The concentrations of Ca were highest in all formerly limed treatments, and a similar response was found for Mg. The highest concentration of P was found in CO plots. Plant species richness was negatively correlated with soil K concentration ($P = 0.015$, $r = -0.52$), however, there was only a trend for P concentration in the soil to impact species richness. Concentrations of other soil nutrients did not influence species richness.

4. Discussion

4.1. Species richness and plant species composition

Our results showed that type of defoliation (cutting, grazing, and their combination) influenced plant species composition, species richness and Shannon (H) species diversity indices more than fertilizer applications (CO treatment) or liming. This is in accordance with previous studies that found that changes in species composition (Köhler et al., 2001) and species richness (Parr and Way, 1988) were more affected by type of disturbance than by minor changes in soil nutrients.

The grazing only management regimes encouraged an increase in grasses at the expense of forbs, regardless of fertilizer applications and liming, leading to decreasing species richness and Shannon (H) species diversity indices. In contrast, hay cutting and hay cutting followed by aftermath grazing showed the highest species richness, linked to increased forb cover, which was likely the result of there being no sward disturbance until the middle of July. Dominance by grasses is a typical response of temperate grassland to frequent sward defoliation (Louault et al., 2005; Pavlů et al., 2007; Ludvíková et al., 2015). Likewise, grazing management in comparison to cutting has been shown to promote dominance of grasses at the expense of forbs (Krahulec et al., 2001; Dumont et al., 2011; Mládková et al. 2015), and therefore mowing is generally advocated in situations where species richness maintenance is the main goal of sward management (Hansson and Fogelfors, 2000). However, our data show that similar outcomes can be obtained by hay cutting followed by aftermath grazing. Although liming has been shown to frequently have positive effects on species richness in grasslands (de Graaf et al., 1998; Holland et al. 2018), this was not evident during the current study. It is likely that this is a result of the vegetation found in the surroundings of the experiment being adapted to growth in the acidic brown podzolic soils at the site.

4.2. Soil chemical properties

Although it is generally considered that P and N_{tot} concentrations in the soil have negative effects on grassland diversity (Janssens et al., 1998; Crawley et al., 2005; Hejcman et al., 2010; Pruchniewicz and Żołnierz, 2014), the results of our experiment show that N did not have any effect on plant species diversity. Low soil P concentrations in the locality were likely responsible for only a negative tendency of P concentration in the soil to influence species richness. Furthermore, Janssen et al. (1998) hypothesised that while N is the main

nutrient limiting plant diversity, its availability is controlled by P.

It is assumed that K concentrations in the soil do not strongly influence plant species richness (Crawley, 2005) and that a higher K concentration is compatible with high levels of sward diversity (Janssens et al., 1998). In our experiment, K concentration in the soil had a tendency to be higher not only in the control treatment with fertilizer applications, but also in both other grazing only treatments, where a proportion of the nutrients removed were returned in sheep excreta. Cutting with biomass removal resulted in a decrease in soil K concentrations, as has been reported previously (Parr & Way, 1988; Schaffers et al., 1998; Alfaro et al., 2003, 2004; Hejerman et al., 2010; Pavlů et al., 2013; Pavlů et al., 2016). The negative relationship between plant-available K and species richness was likely due to the relatively high concentrations of K, even after the long-term removal of herbage biomass from cut plots. So, although it appears that reductions in species richness are connected to higher soil K concentrations, these decreases may be predominantly linked, once again, to defoliation management.

4.3. Management implications

Restoration of improved permanent pasture is a long-term process that can be successful if several conditions (as summarized by van Diggelen and Marrs, 2003) can be fulfilled: (i) abiotic resource levels are within thresholds of target communities and species, (ii) sufficient viable propagules of target species are available at a rate that supports rapid establishment of desired species, and (iii) appropriate management regimes well adapted to target species requirements are in place.

Although it is generally believed that extensive grazing is a suitable management technique to maintain and restore plant diversity of temperate grassland (e.g. van Diggelen

and Marrs, 2003; Marriot et al., 2009; Jacquemyn et al., 2011; Ludvíková et al., 2015; Moinardeau et al., 2016) it is necessary to consider factors such as type of grazing animal and the grazing system. As indicated previously, sheep can be highly selective grazers and consistently sort the best quality components from within multi-species swards (Garcia et al., 2003). Thus, a low species richness is typical of swards managed by continuous sheep grazing, regardless of grazing intensity (Marriot et al., 2009). While grazing by less selective stock is preferable, economic and socioeconomic pressures have led to a reduction in cattle numbers within marginal areas of the UK and similar regions within the EU, despite targeted support within agri-environment schemes.

While the current study has established the comparative benefits to plant species richness of hay cutting, the practicality of this management option in upland fringe areas is commonly compromised by topography, terrain and climatic conditions (especially rainfall). Alternatives based on delayed or rotational grazing systems should be explored if reliance on support payments to offset reduced productivity is to be avoided.

5. Conclusion

Measurements carried out on this long-term restoration experiment provide several clear messages. The first is that a higher species richness with a high proportion of forbs was observed on treatments in which cutting with biomass removal is included, whereas treatments with only grazing were linked to low species richness with a higher proportion of grasses in the sward. Secondly, K was the only element whose plant available concentration in the soil tended to decrease in response to cutting treatments with herbage removal. Higher concentrations of Ca and Mg in the soil in treatments with former liming had no effect on species richness and plant species composition. Finally, continuous sheep grazing is not

recommended as a means of maintaining or restoring plant diversity within temperate grasslands. Postponing the timing of the first defoliation (hay cutting) to mid growing season, thus allowing forbs to reach the reproductive stage, would be the most effective restoration management option for upland grassland.

CrediT authorship contribution statement

Lenka Pavlů: Investigation, Conceptualization, Methodology, Writing-Original draft. **Vilém V. Pavlů:** Investigation, Formal analysis, Visualization, Writing-Original Draft. **Mariecia D. Fraser:** Investigation, Resources, Supervision, Writing- Reviewing and Editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This study was supported by Natural Resources Wales, a Stapledon Memorial Trust Travel Fellowship awarded to V. P. and funding from the Biotechnology and Biological Sciences Research Council (BBS/E/W/0012843C). Manuscript finalisation for L.P. and V.P. was supported by the Ministry of Agriculture of the Czech Republic (RO0417). The Ecological Continuity Trust's support of the Brignant plots was also appreciated. Thanks to J.E. Vale and J. Corton for technical assistance.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper in the field.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at

XXX

References

ADAS, 1983. *Lime and Fertiliser Recommendations*. No. 5. Grass and Forage Crops.

Ministry of Agriculture, Fisheries and Food Booklet 2430. Northumberland, MAFF, UK.

AOAC, 1984. 14th ed. *Official Methods of Analysis*. Association of Official Analytical Chemists, Washington.

Alfaro, M., Jarvis, S., Gregory, P., 2003. Potassium budgets in grassland systems as affected by nitrogen and drainage. *Soil Use Manage.* 19, 89-95. DOI: 10.1111/j.1475-2743.2003.tb00286.x

Alfaro, M., Jarvis, S., Gregory, P., 2004. Factors affecting potassium leaching in different soils. *Soil Use Manage.* 20, 182-189. DOI: 10.1111/j.1475-2743.2004.tb00355.x

Bakker, J.P., 1989. *Nature management by grazing and cutting*. Kluwer Academic Publishers, Dordrecht, NL.

Bakker, J.P., Elzinga, J.A., de Vries, Y., 2002. Effects of long-term cutting in a grassland system: perspectives for restoration of plant communities on nutrient-poor soils. *Appl. Veg. Sci.* 5, 107–120.

Begon, M., Townsend, C.R., Harper, J.L. (Eds.), 2005 *Ecology: From Individuals to Ecosystems*. Oxford, UK: Wiley-Blackwell.

403 Bonari, G, Fajmon, K., Malenovský, I., Zelený, D., Holuša, J., Jongepierová, I., Kočárek, P.,
 404 Konvička, O., Uříčář, J., Chytrý, M., 2017. Management of semi-natural grasslands
 405 benefiting both plant and insect diversity: The importance of heterogeneity and tradition.
 406 Agric. Ecosyst. Environ. 246, 243-256. <http://dx.doi.org/10.1016/j.agee.2017.06.010>
 407 Crawley, M.J., Johnston, A.E., Silvertown, J., Dodd, M., de Mazancourt, C., Heard, M.S.,
 408 Henman, D.F., Edwards, G.R., 2005. Determinants of Species Richness in the Park Grass
 409 Experiment. Am. Nat. 10, 179-192. DOI: 10.1086/427270
 410 Dell Inc., 2016. *Dell Statistica (data analysis software system)*, version 13.1 software.
 411 dell.com.
 412 Dumont, B., Carrère, P., Giname, C., Farruggia, A., Lanore, L., Tardif, A., Decuq, F.,
 413 Darsonville, O., Louault, F., 2011. Plant-herbivore interactions affect the initial direction of
 414 community changes in an ecosystem manipulation experiment. Basic Appl. Ecol. 12, 187-
 415 194. DOI: [10.1016/j.baae.2011.02.011](https://doi.org/10.1016/j.baae.2011.02.011)
 416 Garcia, F., Carrere, P., Soussana, J.F., Baumont, R., 2003. How do severity and frequency of
 417 grazing affect sward characteristics and the choices of sheep during the grazing season? Grass
 418 Forage Sci. 58, 138-150. DOI: [10.1046/j.1365-2494.2003.00363.x](https://doi.org/10.1046/j.1365-2494.2003.00363.x)
 419 Goulding, K.W.T., 2016. Soil acidification and the importance of liming agricultural soils
 420 with particular reference to the United Kingdom. Soil Use Manag. 32, 390–399. doi:
 421 [10.1111/sum.12270](https://doi.org/10.1111/sum.12270)
 422 de Graaf, M.C., Verbeek, P.J., Bobbink, R., Roelofs, J.G., 1998. Restoration of species-rich
 423 dry heaths: the importance of appropriate soil conditions. Acta Bot. Neerlandica 47,
 424 89-111.
 425 Hansson, M., Fogelfors, H., 2000. Management of a semi-natural grassland; results from 15-
 426 year-old experiment in southern Sweden. J. Veg. Sci. 11, 31-38. DOI: [10.2307/3236772](https://doi.org/10.2307/3236772)

427 Hayes, M.J., Tallowin, J.R.B., 2007. Recreating biodiverse grasslands: long-term evaluation
 428 of practical management options for farmers. In J.J., Hopkins, A.J. Duncan, D.I., McCracken,
 429 S., Peel & J.R.B., Tallowin (Eds.) *High Value Grassland: Providing Biodiversity, a Clean*
 430 *Environment and Premium Products. British Grassland Society Occasional Symposium*
 431 *No.38*. British Grassland Society (BGS) (pp. 135-140), Reading, UK.

432 Hejzman, M., Češková, M., Schellberg, J., Pätzold, S., 2010. The Rengen grassland
 433 experiment: Effect of soil chemical properties on biomass production, plant species
 434 composition and species richness. *Folia Geobot.* 45, 125-142. DOI: 10.1007/s12224-010-
 435 9062-9

436 Holland, J.E., Bennett, A.E., Newton, A.C., White, P.J., McKenzie, B.M., George, T.S.,
 437 Pakeman, R.J., Bailey, J.S., Fornara, D.A., Hayes, R.C., 2018. Liming impacts on
 438 soils, crops and biodiversity in the UK: a review. *Sci. Total Environ.* 610-611,
 439 316–332. doi.org/10.1016/j.scitotenv.2017.08.020

440 Humbert, J-Y., Dwyer, J.M., Andrey, A., Raphaël, A., 2016. Impacts of nitrogen addition on
 441 plant biodiversity in mountain grasslands depend on dose, application duration and climate: a
 442 systematic review. *Glob. Change Biol.* 22, 110-120. doi: 10.1111/gcb.12986

443 Janssens, F., Peeters, A., Tallowin, J.R.B., Bakker, J.P., Bekker, R.M., Fillat, F., Oomes,
 444 M.J.M., 1998. Relationship between soil chemical factors and grassland diversity. *Plant Soil.*
 445 202, 69-78. DOI: 10.1023/A:1004389614865

446 Jacquemyn, H., Van Mechelen, C., Brys, R., Honnay, O., 2011. Management effects on the
 447 vegetation and soil seed bank of calcareous grasslands: An 11-year experiment. *Biol.*
 448 *Conserv.* 144, 416-422. DOI: 10.1016/j.biocon.2010.09.020

449 Jongepierová, I., Prach, K., Tichý, L., Malenovský, I., 2018. Secondary grasslands and
 450 heatlands, in: Jongepierová, I., Pešout, P., Prach, K., (Eds.) *Ecological restoration in the*
 451 *Czech Republic II*. Nature Conservation Agency of the Czech Republic, Prague, pp. 63-71. .

452 Klaus, V.H., Hölzel, N., Boch, S., Müller, J., Socher, S.A., Prati, D., Fischer, M.,
 453 Kleinebecker, T., 2013. Direct and indirect associations between plant species richness and
 454 productivity in grasslands: regional differences preclude simple generalization of
 455 productivity-biodiversity relationships. *Preslia*. 85, 97-112. DOI: 10.7892/boris.38709
 456 Klaus, V.H., Kleinebecker, T., Hölzel, N., Blüthgen, N., Boch, S., Müller, J., Socher, S.A.,
 457 Prati, D., Fischer, M., 2011. Nutrient concentrations and fibre contents of plant community
 458 biomass reflect species richness patterns along a broad range of land-use intensities among
 459 agricultural grasslands. *Perspect. Plant Ecol. Syst.* 13, 287-295.
 460 doi:10.1016/j.ppees.2011.07.001
 461 Köhler, B., Ryser, P., Güsewell, S., Gigon, A., 2001. Nutrient availability and limitation in
 462 traditionally mown and in abandoned limestone grasslands: a bioassay experiment. *Plant Soil*.
 463 230, 323-332. DOI: b>10.1023/A:1010335825818
 464 Krahulec, F., Skálová, H., Herben, T., Hadincová, V., Wildová, R., Pecháčková, S.,
 465 2001. Vegetation changes following sheep grazing in abandoned mountain meadows. *Appl.*
 466 *Veg. Sci.* 4, 97-102. doi:10.1111/j.1654-109X.2001.tb00239.x
 467 Kubát, K., Hrouda, L., Chrtěk, J., Kaplan, Z., Kirschner, J., Štěpánek, J. (Eds.) 2002. Klíč ke
 468 květeně České republiky (Key to the flora of the Czech Republic). Academia, Prague, CZ.
 469 Louault, F., Pillar, V.D., Aufrère, J., Garnier, E., Soussana, J.F., 2005. Plant traits and
 470 functional types in response to reduced disturbance in a semi-natural grassland. *J. Veg. Sci.*,
 471 16, 151-161. DOI: 10.1658/1100-9233(2005)016[0151:PTAFTI]2.0.CO;2
 472 Ludvíková, V., Pavlů, V., Pavlů, L., Gaisler, J., Hejcman M., 2015. Sward-height patches
 473 under intensive and extensive grazing density in *Agrostis capillaris* grassland. *Folia Geobot.*
 474 50, 219-228. DOI 10.1007/s12224-015-9215-y
 475 Marriott, C.A., Hood, K., Fisher, J.M., Pakeman, R.J., 2009. Long-term impacts of extensive
 476 grazing and abandonment on the species composition, richness, diversity and productivity of

477 agricultural grassland. *Agric. Ecosyst. Environ.* 134 (3), 190-200. DOI:
 478 10.1016/j.agee.2009.07.002
 479 Mehlich, A., 1984. Mehlich 3 soil test extractant: a modification of Mehlich 2 extractant.
 480 *Commun. Soil Sci. Plant Anal.* 15, 1409-1416.
 481 Mládková, P., Mládek, J., Hejduk, S., Hejzman, M., Cruz, P., Jouany, C., Pakeman, R.J.,
 482 2015. High-nature-value grasslands have the capacity to cope with nutrient impoverishment
 483 induced by mowing and livestock grazing. *J. Appl. Ecol.* 52, 1073-1071. DOI: 10.1111/1365-
 484 2664.12464
 485 Moinardeau, C., Mesléard, F., Dutoit, T., 2016. Using Different Grazing Practices for
 486 Increasing Plant Biodiversity in the Dykes and Embankments Along the Rhône River
 487 (Southern France). *J. Environ. Manage.* 58, 984-997. DOI: 10.1007/s00267-016-0744-9
 488 Parr, T.W., Way, J.M., 1988. Management of roadside vegetation: the long-term effects of
 489 cutting. *J. Appl. Ecol.* 25, 1073-1087. DOI: 10.2307/2403767
 490 Pavlů, V., Hejzman, M., Pavlů, L., Gaisler, J., 2007. Restoration of grazing management and
 491 its effect on vegetation in an upland grassland. *Appl. Veg. Sci.* 10, 375-382. DOI:
 492 10.1111/j.1654-109X.2007.tb00436.x
 493 Pavlů, L., Gaisler, J., Hejzman, M., Pavlů, V., 2016. What is the effect of long-term mulching
 494 and traditional cutting regimes on soil and biomass chemical properties, species richness and
 495 herbage production in *Dactylis glomerata* grassland? *Agric. Ecosyst. Environ.* 217, 13-21.
 496 DOI: 10.1016/j.agee.2015.10.026
 497 Pavlů, L., Pavlů, V., Gaisler, J., Hejzman, M., 2013. Relationship between soil and biomass
 498 chemical properties, herbage yield and sward height in cut and unmanaged mountain hay
 499 meadow (*Polygono-Trisetion*). *Flora*, 208, 599-608. DOI: 10.1016/j.flora.2013.09.003
 500 Pavlů, V., Schellberg, J., Hejzman, M., 2011. Cutting frequency vs. N application: effect of a
 501 20-year management in *Lolio- Cynosuretum* grassland. *Grass Forage Sci.* 66, 501-515. DOI:

10.1111/j.1365-2494.2011.00807.x

Perring, M.P., Edwards, G., de Mazancourt, C., 2009. Removing phosphorus from ecosystems through nitrogen fertilization and cutting with removal of biomass. *Ecosystems* 12, 1130-1144. <http://www.jstor.org/stable/25622870>.

Pegtél, D. M., Akker, J. P., Verweij, G. L., Fresco, L. F. M., 1996. N, P and K deficiency in chronosequential cut summer – dry grasslands on gley podsol after the cessation of fertilizer application. *Plant Soil*. 178, 121-131.

Pruchniewicz, D., Żołnierz, L., 2014. The influence of environmental factors and management methods on the vegetation of mesic grasslands in a central European mountain. *Flora* 209, 687-692. DOI: 10.1016/j.flora.2014.09.001

Schaffers, A.P., Vesseur, M.C., Sýkora, K.V., 1998. Effects of delayed hay removal on the nutrient balance of roadside plant communities. *J. Appl. Ecol.* 35, 349-364. DOI: 10.1046/j.1365-2664.1998.00316.x

Smith, R. S., Shiel, R. S., Bardgett, R. D., Millward, D., Corkhill, P., Evans, P., Quirk, H., Hobbs, P. J., Kometa, S. T. 2008. Long – term change in vegetation and soil microbial communities during the phased restoration of traditional meadow grassland. *J. Appl. Ecol.* 45, 670-679. doi.org/10.1111/j.1365-2664.2007.01425.x

Spiegelberger T., Hegg O., Matthies D., Hedlund K., Schaffner U., 2006. Long-term effects of short-term perturbation in a subalpine grassland. *Ecology* 87, 1939-1944. DOI: org/10.1890/0012-9658(2006)87[1939:LEOSPI]2.0.CO;2

Storkey, J., Macdonald, A., Poulton, P., Scott, T., Köhler, I., Schnyder, H., Goulding, K., Crawley, M., 2015. Grassland biodiversity bounces back from long-term nitrogen addition. *Nature* 528, 40-404. doi:10.1038/nature16444

ter Braak, C.J.F., Šmilauer, P., 2012. *Canoco 5, Windows release (5.00). [Software for canonical community ordination]*. Microcomputer Power, Ithaca, NY, US.

527 Titěra, J., Pavlů, V.V., Pavlů, L., Hejman, M., Gaisler, J., Schellberg, J., 2020. Response of
528 grassland vegetation composition to different fertilizer treatments recorded over ten years
529 following 64 years of fertilizer applications in the Rengen Grassland Experiment. Appl. Veg.
530 Sci. <https://doi-org.infozdroje.czu.cz/10.1111/avsc.12499>.
531 Van Diggelen, R., Marrs, R., 2003. Restoring Plant Communities - Introduction. Appl. Veg.
532 Sci. 6, 106-110.
533 Verhoeven, K.J., Simonsen, K.L., McIntyre, L.M., 2005. Implementing false discovery rate
534 control: increasing your power. Oikos, 108, 643-647. doi:10.1111/j.0030-1299.2005.13727.x
535

536 **Table 1**

537 Cover (%) of the most abundant vascular plant species in alphabetical order; cover (%) of total graminoids and total forbs;
 538 number of all plants species and plant species $\geq 1\%$; Shannon index (H) and Shannon species evenness index (J) under the
 539 different treatments in 2012. Treatment abbreviations: CO = control, continuing the previous site management with liming,
 540 fertilising and grazing by sheep, GL+ = sheep grazing with liming application, GL- = sheep grazing without liming application,
 541 HL+ = hay cutting only with liming application, HL- = hay cutting only without liming application, HGL+ = hay cutting
 542 followed by aftermath sheep grazing with liming application, HGL- = hay cutting followed by aftermath sheep grazing
 543 without liming application. Numbers represent average of three replicates $\pm$ standard error of the mean (SE); F -ratio = F -
 544 statistics for the test of a particular analysis; P -value = corresponding probability value. In cases of significant differences
 545 obtained by linear mixed-effects modelling after table-wise Benjamini-Hochberg's FDR correction (highlighted in bold), a
 546 post hoc comparison using the Tukey's HSD test was applied to identify significant differences between treatments.
 547 Differences are indicated by different small letters within rows.

	Treatment								
	F -ratio	P -value	CO	GL+	GL-	HL+	HL-	HGL+	HGL-
Number of all plant species	15.43	<0.001	8.3 $\pm$ 1.9 c	9.3 $\pm$ 1.5 c	11.3 $\pm$ 0.3 bc	17.3 $\pm$ 1.2 ab	17.3 $\pm$ 0.9 ab	19.3 $\pm$ 1.2 a	20.3 $\pm$ 0.7 a
Number of plant species $\geq 1\%$	14.39	<0.001	6.0 $\pm$ 1.5 b	6.3 $\pm$ 0.3 b	6.7 $\pm$ 0.9 b	11.3 $\pm$ 1.2 a	11.7 $\pm$ 0.3 a	13.3 $\pm$ 0.3 a	14.0 $\pm$ 0.6 a
Shannon (H) diversity index	14.05	<0.001	1.4 $\pm$ 0.2 c	1.6 $\pm$ 0.1 c	1.7 $\pm$ 0.1 bc	2.2 $\pm$ 0.1 a	2.1 $\pm$ 0.1 ab	2.3 $\pm$ 0.1 a	2.4 $\pm$ 0.1 a
Shannon (J) evenness index	1.70	0.20	0.7 $\pm$ 0.04	0.7 $\pm$ 0.04	0.7 $\pm$ 0.05	0.8 $\pm$ 0.03	0.7 $\pm$ 0.01	0.8 $\pm$ 0.01	0.8 $\pm$ 0.01
Total graminoids	101.76	<0.001	116.6 $\pm$ 1.9 a	117.3 $\pm$ 1.6 a	112.8 $\pm$ 3.9 a	52.1 $\pm$ 3.8 b	55.6 $\pm$ 4.1 b	58.6 $\pm$ 4.0 b	46 $\pm$ 3.9 b
Total forbs	159.35	<0.001	2.8 $\pm$ 1.5 b	2.5 $\pm$ 1.5 b	3.3 $\pm$ 1.1 b	65.7 $\pm$ 3.3 a	62.2 $\pm$ 2.3 a	59.5 $\pm$ 4.8 a	63.3 $\pm$ 2.7 a
<i>Anthoxanthum odoratum</i>	4.08	0.018	0.43 $\pm$ 0.4 b	13.3 $\pm$ 5.2 ab	17.7 $\pm$ 2.4 a	10.1 $\pm$ 5.5 ab	12.2 $\pm$ 4.4 ab	10.3 $\pm$ 2.2 ab	8.9 $\pm$ 4.8 ab
<i>Agrostis capillaris</i>	4.93	0.009	32.5 $\pm$ 4.9 ab	36.0 $\pm$ 5.1 ab	43.2 $\pm$ 9.6 a	11.1 $\pm$ 2.3 b	25.2 $\pm$ 8.5 ab	18.9 $\pm$ 4.4 ab	15.8 $\pm$ 3.6 b
<i>Cynosurus cristatus</i>	2.26	0.10	0.0	0.7 $\pm$ 0.7	0.3 $\pm$ 0.3	1.1 $\pm$ 0.95	0.4 $\pm$ 0.4	3.3 $\pm$ 1.3	0.4 $\pm$ 0.2
<i>Festuca rubra</i>	3.26	0.038	29.2 $\pm$ 4.5	28.7 $\pm$ 0.7	21.8 $\pm$ 3.8	18.6 $\pm$ 2.1	12.0 $\pm$ 6.1	16.9 $\pm$ 6.5	14.4 $\pm$ 5.0
<i>Holcus lanatus</i>	3.19	0.041	8.9 $\pm$ 2.9	10.2 $\pm$ 0.6	7.4 $\pm$ 0.4	6.1 $\pm$ 1.1	3.0 $\pm$ 2.02	6.7 $\pm$ 0.6	3.0 $\pm$ 1.5
<i>Lolium perenne</i>	1.84	0.171	1.8 $\pm$ 1.8	0.3 $\pm$ 0.3	5.6 $\pm$ 4.0	0.0	0.0	0.03 $\pm$ 0.03	0.003 $\pm$ 0.003
<i>Poa pratensis</i>	7.15	0.002	43.7 $\pm$ 14.2 a	28.2 $\pm$ 8.4 a	16.5 $\pm$ 5.6 a	2.0 $\pm$ 1.1 b	1.4 $\pm$ 0.9 b	1.1 $\pm$ 0.6 b	1.0 $\pm$ 0.3 b
<i>Trifolium repens</i>	1.01	0.457	0.4 $\pm$ 0.4	1.5 $\pm$ 1.2	1.1 $\pm$ 0.7	0.0	0.0	3.0 $\pm$ 2.3	0.8 $\pm$ 0.4

<i>Cerastium holosteoides</i>	3.33	0.036	0.0	0.0	0.0	0.5 ± 0.3	0.9 ± 0.3	0.13 ± 0.13	0.4 ± 0.4
<i>Crepis capillaris</i>	2.10	0.129	0.0	0.0	0.0	6.3 ± 4.1	1.7 ± 1.7	3.9 ± 1.9	5.3 ± 2.7
<i>Leontodon autumnalis</i>	1.97	0.148	0.0	0.0	0.0	3.9 ± 3.9	7.0 ± 3.6	1.1 ± 1.1	0.2 ± 0.2
<i>Plantago lanceolata</i>	10.53	<0.001	0.0 b	0.0 b	0.0 b	9.3 ± 3.9 ab	14.6 ± 2.1 ab	19.7 ± 6.0 a	22.0 ± 1.1 a
<i>Ranunculus acris</i>	8.98	<0.001	0.0 b	0.03 ± 0.03 b	0.0 b	2.7 ± 1.4 ab	1.4 ± 0.8 b	3.0 ± 0.4 ab	5.4 ± 1.3 a
<i>Ranunculus repens</i>	10.28	<0.001	0.1 ± 0.1 c	0.03 ± 0.03 c	0.4 ± 0.1 c	25.5 ± 7.0 a	18.6 ± 4.3 ab	13.1 ± 3.8 abc	8.3 ± 2.3 bc
<i>Rhinanthus minor</i>	1.24	0.35	0.0	0.0	0.0	0.6 ± 0.5	1.9 ± 1.9	0.3 ± 0.2	3.2 ± 2.6
<i>Rumex acetosa</i>	31.93	<0.001	1.0 ± 0.6 c	0.1 ± 0.1 c	1.4 ± 1.3 c	16.2 ± 2.4 a	14.7 ± 1.0 a	12.1 ± 2.6 ab	7.4 ± 1.6 b
<i>Taraxacum</i> spp.	2.33	0.099	0.0	0.0	0.0	0.4 ± 0.2	0.5 ± 0.2	1.3 ± 0.6	1.7 ± 1.0

548

549

Table 2

Results of redundancy analyses for six different H0 analyses (A1-A3 for vegetation, A4-A6 for soil); % expl. = explained variation by axis 1 (adjusted explained variation by all ordination axes), a measure of the explanatory power of the explanatory variables; *F*-ratio = *F*-statistics for the test of a particular analysis; *P*-value = corresponding probability value obtained by the Monte Carlo permutation test. Treatment abbreviations (CO, GL+, GL, HL+, HL-, HGL+, HGL-) are defined in Table 1.

Analysis	Expl. var.	Covariables	% expl.	<i>F</i> -ratio	<i>P</i> -value
Vegetation					
A1 Different grassland managed regimes have no effect on plant species composition	CO, HGL+, HGL-, HL+, HL-, GL+, GL-	blocks	58.8 (77.2)	18.6 (7.3)	0.001 (0.001)
A2 Different defoliation management regimes have no effect on plant species composition	Grazing, Hay, Grazing+Hay	blocks	58.6 (65.3)	24.0 (16.0)	0.001 (0.001)
A3 Liming has no effect on plant species composition	Limed+, Limed-	blocks	5.8	1.1	0.309
Soil					
A4 Different grassland managed regimes have no effect on soil properties	CO, HGL+, HGL-, HL+, HL-, GL+, GL-	blocks	44.8 (65.5)	10.6 (4.1)	0.002 (0.001)
A5 Different defoliation management regimes have no effect on soil properties	Grazing, Hay, Grazing+Hay	blocks	13.5 (15.4)	2.7 (1.5)	0.167 (0.179)
A6 Liming has no effect on soil properties	Limed+	blocks	42.3	13.2	0.001

557 **Table 3**

558 Soil characteristics per plot under the different treatments in 2012. Treatment abbreviations (CO, GL+, GL, HL+, HL-, HGL+,
559 HGL-) are defined in Table 1. Numbers represent average of three replicates $\pm$ standard error of the mean (SE); F -ratio = F -
560 statistics for the test of a particular analysis; P -value = corresponding probability value. In cases of significant differences
561 obtained by linear mixed-effects modelling after table-wise Benjamini-Hochberg's FDR correction (highlighted in bold), the
562 post hoc comparison using the Tukey's HSD test was applied to identify significant differences between treatments.
563 Differences are indicated by different small letters within rows.

				Treatment						
	Characteristics	F -ratio	P -value	CO	GL+	GL-	HL+	HL-	HGL+	HGL-
Soil up to 7.5 cm	pH	12.78	<0.001	4.73 $\pm$ 0.05 ab	4.89 $\pm$ 0.03 a	4.54 $\pm$ 0.10 bc	4.91 $\pm$ 0.01 a	4.41 $\pm$ 0.09 c	4.74 $\pm$ 0.05 ab	4.43 $\pm$ 0.04 c
	C _{org} (%)	1.10	0.430	8.28 $\pm$ 0.66	7.18 $\pm$ 0.77	7.46 $\pm$ 0.19	6.83 $\pm$ 0.18	7.02 $\pm$ 0.27	7.51 $\pm$ 0.16	7.27 $\pm$ 0.13
	N _{tot} (%)	0.41	0.854	0.51 $\pm$ 0.11	0.53 $\pm$ 0.05	0.56 $\pm$ 0.02	0.66 $\pm$ 0.11	0.55 $\pm$ 0.03	0.58 $\pm$ 0.01	0.57 $\pm$ 0.02
	P mg kg ⁻¹	4.11	0.018	25.01 $\pm$ 5.38 a	13.78 $\pm$ 1.96 b	16.00 $\pm$ 1.67 ab	16.65 $\pm$ 1.02 ab	14.57 $\pm$ 0.24 b	13.16 $\pm$ 1.09 b	13.28 $\pm$ 0.73 b
	K mg kg ⁻¹	3.22	0.040	172.53 $\pm$ 7.11	153.07 $\pm$ 17.27	175.91 $\pm$ 24.86	121.63 $\pm$ 9.26	121.72 $\pm$ 4.95	141.47 $\pm$ 3.99	122.62 $\pm$ 6.41
	Ca mg kg ⁻¹	12.62	<0.001	1395 $\pm$ 136 a	1390 $\pm$ 115 a	813 $\pm$ 112 bc	1415 $\pm$ 123 a	773 $\pm$ 213 c	1261 $\pm$ 73 ab	714 $\pm$ 144 c
	Mg mg kg ⁻¹	6.75	0.002	278.83 $\pm$ 17.92 ab	290.38 $\pm$ 25.82 a	207.31 $\pm$ 17.36 abc	302.16 $\pm$ 32.70 a	183.31 $\pm$ 32.39 bc	269.93 $\pm$ 20.57 abc	177.63 $\pm$ 17.81 c
	C:N	2.07	0.103	18.0 $\pm$ 3.6	13.7 $\pm$ 0.7	13.3 $\pm$ 0.8	11.6 $\pm$ 1.8	12.9 $\pm$ 0.5	12.9 $\pm$ 0.2	12.8 $\pm$ 0.4

564

565

Fig. 1. Species richness under different defoliation management regimes: sheep grazing, hay cutting, hay cutting followed by aftermath sheep grazing. Error bars represent standard error of the mean. Significant differences ($P < 0.05$) according to the Tukey post-hoc test are indicated by different letters.

Fig. 2. Ordination diagram representing the results of redundancy analysis showing changes in plant species composition, treatments were used as predictors. (see Table 2, Analysis 1 for details).

Treatment abbreviations: CO = control, continuing the previous site management with liming, fertilising and grazing by sheep, GL+ = sheep grazing with liming application, GL- = sheep grazing without liming application, HL+ = hay cutting only with liming application, HL- = hay cutting only without liming application, HGL+ = hay cutting followed by aftermath sheep grazing with liming application, HGL- = hay cutting followed by aftermath sheep grazing without liming application. Species abbreviations are based on the first four-letter of genera and the four-letter of species name: Achilmile = *Achillea millefolium*, Agrocapi = *Agrostis capillaris*, Aloppra = *Alopecurus pratensis*, Anthodor = *Anthoxanthum odoratum*, Betooffi = *Betonica officinalis*, Brommoll = *Bromus mollis*, Cardprat = *Cardamine pratensis*, Ceraholo = *Cerastium holosteoides*, Cirspalu = *Cirsium palustre*, Crepcapi = *Crepis capillaris*, Cynocris = *Cynosurus cristatus*, Euphspp = *Euphorbia spp.*, Festrubr = *Festuca rubra*, Holclana = *Holcus lanatus*, Holcmoll = *Holcus mollis*, Hyporadi = *Hypochoeris radicata*, Leonautu = *Leotodon autumnalis*, Lolipere = *Lolium perenne*, Luzucamp = *Luzula campestris*, Planlanc = *Plantago lanceolata*, Phleprat = *Phleum pratense*, Poaprat = *Poa pratensis*, Poteerec = *Potentilla erecta*, Ranuacri = *Ranunculus acris*, Ranurepe = *Ranunculus repens*, Rhinmino = *Rhinanthus minor*, Rumeacet = *Rumex acetosa*, Stelgram =

591 *Stellaria graminea*, *Taraspp* = *Taraxacum spp.*, *Triferep* = *Trifolium repens*, *Veroserp* =
592 *Veronica serpyllifolia*, *Veroarve* = *Veronica arvensis*, *Urtidloi* = *Urtica dioica*
593

594 **Fig. 3.** Ordination diagram representing the results of redundancy analysis showing changes
595 in nutrient concentrations in the soil at a depth of 0-7.5 cm, treatments were used as
596 predictors. (see Table 2, Analysis 4 for details). Treatment abbreviations (CO, GL+, GL,
597 HL+, HL-, HGL+, HGL-) are defined in Figure 1. Abbreviations: pH-soil acidity, Corg-
598 organic carbon; Ntot-total nitrogen in the soil; P, K, Mg, Ca-plant available nutrients; C:N-
599 ratio in the soil

600
601
602
603
604
605
606
607

608
609

610
611
612

Figure 1

613
614

615 Figure 2
616

617
618
619

620 Figure 3