Aberystwyth University

Incipient genetic isolation of a temperate migratory coastal Sciaenid fish (Argyrosomus inodorus) within the Benguela Cold Current system

Henriques, Romina; Potts, Warren; Sauer, Warwick; Shaw, Paul

Published in:
Marine Biology Research

DOI:
[10.1080/17451000.2014.952309](https://doi.org/10.1080/17451000.2014.952309)

Publication date:
2015

Citation for published version (APA):

Henriques, R., Potts, W., Sauer, W., & Shaw, P. (2015). Incipient genetic isolation of a temperate migratory coastal Sciaenid fish (*Argyrosomus inodorus*) within the Benguela Cold Current system. *Marine Biology Research*, 11(4), 423-429. <https://doi.org/10.1080/17451000.2014.952309>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Title: Incipient genetic isolation of a temperate migratory coastal Sciaenid fish (*A. inodorus*) within the Benguela Cold Current system

Authors: Romina Henriques^{1,2}, Warren M. Potts², Warwick H.H. Sauer², Paul W. Shaw^{1,3}

Affiliations: ¹Centre for Ecology, Evolution and Behavior, School of Biological Sciences, Royal Holloway University of London, Egham, U.K.; ²Department of Ichthyology and Fisheries Science, Rhodes University, Grahamstown, South Africa; ³Institute of Biological, Environmental and Rural Sciences (IBERS), Aberystwyth University, Aberystwyth, U.K.

Corresponding author and present address:

Romina Henriques
Evolutionary Genomics Group
Department of Botany and Zoology
Stellenbosch University - Private Bag X1
Matieland
7602
South Africa
Email: rhenriques@sun.ac.za

This work was supported by the Fundação para a Ciência e Tecnologia (FCT – Portugal), under Grant number SFRH/BD/36176/2007.

Abstract

The Benguela Current is considered to be a major biogeographical barrier for tropical and warm-temperate marine fish, but there is limited knowledge regarding its influence on population sub-structuring of cold-tolerant species. Employing genetic variation within the mitochondrial DNA Control Region and six cross-specific nuclear microsatellite markers, a preliminary study was conducted to investigate population substructuring in *Argyrosomus inodorus*, a highly exploited, cool-temperate migratory species, across the Benguela Current region. Results revealed evidence of incipient genetic differentiation (mtDNA $\phi_{ST} = 0.092$; nuclear $F_{ST} = 0.036$ and $D_{ST} = 0.104$, $P < 0.005$) between the two sampling sites, suggesting the presence of two regional populations. Estimates of contemporary migration rates between populations were low, and similar in range to those reported in tagging surveys. Although preliminary, these results suggest that the oceanographic features of the Benguela Current may have influenced the evolutionary history of *A. inodorus*, and that the species is likely to be composed of two populations in the Benguela region. As the species is considered overexploited both in Namibia and South Africa, information on the distribution, population dynamics and long-term dispersal patterns across the Benguela Current region would provide a comprehensive evaluation of genetic structure, which should be incorporated into fishery management arrangements. .

Key-words: *Argyrosomus inodorus*, population structure, Benguela Current, isolation

Introduction

The Benguela Cold Current system, located in the southern Atlantic, features cold sea surface temperatures bounded to the north and south by tropical currents (the Angola and Agulhas currents, respectively), and a perennial upwelling cell off central Namibia that divides the region into two sub-systems with different characteristics (Shannon 1985; Hutchings et al. 2009). The colder sea surface temperatures of the Benguela Current have been considered an important bio-geographical barrier, isolating tropical and warm-temperate fauna of the Atlantic and Indo-Pacific Oceans (Avisé 2000; Floeter et al. 2008). However, recent studies revealed that other oceanographic features such as the perennial upwelling cell may also play an important role in shaping the population structure of warm-temperate fish populations within the Benguela system, as complete disruption of gene flow was documented both in *Lichia amia* and *Atractoscion aequidens* (Henriques et al. 2012; 2014). Little is known, however, regarding the influence of the Benguela system on genetic population connectivity of cold water tolerant species.

Argyrosomus inodorus is a migratory, benthopelagic sciaenid fish, endemic to the southeastern Atlantic (Griffiths & Heemstra 1995). Distribution range is restricted to cold-temperate waters (13°C-16°C), from the nearshore environment to depths of 100m, between Cape Frio and Meob Bay in Namibia, and between Cape Point and East London in South Africa (Griffiths & Heemstra 1995; Griffiths 1997; Kirchner & Holtzhausen 2001). The species distribution overlaps with those of the congeneric *A. coronus* in northern Namibia, and with *A. japonicus* along the southern and Eastern Cape coasts of South Africa (Griffiths & Heemstra 1995). Contrary to *A. inodorus*, *A. coronus* and *A. japonicus* are considered warm-temperate species, occurring preferentially in sea surface temperatures of 16°-19°C (Potts et al 2010) and 21°-25°C (Heemstra & Heemstra 2004), respectively. As *A. inodorus* is absent from the west coast of South Africa and there is no evidence for significant migration between the two areas of occurrence (Kirchner & Holtzhausen 2001), the species has been managed as two independent stocks. Life history characteristics appear to corroborate the hypothesis of two isolated and locally adapted populations, as features such as maximum size and size at maturity of Namibian and South African *A. inodorus* are significantly different, although no evidence of differentiation was observed within either region (Griffiths 1997; Holtzhausen et al. 2001). *A. inodorus* is a critical component of multiple coastal

fishery sectors, and exploitation pressure throughout its distribution has led to the species becoming severely depleted, with spawning stocks estimated to be 69% of unexploited values (FAO 2012). In Namibia, *A. inodorus* is harvested by the commercial and recreational fishery sectors and although approximately the same numbers of fish are captured in each sector, the average size captured in the commercial sector is larger (Kirchner 1998). While *A. inodorus* is only regulated through an input control (number of permits) the recreational fishery catch, which comprises 70% of the total recreational catch, is regulated through bag- and size-limits (Kirchner & Beyer, 1999; Holtzhausen et al. 2001). In South Africa, a 2012 survey reported that total landings of *A. inodorus* exceeded 400t per year (DAFF 2012). To establish sustainable management measures, it is necessary to understand how *A. inodorus* populations are structured across the Benguela Current region and whether migration between the two centers is absent. To date, no comprehensive genetic survey has been carried out in *A. inodorus*, with the exception of a genetic identification study in 1997 to differentiate between *A. inodorus* and *A. coronus*, based on allozymes (van der Bank & Kirchner 1997), and a more recent study on shifts of abundance of these two species in central Namibia (Potts et al. 2014).

The distribution range and life history features of this species suggest that, as observed for warm-temperate species, the oceanographic features of the Benguela Current may influence the population structure and gene flow across the region. The aim of this study was to conduct a preliminary assessment of genetic diversity, population substructuring and connectivity between the two putative populations of *A. inodorus* across the Benguela Current, using both mitochondrial DNA (mtDNA) and nuclear microsatellite DNA markers, in order to test whether the regional oceanographic features influence population connectivity in this cold-temperate fish species.

Methods

Sampling

A total of 80 fish were captured by rod-and-line fishing from the shore, by local collaborators in two areas: the West Coast Recreational Area in Namibia ($n = 40$) and the Eastern Cape Province in South Africa ($n = 40$), representing the two centres of

abundance of the species (Figure 1). A clip of the pectoral fin was removed immediately after capture and stored in 95% ethanol.

Genetic screening

DNA extraction was performed using a standard phenol:chlorophorm method (Sambrook et al. 1989). Genetic variation was assessed as DNA sequence polymorphism in a fragment of the mtDNA Control Region (CR) and allele frequencies at six microsatellite loci isolated from *Argyrosomus japonicus* (Archangi et al. 2009). A total of 36 *A. inodorus* individuals were amplified by polymerase chain reaction (PCR) for CR, using primers and protocols of Appleyard et al. (2002). PCR products were purified with an enzymatic digestion, consisting of 0.5U of EXO1 (NewEngland biolabs) and 1u of shrimp alkaline phosphatase (SAP) in 1x supplied buffer (Fermentas), and sequenced in the forward direction using the same amplification primers, by Macrogen Inc. (South Korea). Sequences were visually inspected and a multiple alignment was performed in CLUSTAL X (Thompson et al. 1997), as implemented in BioEdit 7.0.1 (accession numbers: JX191998-2033).

Forty individuals per sampling site were screened at six microsatellite loci (UBA5, UBA40, UBA50, UBA91, UBA853 and UBA854). Optimized PCR mixes included 1x NH₄Cl buffer, 2mM of MgCl₂, 0.2mM of dNTPs, 0.5pmol of each primer, 0.2U of Taq polymerase (Bioline UK) and 50-100ng of extracted DNA, in a final volume of 10 µl. The Archangi et al. (2009) protocols were modified to ensure accurate amplification: annealing temperatures and number of cycles (UBA91 T_a = 52°C, remaining loci T_a = 48°C, with 35 cycles), and removal of the final extension step of 72°C for 10min. PCR fragments from multiple loci were combined and genotyped on an AB3500 Genetic Analyzer (Applied Biosystems). Alleles were scored as PCR product size in base pairs, and scores were determined against an internal size marker (LIZ 600), using GeneMapper 4.0 (ABIPrism). In order to ensure accurate allele size scoring between runs, individuals with known allele sizes were used in each run as positive controls.

Data analyses

The CR dataset was assessed for levels of haplotype (h) and nucleotide (π) diversity, and fits to neutrality tests: Ewens-Watson's F , Tajima's D and Fu's FS , as implemented in ARLEQUIN (Excoffier et al. 2005). Determination of the most suitable nucleotide substitution model was performed in jModelTest (Posada, 2008). Preliminary inference of population connectivity of *A. inodorus* across the Benguela Current region was estimated as ϕ_{ST} in ARLEQUIN (Excoffier et al. 2005), with a significance level of $P < 0.05$ determined by 10,000 permutations. Haplotype networks were reconstructed to evaluate intraspecific relationships among haplotypes, using the Median-Joining (MJ) algorithm implemented in NETWORK (Bandelt et al. 1999).

Microsatellite genotypic frequencies were tested for deviation from Hardy-Weinberg expectations of random mating and from linkage equilibrium, as implemented in GENEPOP (Raymond & Rousset 1995). The occurrence of amplification errors such as large allele drop out and stuttering, and estimation of null allele frequencies were assessed in MICROCHECKER (van Oosterhout et al. 2006). Levels of genetic diversity were estimated as number of alleles (N_a), allelic richness (AR), observed and expected heterozygosity (H_O and H_E), and Wright's inbreeding coefficient (F_{IS}), in ARLEQUIN (Excoffier et al. 2005). A preliminary analysis to investigate the statistical power of the dataset to infer population substructuring was conducted in POWSIM (Ryman & Palm 2006). Simulations were conducted for six loci and two populations ($n = 40$, $n = 40$), using the estimated allelic frequencies as the baseline for the ancestral population. Runs were performed using multiple combinations of effective population size (N_e) and number of generations (t) to generate a population differentiation of $F_{ST} = 0.05$, $F_{ST} = 0.02$ and $F_{ST} = 0.01$. Each simulation was run for 1,000 replicates, and power was estimated as the proportion of tests that indicated significant genetic divergence (Ryman & Palm, 2006). Genetic differentiation was measured as Weir & Cockerham (1984) F_{ST} estimator, as implemented in FreeNA (Chapuis & Estoup 2007), with significance and 95% confidence intervals estimated after jackknifing. For comparison purposes, genetic differentiation was also measure using Jost's D_{est} estimator, which is independent of the levels of genetic diversity, in SMOGD (Crawford 2010). Contemporary estimates of long-term average migration rates between the two sampling sites were performed for the microsatellite dataset using two complementary approaches: the classical method based on F_{ST} values ($F_{ST} = 1/(4N_{em} + 1)$) (Excoffier et al. 2005), and by employing the coalescent-based approach of MIGRATE (Beerli 2009). In MIGRATE, the Bayesian approach was implemented, enforcing a full migration model, with three replicates run

for each dataset (Beerli 2009). Each analysis was performed with four connected chains, using static heating (1,000,000, 3, 1.5, 1), a burn-in period of 10,000 steps, followed by 90,000 steps, and parameters were recorded every 100 steps. Estimates of migration rates (m) were obtained from M ($M = m\mu$) and Θ ($\Theta = 4N_e\mu$) (Beerli, 2009). In order to obtain estimates of migration rates per generation (and not scaled by mutation) three general mutation rates were used: 0.1%, 0.5% and 1% per generation.

Results

Population structure and phylogeography

Sequencing of mtDNA CR yielded a fragment of 704 base pairs (bp). The 36 individuals screened displayed 32 haplotypes defined by 34 variable nucleotide sites, of which 16 sites were parsimony informative. The Tamura-Nei nucleotide substitution model was identified as the most suitable for the mtDNA dataset. Haplotype diversity was high ($h = 0.991$), whilst nucleotide diversity was low ($\pi = 0.006$), with Namibian samples exhibiting higher values than the South African samples (Table 1). Deviations from the assumptions of selection neutrality were observed in Fu's FS for both populations, but not with either Ewens-Watsonson F or the Tajima's D tests (Table 1). As Fu's FS is known to be sensitive to abrupt demographic changes it is likely that the observed deviation to neutrality resulted from past population size changes, rather than reflecting selection effects. Genetic differentiation (ϕ_{ST}) between samples was low but statistically significant ($\phi_{ST} = 0.092$, $P < 0.05$), although haplotype relationships did not show an obvious geographical pattern (Figure 2): most individuals were represented by unique haplotypes with no obvious clustering of related haplotypes into Namibian or South African groups (Figure 2). The majority of haplotypes were closely related, differing from one another by one to two mutation steps, with the exception of two HEN individuals that were divergent by 10 mutation steps (Figure 2).

None of the six microsatellite loci exhibited evidence of amplification errors, and all displayed genotype frequencies that confirmed with Hardy-Weinberg and linkage equilibrium expectations (Table 2). Levels of genetic diversity in terms of heterozygosity and allelic richness were high (overall values of $H_E = 0.774$ and $AR = 13.7$), with both samples displaying very similar values at individual loci and overall

(Table 2). Number of private alleles varied between 1 and 7, per locus and region (Table 2). Analyses of statistical power of the dataset revealed that the loci and sample sizes used in this study could statistically detect genetic differentiation as low as $F_{ST} = 0.001$ in 99% of tests. As for the mtDNA data, nuclear genetic differentiation between the Namibian and South African samples was significantly greater than zero ($F_{ST} = 0.036$, $P < 0.05$), with Jost's D_{est} indicating a slightly higher level of differentiation ($D_{est} = 0.104$, $P < 0.05$). Estimates of contemporary migration rates per generation between the two geographical populations were low, independently of the method used or mutation rate considered (F_{ST} -based: $N_{em} = 6$; MIGRATE: $m_{2 \rightarrow 1} = 0.0014$; $m_{1 \rightarrow 2} = 0.0011$ for $\mu = 0.1\%$ per generation).

Discussion

Despite the preliminary nature of the present study, due to the limited number of sampling sites available, similarly high levels of genetic diversity and evidence for shallow but significant genetic differentiation between the two regional populations (Namibia and South Africa) of *Argyrosomus inodorus* was found. The observed mitochondrial and nuclear genetic diversity ($h = 0.991$, $\pi = 0.006$; $H_O = 0.771$, $H_E = 0.764$) were comparable with other commercially exploited fish species occurring in the Benguela Current region, such as *Argyrosomus japonicus* ($h = 0.96$, $\pi = 0.009$ – Kloppe 2005), *Lichia amia* ($h = 0.991$, $\pi = 0.006$ – Henriques et al. 2012), *Atractoscion aequidens* ($h = 0.853$, $\pi = 0.005$; $H_E = 0.889$ – Henriques et al. 2014) and *Rhabdosargus holubii* ($h = 0.91$, $\pi = 0.006$ – Oosthuizen 2007). High genetic diversity and shallow population structure are common features of marine teleosts, even in abundant, commercially exploited species. These are thought to result from historically high effective population sizes and/or high levels of gene flow between adjacent populations (Waples 1998). Interestingly, the observed genetic divergence between the Namibian and South African *A. inodorus* populations (mtDNA $\phi_{ST} = 0.092$; nuclear $F_{ST} = 0.036$ and $D_{est} = 0.104$, $P < 0.05$) was higher than that reported for other migratory sciaenids such as *Micropogonias undulatus* ($\phi_{ST} = 0.046$ – Lankford et al. 1999) and *Sciaenops ocellatus* ($\phi_{ST} = 0.057$ – Gold & Richardson 1998), but substantially lower than observed for other fish species with similarly disjunct distributions across the Benguela Current region (*L. amia*, $\phi_{ST} = 0.9$ – Henriques et al. 2012; *A. aequidens*, $\phi_{ST} =$

0.902, $F_{ST} = 0.055$ – Henriques et al. 2014). These results, combined with estimates of the number of contemporary migrants ($N_{em} = 0.0014 - 6$ per generation, depending on the method used), suggest a limited level of gene flow between Namibian and South African *A. inodorus* populations, and support the presence of incipient population differentiation. The present findings concur with tagging studies conducted for the species, where only two of 17,353 *A. inodorus* tagged in Namibia were recaptured in South Africa, suggesting that connectivity between populations may be limited (Kirchner & Holzhauzen 2001). Therefore, the low but significant genetic differentiation displayed by *A. inodorus* is likely to result from a present-day disjunct population distribution, with occasional migrants, and historically high levels of effective population size, rather than substantial gene flow between Namibia and South Africa.

As with other fish species distributed around southwestern Africa (e.g. *L. amia* – Henriques et al. 2012; *A. aequidens* – Henriques 2012; *Albula* spp. – Colborn et al. 2001), the distribution break in *A. inodorus* appears to correspond with the areas of cold water upwelling off southern Namibia and the west coast of South Africa (Griffiths & Heemstra, 1995; Griffiths 1997; Kirchner & Holtzhausen 2001). Although the limited sampling precludes the drawing of definitive conclusions, the reported genetic divergence and breakdown of gene flow across the Benguela Current suggests that the oceanographic features of the system, namely the cold water region, may be contributing to disrupt both adult and larval dispersal of *A. inodorus*, and supports the hypothesis of two isolated populations with limited migration between them. As the species is considered overexploited both in Namibia and South Africa, information on the distribution, population dynamics and long-term dispersal patterns across the Benguela Current region would provide a comprehensive evaluation of genetic structure, which should be incorporated into fishery management arrangements.

Acknowledgments

The present work and R. Henriques were supported by a Fundação da Ciência e Tecnologia – FCT grant (ref. SFRH/BD/36176/2007). The authors would like to thank Steve “Spyker” Krugger for assisting in sample collection in Namibia, and the

University of Namibia, and the Ministry of Fisheries in Namibia, for their invaluable support.

References

- Appleyard SA, Ward RD, Williams R. 2002. Population structure of the Patagonian toothfish around Heard, McDonald and Macquarie Islands. *Antarctic Science* 14:364-373.
- Archangi B, Chand V, Mather PB. 2009. Isolation and characterization of 15 polymorphic microsatellite DNA loci from *Argyrosomus japonicus* (mulloway), a new aquaculture species in Australia. *Molecular Ecology Resources* 9:412-414.
- Avice JC. 2000. *Phylogeography: The History and Formation of Species*. Cambridge: Harvard University Press. 447 pages.
- Bandelt HJ, Forster P, Rohl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16:37-48.
- Beerli P. 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use? In: Bertorelle G, Bruford MW, Hauffe HC, Rizzoli A, Vernesi C, editors. *Population Genetics for Animal Conservation*. Cambridge: Cambridge University Press, p 42-79.
- Chapuis MP, Estoup A. 2007. Microsatellite null alleles and estimation of population differentiation. *Molecular Biology and Evolution* 24:621-631.
- Colborn J, Crabtree RE, Shaklee JB, Pfeiler E, Bowen BW. 2001. The evolutionary enigma of bonefishes (*Albula* spp.): cryptic species and ancient separations in a globally distributed shorefish. *Evolution* 55:807-820.
- Crawford NG. 2010. SMOGD: software for the measurement of genetic diversity. *Molecular Ecology Resources* 10:556-557.
- DAFF. 2012. *Status of the South African marine fishery resources 2012*. Cape Town: DAFF. 71 pages.
- Excoffier L, Laval G, Schneider S. 2005. ARLEQUIN (version 3.0): An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics* 1:47-50.
- FAO. 2012. *The State of World Fisheries and Aquaculture*. Rome: FAO. 209 pages.

- Floeter SR, Rocha LA, Robertson DR, Joyeux JC, Smith-Vaniz WF, Edwards AJ et al. 2008. Atlantic reef fish biogeography and evolution. *Journal of Biogeography* 35:22-47.
- Gold JR, Richardson LR. 1998. Mitochondrial DNA diversification and population structure in fishes from the Gulf of Mexico and western Atlantic. *Journal of Heredity*, **89**, 404-414.
- Griffiths MH. 1997. The life history and stock separation of silver kob, *Argyrosomus inodorus*, in South African waters. *Fishery Bulletin* 95:47-67.
- Griffiths MH, Heemstra PC. 1995. A contribution to the taxonomy of the marine fish genus *Argyrosomus* (Perciformes: Sciaenidae), with description of two new species from southern Africa. *Ichthyological Bulletin of the J.L.B. Smith Institute of Ichthyology* 1-40.
- Heemstra P, Heemstra E. 2004. *Coastal Fishes of Southern Africa*. Grahamstown: NISC, SAIAB. 488 pages.
- Henriques R, Potts WM, Sauer WHH, Shaw PW. 2012. Evidence of deep genetic divergence between populations of an important recreational fishery species, *Lichia amia* L. 1758, around southern Africa. *African Journal of Marine Science* 34:585-591.
- Henriques R, Potts WM, Sauer WHH, Santos CV, Shaw PW. 2014. Population connectivity and phylogeography of a coastal fish, *Atractoscion aequidens* (Sciaenidae), across the Benguela Current region: evidence of an ancient vicariant event. *PLOS One* 9(2): e87907. doi:10.1371/journal.pone.0087907
- Holtzhausen JA, Kirchner CH, Voges SF. 2001. Observations on the linefish resources of Namibia, 1990-2000, with special reference to West Coast steenbras and silver kob. *South African Journal of Marine Science-Suid-Afrikaanse Tydskrif Vir Seewetenskap* 23:135-144.
- Hutchings L, Van der Lingen CD, Shannon LJ, Crawford RJM, Verheye HMS, Bartholomae CH, et al. 2009. The Benguela Current: An ecosystem of four components. *Progress in Oceanography* 83:15-32.
- Kirchner CH. 1998. Population dynamics of the exploited silver kob (*Argyrosomus inodorus*) in Namibian waters. PhD thesis, University of Port Elizabeth, South Africa. 276 pages.

- Kirchner CH, Beyer JE. 1999. Estimation of total catch of silver kob *Argyrosomus inodorus* by recreational shore-anglers in Namibia using a roving-roving creel survey. South African Journal of Marine Science-Suid-Afrikaanse Tydskrif Vir Seewetenskap 21:191-199.
- Kirchner CH, Holtzhausen JA. 2001. Seasonal movements of silver kob, *Argyrosomus inodorus*, (Griffiths and Heemstra) in Namibian waters. Fisheries Management and Ecology 8:239-251.
- Klopper AW. 2005. Intraspecific genetic variation in the percoid teleosts, *Argyrosomus japonicus* (Temminck & Schlegel, 1843) and *Pomadasys commersonnii* (Lacepede, 1801) as inferred from the mitochondrial control region. Master Thesis. University of Pretoria, South Africa. 144 pages.
- Lankford TE, Targett TE, Gaffney PM. 1999. Mitochondrial DNA analysis of population structure in the Atlantic croaker, *Micropogonias undulatus* (Perciformes : Sciaenidae). Fishery Bulletin 97:884-890.
- Oosthuizen CJ. 2007. Genetic variation within the Cape Stumpnose *Rhabdosargus holubii* Steindachner (Teleostei: Sparidae). Master Thesis. University of Pretoria, South Africa. 182 pages.
- Posada D. 2008. jModelTest: Phylogenetic model averaging. Molecular Biology and Evolution 25:1253-1256.
- Potts WM, Henriques R, Santos CV, Munnik K, Ansorge I, Dufois F, Booth AJ, Kirchner C, Sauer WHH, Shaw PW. 2014. Ocean warming, a rapid distributional shift and the hybridization of a coastal fish species. Global Change Biology, DOI: 10.1111/gcb.12612.
- Raymond M, Rousset F. 1995. GENEPOP version-1.2: Population genetics software for exact tests and ecumenicism. Journal of Heredity 86:248-249.
- Ryman N, Palm S. 2006. POWSIM: a computer program for assessing statistical power when testing for genetic differentiation. Molecular Ecology Notes 6:600-602.
- Sambrook J, Fritscher EF, Maniatis T. 1989. Molecular cloning: a laboratory manual. New York: Cold Spring Harbor Laboratory Press. 2028 pages.
- Shannon LV. 1985. The Benguela ecosystem. 1. Evolution of the Benguela, physical features and processes. Oceanography and Marine Biology 23:105-182.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence

alignment aided by quality analysis tools. Nucleic Acids Research 25:4876-4882.

Van der Bank H, Kirchner CH. 1997. Biochemical genetic markers to distinguish two sympatric and morphologically similar Namibian marine fish species, *Argyrosomus coronus* and *Argyrosomus inodorus* (Perciformes: Sciaenidae). Journal of African Zoology 111: 441-448.

Van Oosterhout C, Weetman D, Hutchinson WF. 2006. Estimation and adjustment of microsatellite null alleles in nonequilibrium populations. Molecular Ecology Notes 6:255-256.

Waples RS. 1998. Separating the wheat from the chaff: Patterns of genetic differentiation in high gene flow species. Journal of Heredity 89:438-450.

Tables

Table 1: Estimates of mitochondrial genetic diversity levels and neutrality tests for *A. inodorus* CR: *n* – number of individuals; *H* – number of haplotypes; *h* – haplotype diversity; π - nucleotide diversity; *F* – Ewens-Waterson neutrality test; *D* – Tajima neutrality test; *FS* – Fu neutrality test. Significant departures from expectations (*P* < 0.05) in bold.

	HEN	EastC	Overall
<i>n</i>	18	18	36
<i>H</i>	18	14	32
<i>h</i>	1.000	0.968	0.991
π	0.008	0.004	0.006
<i>F</i>	-	0.862	0.966
<i>D</i>	-1.486	0.324	-1.554
<i>FS</i>	-14.762	-10.099	-25.652

387 **Table 2:** Genetic diversity in *A. inodorus* samples at six cross-specific microsatellite
 388 loci: n – number of individuals genotyped; NA – number of alleles; AR – allelic
 389 richness; PA – number of private alleles H_E – expected heterozygosity; H_O – observed
 390 heterozygosity; F_{IS} – inbreeding coefficient. Significant deviations to Hardy-Weinberg
 391 expectations in bold.

		HEN	EastC	Overall
UBA5	n	40	40	80
	NA	11	11	13
	AR	10.803	10.925	10.52
	PA	1	2	3
	H_E	0.819	0.825	0.839
	H_O	0.875	0.825	0.850
	F_{IS}	-0.047	0.003	-0.007
UBA40	n	40	39	79
	NA	8	7	8
	AR	7.951	7.000	7.452
	PA	1	0	1
	H_E	0.765	0.807	0.790
	H_O	0.750	0.795	0.772
	F_{IS}	0.038	0.037	0.028
UBA50	N	39	40	79
	NA	14	15	16
	AR	13.974	14.899	14.846
	PA	1	2	3
	H_E	0.887	0.896	0.914
	H_O	0.821	0.800	0.810
	F_{IS}	0.066	0.038	0.120

UBA91	<i>n</i>	40	40	80
	<i>NA</i>	5	3	5
	<i>AR</i>	4.902	3.000	3.962
	<i>PA</i>	1	1	2
	<i>H_E</i>	0.361	0.387	0.375
	<i>H_O</i>	0.275	0.475	0.375
	<i>F_{IS}</i>	0.182	-0.207	0.006
UBA853	<i>n</i>	40	40	80
	<i>NA</i>	13	14	17
	<i>AR</i>	12.799	12.924	14.530
	<i>PA</i>	3	4	7
	<i>H_E</i>	0.831	0.872	0.876
	<i>H_O</i>	0.925	0.900	0.913
	<i>F_{IS}</i>	-0.100	-0.036	-0.035
UBA854	<i>n</i>	40	40	80
	<i>NA</i>	9	7	19
	<i>AR</i>	7.604	11.899	15.152
	<i>PA</i>	7	1	8
	<i>H_E</i>	0.881	0.776	0.860
	<i>H_O</i>	0.975	0.675	0.825
	<i>F_{IS}</i>	-0.038	0.130	0.047
Average all loci	<i>n</i>	40	40	80
	<i>NA</i>	10	9.500	11.333
	<i>AR</i>	9.004	9.833	13.667
	<i>PA</i>	14	10	24
	<i>H_E</i>	0.757	0.760	0.776
	<i>H_O</i>	0.770	0.745	0.758
	<i>F_{IS}</i>	0.000	0.012	0.027

Figure Legends

Figure 1: Sampling strategy for *A. inodorus* across the Benguela Current region, highlighting sampling sites, and their position relative to the major oceanographic features of the system: position of the Benguela and Agulhas Currents, central Namibia upwelling cell, and continental platform width.

Figure 2: Haplotype network for *A. inodorus* across the Benguela Current region, based on 704bp of mtDNA CR sequences: $\ominus$ = HEN; $\oplus$ = EastC. Branch lengths are proportional to number of nucleotide differences, and node sizes are proportional to the number of individuals. Red dots represent unsampled inferred haplotypes.