

Aberystwyth University

*Spatial genetic structure and dispersal of the cacao pathogen *Moniliophthora perniciosa* in the Brazilian Amazon*

Artero, A. S.; Silva, J. Q.; Albuquerque, P. S. B.; Bressan, E. A.; Leal Jr., G. A.; Sebben, A. M.; Griffith, G. W.; Figueira, A.

Published in:
Plant Pathology

DOI:
[10.1111/ppa.12644](https://doi.org/10.1111/ppa.12644)

Publication date:
2017

Citation for published version (APA):

Artero, A. S., Silva, J. Q., Albuquerque, P. S. B., Bressan, E. A., Leal Jr., G. A., Sebben, A. M., Griffith, G. W., & Figueira, A. (2017). Spatial genetic structure and dispersal of the cacao pathogen *Moniliophthora perniciosa* in the Brazilian Amazon. *Plant Pathology*, 358-371. <https://doi.org/10.1111/ppa.12644>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Received Date : 09-Nov-2015

Revised Date : 17-Oct-2016

Accepted Date : 18-Oct-2016

Article type : Original Article

Spatial genetic structure and dispersal of the cacao pathogen *Moniliophthora perniciosa* in the Brazilian Amazon

A.S. Artero^{1,&}; J. Q. Silva^{1,#a,&}; P.S.B. Albuquerque²; E.A. Bressan¹; G.A. Leal Jr.^{1,#b}; A.M. Sebbenn³,
G.W. Griffith⁴; A. Figueira^{1*}

¹Universidade de São Paulo, Centro de Energia Nuclear na Agricultura, Piracicaba, São Paulo, Brazil;

²Comissão Executiva do Plano da Lavoura Cacaueira, ERJOH, Marituba, Pará, Brazil;

³Instituto Florestal de São Paulo, Estação Experimental de Tupi, Piracicaba, São Paulo, Brazil;

⁴Aberystwyth University, Cledwyn Building, Institute of Biological, Environmental and Rural Sciences, Penglais, Ceredigion, SYD3 3DD, Wales, United Kingdom;

^{#a}Current address: Universidade Federal do Recôncavo da Bahia, Centro de Ciências Agrárias, Ambientais e Biológicas, Campus de Cruz das Almas, Cruz das Almas, Bahia, Brazil;

^{#b}Current address: Universidade Federal de Alagoas, Centro de Ciências Agrárias, Rio Largo, Alagoas, Brazil.

[&]These authors contributed equally to this work.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ppa.12644

This article is protected by copyright. All rights reserved.

*Corresponding author

e-mail: figueira@cena.usp.br

Running head: *Moniliophthora perniciosa* diversity

Keywords: *Crinipellis*; *Marasmius*; *Theobroma*; witches' broom disease.

Abstract

Moniliophthora perniciosa is the causal agent of witches' broom in *Theobroma cacao* (cacao). Three biotypes of *M. perniciosa* are recognized, differing in host specificity, with two causing symptoms on cacao or Solanaceae species (C- and S-biotypes), and the third found growing endophytically on lianas (L-biotype). Our objectives were to clarify the genetic relationship between the three biotypes, and to identify for the C-biotype, those regions in the Brazilian Amazon with the greatest genetic diversity. Phylogenetic reconstruction based on the rRNA ITS regions showed that the C- and S-biotypes formed a well-supported clade separated from the L-biotype. Analysis of 131 isolates, genotyped at 11 microsatellite loci found that S- and especially L-biotypes showed a higher genetic diversity. A significant spatial genetic structure was detected for the C-biotype populations in Amazonia for up to 137 km, suggesting isolation by distance mode of dispersal. However, in regions containing extensive cacao plantings, C-biotype populations were essentially 'clonal', as evidenced by high frequency of repeated multilocus genotypes. Among the Amazonian C-biotype populations, Acre and West Amazon displayed the largest genotypic diversity and might be part of the center of diversity of the fungus. The pathogen dispersal may have followed the direction of river flow downstream from Acre, Rondonia and West Amazon eastward to the rest of the Amazon valley, where cacao is not endemic. The Bahia population exhibited the lowest genotypic diversity, but high allele richness, suggesting multiple invasions, with origin assigned to Rondonia and West Amazon, possibly through isolates from the Lower Amazon population.

Introduction

Little attention has been given to the evolutionary biology and population genetics of plant pathogens (Giraud *et al.*, 2008), but establishing basic processes in the life history of parasites may assist in developing strategies to avoid or reduce their impact in agriculture and manage potentially harmful invasions. The dispersal ability of pathogens, especially fungi, defines the genetic differentiation between populations and the potential gene flow between populations at global, intracontinental or regional scale. Fungal mating system plays an important role in dispersal ability since reproductive strategies, such as selfing or clonality, may be beneficial for pathogens as a rapid means to propagate the fittest individuals, without requiring the presence of a compatible mating partner.

Moniliophthora perniciosa (Marasmiaceae *sensu stricto*; Aime & Phillips-Mora, 2005) is the causal agent of the witches' broom disease of *Theobroma cacao* L (cacao). Its basidiospores infect actively growing meristematic tissues (shoot apices, single flowers, flower cushions, and developing pods), inducing a range of symptoms depending on organ infected and stage of development (Purdy & Schmidt, 1996). Hypertrophic and hyperplastic growth of infected buds, called 'brooms', are the most remarkable symptoms, but economic losses derive mainly from infected pods and debilitated trees. It was originally believed that *M. perniciosa* infects only cacao and *Theobroma* congeners, and species from the sister genus *Herrania* (Purdy & Schmidt, 1996), all from the Malvaceae *sensu lato* (formerly Sterculiaceae). However, since the 1980s, it has been found infecting living tissues of plants belonging to diverse unrelated hosts, causing distinctive witches' broom symptoms in many of these.

Griffith & Hedger (1994a) proposed that *M. perniciosa* contains at least three discrete biotypes (C-, S-, and L-biotypes). The fungus exhibits host specificity, with only very limited development of symptoms (tissue swelling) in cross-inoculations among distinct biotypes and hosts. Strains that infect cacao and relatives are termed C-biotype (Griffith & Hedger, 1994a), whilst the S-biotype naturally infects various weedy solanaceous species (e.g. *Solanum rugosum*). However, upon artificial inoculation, it can cause broom symptoms on cultivated species, such as tomato and pepper (Bastos & Evans, 1985). The L-biotype has not been shown to cause witches' broom symptoms and has typically found forming basidiocarps in living woody liana vines and associated debris (Griffith & Hedger, 1994b). In coastal and Amazonian Ecuador, the L-biotype was consistently found on vines of *Arrabidaea verrucosa* (Bignoniaceae) (Griffith & Hedger, 1994b).

More detailed investigation of the breeding biology of the three biotypes (Griffith & Hedger, 1994b) showed that the L-biotype is with a multiallelic tetrapolar outcrossing mechanism, as is widely found among agaric fungi, such that formation of the fertile dikaryotic mycelium bearing clamp connections, first requires mating between two compatible monokaryons. In contrast, the C- and S- biotypes exhibit an unusual non-outcrossing breeding strategy found only rarely among agaric fungi whereby haploid, uninucleate basidiospores germinated on agar media are able to autodikaryotize over a period of 1-3 weeks following germination. However, in nature basidiospores germinate and infect host tissues forming a swollen and multinucleate 'biotrophic' mycelium within green brooms, with the typical clamped dikaryotic mycelium only appearing as the brooms die (Griffith & Hedger, 1994c). The breeding strategy of the C- and S- biotypes is often described as 'clonal' and we also use this term here. However, we recognize that meiotic division following karyogamy of two identical nuclei is not *clonality* in its strictest sense. Furthermore, in multiply infected brooms, it is possibly that fusion of hyphae derived from genetically different basidiospores could lead to the formation of hetero-dikaryons.

Defining the areas with the largest genetic diversity would help to devise strategies to search for resistance in the Amazon region, putatively considered the center of origin of both *M. perniciosa* (Baker & Holliday, 1957; Purdy & Schmidt, 1996) and its host *T. cacao* (Motamayor *et al.*, 2008; Thomas *et al.*, 2012). The C-biotype is endemic to the Amazon (Purdy & Schmidt, 1996), from where it invaded and devastated local industries in various cacao-producing regions across South America and the Caribbean, often being transported unwittingly. The first report of witches' broom affecting commercial cacao plantations occurred in Suriname in 1895 (Rorer, 1913). Afterwards, the pathogen was sequentially reported occurring in British Guyana (1906), Ecuador (1921), Trinidad (1928), Tobago (1939), and Grenada (1948) (Baker & Holliday, 1957), reaching the largest Brazilian cacao producing region in southern Bahia state in 1989 (Andebrhan *et al.*, 1999). The long distance between Bahia and the Amazon region (over 2,000 km; Figure 1) and some phytosanitary controls likely avoided an earlier invasion by the pathogen. The fungus is presently found in all South American and Caribbean cacao producing countries, with estimated losses from witches' broom disease reaching over 250,000 tonnes.year⁻¹ (Bowers *et al.*, 2001). There are current serious concerns about the potential risk of *M. perniciosa* invasion in West Africa, a region responsible for over 70% of the world cacao production, with potential catastrophic socioeconomic consequences.

The natural distribution of *T. cacao* in Brazil is limited to the Upper Amazon region (Serenio *et al.*, 2006; Motamayor *et al.*, 2008; Thomas *et al.*, 2012), but when cacao became a valuable commercial product during the 17th century, cultivation extended eastward to near the mouth of the

Amazon River, particularly stimulated by Jesuit missionaries (Bartley, 2005). Cacao production was the original basis of the Amazon economy until being supplanted by rubber in the late 19th century. Later, cacao cultivation was encouraged by the Brazilian government in the state of Rondonia and in Central Para state along the Transamazon road (Figure 1), as part of colonization projects during the 1970s and 1980s (Godar *et al.*, 2012). The states of Rondonia and Acre, near the putative center of diversity, contained spontaneous wild cacao stands (possibly from ancient pre-European plantings), with natural incidence of *M. perniciosa* (Bartley, 2005; Thomas *et al.*, 2012). On the other hand, the eastern parts of the Transamazon road (Central Para state) did not contain spontaneous cacao stands (Bartley, 2005). Since the establishment of cacao cultivation in the Transamazon road region (Godar *et al.*, 2012), witches' broom became a major limiting factor for production. But the origin of *M. perniciosa* in the region along the Transamazon road is unclear, as it could have derived either from the old plantations along the Amazon River (~400 km away) or around Belem (~500 km), from local cupuassu (*T. grandiflorum*) stands, or from elsewhere.

The spread of the fungus to Southern Bahia occurred in 1989, apparently in two primary outbreaks (Andebrhan *et al.*, 1999), possibly by either deliberate or accidental intervention by man (Homewood, 1991). The source of the original genotypes remains unknown, but clearly at least two distinct lineages have been consistently detected by various molecular means (Andebrhan *et al.*, 1999; Rincones *et al.*, 2003; Santana *et al.*, 2012). Investigation of the origin of the genotypes in these major cacao-producing regions is useful to inform breeding programs about diversity, dispersal patterns, and structure of these pathogen populations.

Therefore, the objectives of this study were to describe the relationship between the biotypes based on ribosomal gene internal transcribed spacer sequences (ITS), while defining the regions in the Brazilian Amazon with the largest genetic diversity of the pathogenic C-biotype using microsatellite loci, to determine the pathogen genetic population structure, the dispersal pattern in the Amazon and to verify the origin of introduced genotypes in cacao producing regions (Bahia and Transamazon).

Materials and methods

Moniliophthora perniciosa sampling: A systematic collection of isolates from the C-biotype was conducted from infected *T. cacao* trees in 2004 in areas of the Amazonian region where the fungus occurs naturally, including sites in the states of Acre, Amazonas, Pará, and Rondonia in Brazil, and in Ecuador (Figure 1). Infected cacao shoots ('brooms') were collected from trees under cultivated or

natural conditions along main roads at the Brazilian Amazonian, located at a minimum distance of 10 km between sites, with the geographic positioning recorded (Table 1, Table S1). The isolates from Acre, Amazonas, Pará and Rondonia were collected by the staff of the Brazilian Ministry of Agriculture of the “Comissão Executiva do Plano da Lavoura Cacaueira” (CEPLAC). Briefly, infected cacao shoots (‘green brooms’, 20-50 cm long) were collected from trees and taken back to the laboratory at the “Estação de Recursos Genéticos José Haroldo” (ERJOH), Marituba, Para state, where mycelia were cultivated using a selective PDA medium (Potato-Dextrose-Agar containing 100 mg L⁻¹ benzimidazole and 100 mg L⁻¹ streptomycin) from small slices of shoot tissues after surface sterilization with 5% sodium hypochloride solution. All isolates were routinely maintained on PDA media.

C-biotype isolates from Southern Bahia were included to investigate their origins. These isolates from Bahia were kindly provided by Prof. Gonçalo Pereira from the “Universidade Estadual de Campinas - UNICAMP” (isolates 74, 75, 76, and 77), and by Dr. Karina Peres Gramacho from CEPLAC, Itabuna, Bahia state (the remaining fourteen isolates). A total of 111 C-biotype isolates of *M. pernicioso* were analyzed (Table 1, Table S1). In addition, 14 isolates from the S-biotype and six isolates from the L-biotypes, originally collected in Brazil and Ecuador (Table 1) were sourced from the Mycology culture collection at Aberystwyth University or by Dr. Alan Pomella from the Mars Center for Cocoa Studies, Barro Preto, Bahia, Brazil. Isolates were transported under the permit of the Brazilian Ministry of Agriculture (‘Parecer Técnico’ DFCP/CGPP 003/2010).

DNA extraction: Isolates were cultivated on PDA in Petri dishes for two weeks, when hyphae plugs from the growing edge of the culture were used to inoculated 50 mL of medium liquid MYE + G (0.17% malt extract, 0.5% yeast extract, 5% glycerol). These cultures were incubated in the dark under agitation (100 rpm) at room temperature for two weeks. Mycelium was then collected by filtration and stored at -80° C. DNA was extracted from mycelia following a CTAB protocol (Silva *et al.*, 2008). DNA concentration was determined by fluorimetry using DyNA Quant 2000 (Amersham Pharmacia Biotech; Buckinghamshire, UK).

ITS sequence analysis and phylogenetic reconstruction: For genetic analysis, the internal transcribed spacer region (spanning ITS1 and ITS2) was amplified, using the primer pairs ITS1F and ITS4 (White *et al.*, 1990), and sequenced using Sanger BigDye technology. Sequence management was conducted within the Geneious (v.6.1.6) bioinformatics package using MAFFT for sequence alignment (default settings). Phylogenetic reconstruction was conducted using PhyML and the GTR substitution model.

Microsatellite loci analyses: The genetic diversity of the 131 isolates of *M. pernicioso* of the C-, S- and L-biotype (Table 1, Table S1) was analyzed using specific primers for 14 microsatellite loci, including nine loci (*mMpCena3*, *mMpCena4*, *mMpCena8*, *mMpCena11*, *mMpCena12*, *mMpCena16*, *mMpCena19*, *mMpCena22*, and *mMpCena26*) described by Silva *et al.* (2008), and additional five loci (*MsCepec_14*, *MsCepec_15*, *MsCepec_16*, *MsCepec_19*, and *MsCepec_45*) developed by Gramacho *et al.* (2007). The amplification reactions were conducted as previously described (Silva *et al.*, 2008). Amplification products were separated in denaturing sequencing gels (7% polyacrylamide, 7 M urea) ran in Tris-borate buffer at 50 W for 2 h, and visualized by silver staining (Silva *et al.*, 2008). Allele calls were conducted by migration distance in relation to a DNA 100 bp molecular ladder (Fermentas, Waltham, MA, USA).

Genetic diversity analyses: To identify the presence of repeated multilocus genotypes among isolates, we used the genetic identity analysis implemented in Cervus 3.0 (Kalinowski *et al.*, 2007). The likelihood that two individuals taken at random have different multilocus genotypes (D_g) in each population was estimated by: $D_g = [n/(n-1)](1 - \sum p_i^2)$, where n is the sample size and p_i is the frequency of the i^{th} multilocus genotype in each population (Kalinowski *et al.*, 2007). Estimates of D_g were obtained using the Multilocus program (Agapow & Burt, 2001). Biotype and population genetic diversities were determined using conventional parameters: the average number of alleles per locus (A), allele richness estimated for a minimum sample size of 9 or 6 individuals (R) estimated based on the rarefaction method (El Mousadik & Petit, 1996), expected heterozygosity (H_e) and fixation index (F). The statistical significance of the F values was estimated based on Monte Carlo permutation of alleles among individuals and a Bonferroni correction of multiple tests ($\alpha= 0.05$). All these parameters and analyses were estimated using the FSTAT program, version 2.9.3.2 (Goudet, 1995).

Spatial genetic structure analyses and isolation by distance: The isolation by distance (IBD) was investigated using only unique multilocus genotypes from the Amazonian region ($n= 56$) based on the analysis of spatial genetic structure (SGS). Spatial genetic structure was determined using the estimation of the average co-ancestry coefficient (θ_{xy}) between pairs of isolates and SPAGeDi version 1.3 (Hardy & Vekemans, 2002). The estimates of θ_{xy} based on Loiselle *et al.* (1995) have the advantage to not assume Hardy-Weinberg equilibrium and it can be used in populations with inbreeding. To visualise the SGS, θ_{xy} values were averaged over a set of ten distance classes and plotted against distances. The distance classes were chosen to contain the same number of pair of individuals. To test whether there was a significant deviation from random spatial genetic structure;

the 95% confidence interval was calculated for each observed value and each distance class from 1,000 permutations of individuals among locations.

Analysis of genotype assignment: Individual assignment tests were carried out using the Bayesian multilocus approach (Cornuet *et al.*, 1999) implemented in the GeneClass2 program (Piry *et al.*, 2004). Based on Efron (1983), all individuals from the reference dataset were self-classified to the sampled populations using the leave-one-out approach (self-assignment). There was a most likely population in any reference set to which the individuals could always be assigned to. However, the set of reference populations might not include the true population of origin for the control group resulting in a false positive assignment. Therefore, a measure of confidence was needed that the tested individuals truly belonged to a given population (Halkett *et al.*, 2010). This was developed by comparing the likelihood value of the individual test with the likelihood distribution of the population based on allele frequencies of the population, generated from 10,000 sampling runs with replacement. If the observed likelihood of the individual test was outside the distribution, then this indicated that the individual did not belong to that population.

Results

Phylogenetic analysis of the three biotypes: In order to better define the relationship between the three genotypes, the ITS 'DNA barcode' region for all L-biotype and S-biotype and a subset of C-biotype samples was sequenced. Phylogenetic reconstruction based on alignments of sequences from the present study and several deposited in GenBank showed that the C- and S-biotype formed a well-supported clade (Figure 2) with very few polymorphisms variation found within the ITS1/2 spacer regions. However, within this clade, S- and C-biotype isolates were separated but with only poor statistical support. L-biotype isolates were clearly separated from C-/S-biotype samples, with isolates from coastal Ecuador (112-116) forming a distinct well-supported clade, clearly separated from the L-biotype sample from Amazonian Ecuador (117). Several unpublished sequences of the L-biotype of *M. pernicioso* from Brazil are deposited on GenBank (Tarnowski & Ploetz, 2008). The host liana is identified as either Bignoniaceae or Malphiaceae and isolates obtained from these distinct hosts clustered separately with good statistical support. Furthermore, the 'Malphiaceae' L-biotype clade also contained the sequence (AY317137) of the new species *Crinipellis* (= *Moniliophthora*) *brasiliensis* that was reported by Arruda *et al.* (2005) forming brooms on *Heteropterys acutifolia* in Eastern Brazil (Minas Gerais). For the purpose of the present study, these analyses confirm that L-biotype is distinct from the C-/S- biotypes consistent with its outcrossing breeding strategy (Griffith

& Hedger, 1994a). However, further work is needed to determine whether there is host specialization of the L-biotype on Bignoniaceae and Malpigiaceae and also whether the broom formation reported for *M. brasiliensis* by Arruda *et al.* (2005) occurs on hosts in Bignoniaceae.

As all of the C-biotype isolates were obtained from infected tissues, it was necessary to confirm their identity (Table 1, Table S1). The ITS1-5.8S-ITS2 region was amplified (*circa* 750 bp fragment) and digested by either *MspI* or *HinfI* (Arruda *et al.*, 2003). All putative *M. pernicioso* isolates gave an identical PCR-RFLP pattern thus confirming their identity (Figure S1).

Analyses of microsatellite loci from isolates of the three biotypes: Analysis of the 14 microsatellite loci in all 131 isolates revealed size polymorphism at all but one locus, and of these, unambiguous allelic polymorphism was present at 11 loci. A total of 56 alleles were found, with a mean of 5.1 alleles per locus, ranging from the *mMpCena19* locus with a total of 9 alleles, to *MsCepec_15* with only two alleles.

Most isolates were homozygous for nearly all the loci, but heterozygosity was detected for eight loci (*mMpCena3*, *mMpCena4*, *mMpCena8*, *mMpCena11*, *mMpCena19*, *mMpCena22*, *MsCepec_16*, and *MsCepec_19*) for a few isolates. Heterozygous isolates from all three biotypes were observed for *mMpCena3*. For *mMpCena11*, heterozygosity was revealed for isolates from L- and S-biotypes. For another four loci (*mMpCena4*, *mMpCena19*, *MsCepec_16* and *MsCepec_19*), heterozygous individuals were disclosed only from L-biotype. Isolate 113 (Pichilingue, Ecuador) of the L-biotype was the only one exhibiting two alleles for loci *mMpCena4*, *mMpCena19* and *MsCepec_19*. For *MsCepec_16*, only the L-biotype isolate 114 (Pichilingue, Ecuador) was heterozygous. C-biotype isolates from Acre (65 and 66 Mal. Thaumaturgo, 69 and 73 Assis Brasil), Pará (27 Trairão) and Amazonas state (43 and 44 Costa da Conceição, 45 and 46 Ceplac/ERNEG) were heterozygous for *mMpCena8*, whereas isolates from Pará state (1 Anapu, 7, 8 and 9 Medicilândia, 14 Uruará, 15 Alenquer, 17 Santarém, 19, 20 and 21 Cameté, 22 Baião, 23 and 24 Mocajuba) were heterozygous for *mMpCena22* (Table 1, Table S1). For *mMpCena3*, heterozygous C-biotype isolates were identified from Pará (26 Trairão), Amazonas (43 Costa da Conceição, 45 Ceplac/ERNEG), Rondonia (56, 57 and 58 Ouro Preto D'Oeste), and Acre (65 and 66 Mal. Thaumaturgo, 70 Brasileira).

Isolates from the L-biotype displayed a total of 42 alleles, with an average of 3.81 alleles per locus, with eight private alleles (Table S2). For the S-biotype isolates, 41 alleles were identified, with an average of 3.72 alleles per locus and six private alleles. For all 111 C-biotype isolates, a total of 29 alleles were disclosed, with average 2.63 alleles per locus and with only three private alleles.

Identification of repeated multilocus genotypes within populations: The microsatellite analyses indicated that some isolates from the same region of collection shared the same allelic composition for the investigated loci. Therefore, Cervus 3.0 was used to verify and identify isolates with identical multilocus genotypes ('clonal', with exact allelic match). Our results indicated that from the 131 isolates, only 83 corresponded to unique multilocus genotypes, with the remaining 48 shared identical allele composition with one or more isolates (Table 1, Table S3).

Spatial genetic structure (SGS) and isolation by distance of C-biotype isolates: Excluding the seven multilocus genotypes from Bahia (Table 1), where *M. perniciosus* was only recently introduced, and one isolate without proper spatial position, the analysis of the remaining 56 unique multilocus genotypes from the Amazon revealed a significant SGS for up to 137 km (Figure 3), tending to zero or significantly negative values for longer distances, in a typical pattern of isolation by distance.

The SGS analysis allowed us to arbitrarily define seven geographic populations with the isolates from the C-biotype of *M. perniciosus* (Figure 1): Central Pará (Transamazon road region), Pará or Lower Amazon (surrounding the Amazon river mouth), East Amazon (around Manaus), West Amazon (Upper Solimões at the Colombia-Peru border), Rondonia, Acre, and Bahia (Table 1). Among these seven populations, Central Pará, East Amazon, Rondonia, and Bahia were the ones with the largest number of repeated multilocus genotypes (>40%) (Table 1). From the 47 repeated multilocus genotypes identified for the C-biotype, only in three cases identical genotypes were identified in locations outside the original population (Table S3). Isolates from the East Amazon (44 Costa da Conceição and 46 Ceplac-ERNEG, AM) displayed an identical multilocus match with isolate 27 from Trairão, PA (Central Pará) and 73 from Assis Brasil (Acre). Isolate 64 from Marechal Thaumaturgo (Acre) was identical to isolates from Central Pará (6 Medicilândia and 25 Trairão), whereas isolates 107 and 109 from Alenquer (Lower Amazon) were identical to isolate 92 from Tabatinga, AM (West Amazon) (Table S3). For the S-biotype, only one pair of repeated multilocus genotypes was identified (isolates 120 and 121, both from Manaus), and no repeated multilocus genotypes was detected among the six L-biotype isolates (Table 1, Table S3).

Genetic diversity within and between biotypes and populations: The seven arbitrary C-biotype populations of multilocus genotypes contained a distinct number of members, ranging from 10 (Acre) to 20 (Central Pará), while there were only six L-biotype isolates (Table 2). Isolates from the L- and S-biotype displayed significantly higher expected heterozygosity (H_e) compared to the C-biotype isolates (0.69/0.56 vs. 0.29; Table 2). Within the C-biotype, the highest H_e was observed for Pará population (0.39), followed by Bahia (0.32) and Acre (0.31). On the other hand, Acre held the largest genotypic diversity ($D_g = 0.96$), followed by the West Amazon ($D_g = 0.95$), while Bahia

displayed the lowest D_g (0.77). All C-biotype isolates exhibited a high and significant fixation index (Table 2). Allele richness (R) was standardized for a sample size (n) of nine genotypes (Table 2) when considering all genotypes (including identical multilocus genotypes). For the C-biotype, allele richness (R) ranged from 2.14 (Bahia) to 1.71 (West Amazonas), whereas the S- and L-biotype displayed higher values (3.56 and 3.82, respectively). When only the unique multilocus genotypes were analyzed ($n=6$, Table 2), the genetic parameters showed a similar trend for most cases, except that the highest H_e was observed for the Bahia population (0.45), followed by Pará (0.40) and Rondonia (0.37).

Genetic assignment of genotypes to populations: We undertook a genetic assignment test to assign or exclude the possible origin of individual multilocus genotypes to reference populations. When the 13 S-biotype genotypes were analyzed against all the remaining genotypes, nine were assigned at a high score (>99%, Table 3) to isolate 117 of the L-biotype from San Carlos in Amazonian Ecuador, whereas isolate 123 and 129 displayed a lower score (<92%) to the same genotype. Of the two other S-biotype genotypes, the 124 displayed a high assignment score (>99%) to genotype 63 (Ji-Paraná, RO), and 119 showed a lower score (50.1%) to genotype 91 (Aurelino Leal, BA), both from the C-biotype (Table 3). As a group, the S-biotype isolates were assigned at a 100% probability to the L-biotype isolate 117 from San Carlos, Ecuador.

When all 64 unique multilocus genotypes from the C-biotype were tested against those from the L- and S-biotypes (Table S4), 57 were assigned at a high score (>98%) to L-biotype isolate 117 from San Carlos, while two (32 and 93) showed lower score (between 90.0 to 93.4%) to the same genotype (Table S4). The remaining five genotypes were assigned to S-biotype isolate 131 from Manaus (<97.0%). As a group (containing the seven populations), the C-biotype genotypes were assigned to the L-biotype isolate 117 from San Carlos, Ecuador at a 100% score.

Conversely, when all six genotypes of the L-biotype from Ecuador were analyzed for assignment against all the remaining genotypes from the S- and C-biotypes (Table 3), three were assigned with a high score (> 99%) to the S-biotype isolate 131 (Manaus, AM, Brazil), together with the isolate 116 (Janauche), but at a lower score (51.6%). The two other L-biotype genotypes (115 and 117) were assigned at a high score (>92%) to C-biotype genotypes (63 Ji-Paraná, RO or 91 Aurelino Leal, BA, respectively, Table 3). The group of L-biotype isolates matched, with 100% score, the S-biotype isolate 131 from Manaus.

To determine the putative origin, genetic assignment analyses were then conducted for each genotype from each arbitrary individual population of the C-biotype where *M. pernicioso* was originally neither endemic nor its host *T. cacao* was endemic (East Amazon, Central Pará, Lower Amazon, and Bahia). The 10 genotypes from East Amazon were assigned with a score below 96% to genotypes from 92 Tabatinga, AM, 63 Ji-Paraná, RO, and 73 from Assis Brasil, AC (Table 3). As a group, the East Amazon C-biotype isolates were assigned at 94.2% probability to isolate 73 from Assis Brasil, Acre. The 12 genotypes from Central Pará were analyzed in the same way, and six were assigned to genotype 73 from Assis Brasil, Acre, two to genotype 63 Ji-Paraná, Rondonia, and four to genotype 45 Ceplac/ERNEG from East Amazon, which was assigned to genotype 73 Assis Brasil, Acre (Table 3). The Central Pará population showed a high score (99.1%) to the genotype 45 Ceplac/ERNEG.

For the nine genotypes from the Lower Amazon population, three were assigned to genotype 73 from Assis Brasil, Acre (score ranging from 90.7 to 97.7%), whereas four had a 100% score with isolate 110 from Agua Blanca, Ecuador (Table 3). The two other genotypes were assigned at 81.2 and 91.6% probability to isolate 63 from Ji-Paraná, RO and 45 Ceplac/ERNEG, AM, respectively. As group, the Lower Amazon population shared a 100% certainty to derive from genotype 63 from Ji-Paraná, Rondonia.

Among the seven genotypes from Bahia, four genotypes (74, 77, 84 and 91) were assigned to an origin from Tabatinga, AM (92) with scores ranging from 53.4 to 100.0%, while another three (81, 87 and 90) were assigned to genotype 63 from Ji-Paraná, Rondonia, with scores ranging from 71.0 to 98.2%. As a group, the isolates from Bahia were assigned to genotype 92 from Tabatinga, AM at 100% probability.

The same analyses were conducted for the populations from Rondonia, Acre and West Amazonas (Table S5). Of the seven genotypes from Rondonia, two exhibited high probability of assignment (> 99%) to genotype 45 Ceplac/ERNEG, and one to 73 Assis Brasil, Acre. Of the eight genotypes from Acre, four shared scores between 80.7 to 98.3% to genotype 73 Assis Brasil, three shared scores between 92.6 to 97.8% to genotype 63 Ji-Paraná, RO, and one to genotype 45 Ceplac/ERNEG (score 76.1%). All the 11 multilocus genotypes from West Amazon (Upper Solimões) at the border between Brazil, Colombia and Peru, displayed a high score (>99%) with an identical multilocus genotype (isolate 109) from the same population (isolate 92 Tabatinga) (Table S5).

Discussion

The three recognized *M. pernicioso* biotypes exhibit contrasting life styles, ranging from the highly pathogenic C-biotype on cacao, the occasionally pathogenic S-biotype, to the apparently endophytic non-pathogenic L-biotype (Griffith & Hedger, 1994a). Our phylogenetic analyses found the pathogenic broom-forming biotypes (C- and S-) to be closely related and exhibiting little genetic variability, consistent with their primary homothallic (homomictic) breeding strategy, whereas the L-biotype (both Pacific and Amazonian populations) was distinct and more variable due to its bifactorial outcrossing mechanism (Griffith & Hedger, 1994b). Biparental inbreeding might occur in the S- and C-biotypes, possibly by selection for homothallism, linked to a more competitive dispersal ability (i.e. meristem infection by a single spore could lead to later basidiocarp formation without any mating event).

L-biotype isolates exhibited the greatest level of heterozygosity, possessing more alleles ($A=42$), including eight private alleles, the largest allele richness (R), expected heterozygosity (H_e) and genotypic diversity (D_g). Somatic compatibility and molecular analyses (Griffith & Hedger, 1994a; Arruda *et al.*, 2003) have previously suggested that the broom-forming C- and S-biotypes were more similar to each other, while exhibiting a lower level of diversity, in comparison to the L-biotype. Evidence from ITS nucleotide sequences (Arruda *et al.*, 2005), karyotype and microsatellite telomeric amplification analyses suggested that the S-biotype is more diverse than the C-biotype (Rincones *et al.*, 2006). Indeed, our results indicated a lower genetic diversity for the isolates from the C-biotype in comparison to the S- or L-biotypes. It is recognized that a pathogen transition to an agricultural host, even in the cases of sexually reproducing pathogen, imposes a founder effect with important reduction in genetic variation (Stukenbrock & Bataillon, 2010). Such a genetic bottleneck could have arisen in *M. pernicioso* by a switch from outcrossing to non-outcrossing breeding strategy, as suggested by Griffith and Hedger (1994a).

The convergence of most S- and C-biotype multilocus genotypes to the same L-biotype isolate (117) could indicate a shared ancestry of both biotypes. Most of the C-biotype unique multilocus genotypes (92%, 59 out of 64) were assigned to L-biotype isolate 117 from San Carlos, Ecuador, while the remaining five isolates shared higher score against S-biotype isolate 131 from Manaus, AM. These results suggested that both pathogenic biotypes (C- and S-) might have evolved from a more ancestral non-pathogenic form, likely the L-biotype. S-biotype and C-biotype might have evolved by disruptive selection by infection on distinct host species occurring in disturbed forest areas, either as weedy invader (*eg.* Solanaceae species) or as a susceptible host with induced meristematic flushing by the disturbed condition (*Theobroma* and *Herrania*).

The dispersal of *M. pernicioso* depends on the release of the haploid uninucleate basidiospores, the only recognized infective structure of the C-biotype, following meiosis from basidiocarps (Purdy & Schimdt, 1996). Basidiospores are highly sensitive to desiccation, losing viability within hours, but long distance dispersal, most likely during the night, has been reported to up to 50-70 Km in Ecuador (Wheeler & Soarez, 1993). Tropical rainfall regimes do not typically limit infection cycles over the year, but infection tends to be monocyclic within a year (Purdy & Schimdt, 1996), and continuous inoculum release may favor spread over long distances upon the appropriate conditions. However, movement of infected plant material under unwitting human intervention might predominate as a mechanism for longer distance dispersal.

The dispersal ability of pathogens, especially fungi, defines the genetic differentiation among populations and the potential gene flow between distant populations. Here, we detected a significant spatial genetic structure (SGS) for isolates of the C-biotype in the Brazilian Amazon, with significant co-ancestry detected for up to 137 km, suggesting the occurrence of 'isolation by distance' (IBD) mode of dispersal. A large number of repeated multilocus genotypes (47 in 111 isolates) were detected among C-biotype isolates from the same region, especially in regions containing large commercial stands of cacao, such as Central Pará, East Amazonas, Rondonia and Bahia. C-biotype isolates were mostly homozygous. As the C-biotype mycelia is homomictic, dispersal can be achieved without the requirement for mating prior to sexual reproduction, leading to the spread of geographically extensive adapted lineages (Griffith & Hedger, 1994b), similar to 'clones' (Stukenbrock & Bataillon, 2010).

The C-biotype of *M. pernicioso* is endemic to the Upper Amazon valley, where it infects cacao and related species (Purdy & Schmidt, 1996). The travel report of the naturalist Alexandre Rodrigues Ferreira, who visited this region in 1785-1787 described cacao plants exhibiting what appeared to be witches' broom symptoms ('*lagarto*' [= lizard]; Silva, 1987), representing the earliest report of the disease. Subsequently, the introduction of *M. pernicioso* into commercial cacao plantations was documented since the first outbreak in 1895 in Suriname (Rorer, 1913). Within this 100-year period, modern cultivation of cacao had expanded in the Amazon and the pathogen might have spread throughout the area, beyond its original natural distribution.

The Acre and West Amazon populations appeared to contain the greatest genetic diversity within the C-biotype, and both might be part of the center of diversity of the fungus in Brazil. Of these, the Acre population was the more diverse and was the only area where heterozygosity (at two loci) was detected within the C-biotype populations. Such heterozygosity could have arisen following multiple infections of meristems by genetically different spores. The West Amazon population also presented an elevated genotypic diversity, but the lowest allele richness, and together with the Acre population, both contained the least repeated multilocus genotypes. Most of the multilocus genotypes from Acre tended to be assigned to genotypes from Acre, or to a lesser extent, to Rondonia (isolate 63), corroborating that Acre might be part of the origin of many multilocus genotypes from the Amazon.

The genetic assignment test indicated that the putative origin of isolates from the East Amazon, Lower Amazon and Bahia populations was from the West Amazon. Some genotypes from Rondonia also pointed to a West Amazon origin. The Acre and West Amazon populations corresponded geographically to the regions that presumably encompass the center of diversity of *T. cacao* (Sereno *et al.*, 2006; Motamayor *et al.*, 2008; Thomas *et al.*, 2012), therefore these two populations appeared to contain the largest genetic diversity of *M. perniciosa*, and might be part of the center of genetic diversity and dispersal of the fungus.

The Rondonia population appeared to have an important role in the dispersal of isolates, as individual multilocus genotypes from the East Amazon, Central Pará, Lower Amazon, and Bahia traced back to isolate 63 from Ji-Paraná. But Rondonia showed no private alleles and a limited number of heterozygous isolates for *mMpCena3*. According to GeneClass, the genotypes from Rondonia derived from Acre (73 Assis Brasil) or West Amazon (92 Tabatinga). Rondonia became an important cacao producing region since the 1970s, with large areas of cultivation, from where the isolates were collected, which might have limited the genetic diversity of the pathogen.

The spread of the pathogen elsewhere occurred probably in association with cacao dispersal, since the host species did not occur naturally outside the Upper Amazon region. The movement of the pathogen might have followed the direction of river flow downstream from Acre, Rondonia and West Amazon eastwards to the rest of the Amazon valley to regions where cacao is not native, and where modern cultivation started around 300 years ago (Bartley, 2005). The Central Pará population appeared to have derived exclusively from introductions of genotypes from Acre and Rondonia, but not West Amazon, suggesting that the pathogen was either introduced directly from Rondonia and Acre, or from the Santarém region, a traditional region of cacao cultivation, where the West Amazon

genotypes were not primarily detected. Introduction from the Lower Amazon was less likely as West Amazon origins were also detected in that population.

The Bahia population exhibited the lowest genotypic diversity, but high allele richness, suggesting multiple introductions. The multilocus genotypes from Bahia, a recent and well-documented invasion of *M. pernicioso* (Andebrhan *et al.*, 1999), displayed the clear migration from two origins, with ancestry to genotype 63 from Ji-Paraná (Rondonia) and 92 from Tabatinga (West Amazon). Cacao was introduced in Southern Bahia in the 18th century, and the region managed to be free of *M. pernicioso* for over 200 years. In 1989, two independent foci of *M. pernicioso* occurrence were reported at Uruçuca and Camacan municipalities, 120 km apart (Andebrhan *et al.*, 1999). Our results clearly indicated two major origins for the genotypes introduced in Southern Bahia. The presence of two distinct lineages of *M. pernicioso* isolates in Bahia was originally demonstrated by Andebrhan *et al.* (1999) and later confirmed by many other genetic analyses (Rincones *et al.*, 2003; Ploetz *et al.*, 2005; Santana *et al.*, 2012). Thus, it is possible that this invasion occurred based on isolates from the Lower Amazon population (Pará), an area with an easier and more intense contact with Southern Bahia.

The microsatellite analysis deployed in the present study has demonstrated only limited genetic variability within C-biotype populations of *M. pernicioso*, consistent with non-outcrossing (clonal) spread. The presence of genetically very similar strains in geographically distant areas is consistent with known and suspected incidences of translocation of the pathogen by man. However, the origins of the C-biotype remain uncertain but it may have evolved from an ancestral population infecting solanaceous hosts. However, since these S-biotype populations are currently under-sampled, additional collections from more diverse locations could shed light on the origins of this destructive pathogen.

Acknowledgments

The authors wish to thank the “São Paulo Research Foundation” (FAPESP) for financial support (03/11483-0, 07/07175-0) and CEPLAC for technical support. ASA, PSBA, GAL, AMS, and AF were recipients of Brazilian National Research Council (CNPq) fellowships. Technical assistance by Edivaldo Pimentel is greatly appreciated. GWG acknowledges funding from Cocoa Research UK Ltd.

References

- Agapow PM, Burt A, 2001. Indices of multilocus linkage disequilibrium. *Molecular Ecology Notes* **1**, 101-2.
- Aime MC, Phillips-Mora W, 2005. The causal agent of witches' broom and frosty pod rot of cocoa (chocolate, *Theobroma cacao*) form a new lineage of Marasmiaceae. *Mycologia* **97**, 1012-22.
- Andebrhan T, Figueira A, Yamada MM, Furtek DB, 1999. Molecular fingerprinting suggests two primary outbreaks of witches' broom disease (*Crinipellis pernicioso*) of *Theobroma cacao* in Bahia, Brazil. *European J Plant Pathology* **105**, 167-75.
- Arruda MCC, Ferreira MASV, Miller RNG, Resende MLV, Felipe MSS, 2003. Nuclear and mitochondrial rDNA variability in *Crinipellis pernicioso* from different geographic origins and hosts. *Mycological Research* **107**, 25–37.
- Arruda MCC, Sepulveda GFC, Miller RNG, Ferreira MASV, Santiago DVR, Resende MLV, Dianese JC, Felipe MSS, 2005. *Crinipellis brasiliensis*, a new species based on morphological and molecular data. *Mycologia* **97**, 1348-61.
- Baker RED, Holliday P, 1957. Witches' broom disease of cacao (*Marasmius pernicioso* Stahel). Phytopathological papers no. 2. The Commonwealth Mycological Institute, Kew, 42 p.
- Bartley BGD, 2005. The genetic diversity of cacao and its utilization. CABI Publishing, Oxforshire, UK.
- Bastos CN, Evans HC, 1985. A new pathotype of *Crinipellis pernicioso* (witches' broom disease) on solanaceous hosts. *Plant Pathology* **34**, 306–12.
- Bowers JH, Bailey BA, Hebbar PK, Sanogo S, Lumsden RD, 2001. The impact of plant diseases on world chocolate production. *Plant Health* doi: 10.1094/PHP-2001-0709-01-RV.
- Cornuet JM, Piry S, Luikart G, Estoup A, Solignac M, 1999. New methods employing multilocus genotypes to select or exclude populations as origins of individuals. *Genetics* **153**, 1989-2000.
- Efron B, 1983. Estimating the error rate of a prediction rule-improvement on cross-validation. *J American Statistical Association* **78**, 316-31.
- El Mousadik A, Petit RJ, 1996. Chloroplast DNA phylogeography of the argan tree of Morocco. *Molecular Ecology* **5**, 547–55.

Giraud T, Refrégier G, Le Gac M, de Vienne DM, Hood ME, 2008. Speciation in fungi. *Fungal Genetics Biology* **45**, 791–802.

Godar J, Tizado EJ, Pokorny B, 2012. Who is responsible for deforestation in the Amazon? A spatially explicit analysis along the Transamazon Highway in Brazil. *Forest Ecology Management* **267**, 58–73.

Goudet J, 1995. FSTAT version 1.2: a computer program to calculate *F*-statistics. *J Heredity* **86**, 485–6. <http://www.unil.ch/popgen/software/fstat.htm>.

Gramacho KP, Risterucci AM, Lanaud C, Lima LS, Lopes UV, 2007. Characterization of microsatellites in the fungal plant pathogen *Crinipellis pernicioso*. *Molecular Ecology Notes* **7**, 153-5.

Griffith GW, Hedger JN, 1994a. The breeding biology of biotypes of the witches' broom pathogen of cocoa, *Crinipellis pernicioso*. *Heredity* **72**, 278-89.

Griffith GW, Hedger JN, 1994b. Spatial distribution of mycelia of the liana (L-) biotype of the agarica *Crinipellis pernicioso* (Stahel) Singer in tropical forest. *New Phytologist* **127**, 243-59.

Griffith GW, Hedger JN, 1994c. Dual culture of *Crinipellis pernicioso* and potato callus. *European Journal of Plant Pathology* **100**, 371-9.

Halkett F, Rivas CD, Platero GG, Zapater MF, Abadie C, Carlier J, 2010. Isolation by distance and genetic discontinuities in recently established populations of the plant pathogenic fungus *Mycosphaerella fijiensis*. *Molecular Ecology* **19**, 3909-23.

Hardy O, Vekemans X, 2002. SPAGeDi 1.2: a versatile computer program to analyse spatial genetic structure at the individual or population levels. *Molecular Ecology Notes* **2**, 618-20. <http://www.ulb.ac.be/sciences/lagev/software.html>.

Homewood B, 1991. Fungus threat to Brazil's 300 million cocoa trees. *New Scientist* **1778** (July 1991), 16.

Kalinowski ST, Taper ML, Marshall TC, 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* **16**, 1099-106.

Loiselle BA, Sork VL, Nason J, Graham C, 1995. Spatial genetic structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American J Botany* **82**, 1420-5.

- Motamayor JC, Lachenaud P, Mota JWS, Loor R, Kuhn DN, Brown JS, Schnell RJ, 2008. Geographic and genetic population differentiation of the Amazonian chocolate tree (*Theobroma cacao* L). *Plos One* **3**, e3311.
- Piry S, Alapetite A, Cornuet JM, Paetkau D, Baudouin L, Estoup A, 2004. GENECLASS 2: a software for genetic assignment and first-generation migrant detection. *J Heredity* **95**, 536-9.
- Ploetz RC, Schnell RJ, Ying Z, Zheng Q, Olano CT, Motamayor JC, Johnson ES, 2005. Analysis of molecular diversity in *Crinipellis pernicioso* with AFLP markers. *European J Plant Pathology* **111**, 317-32.
- Purdy LH, Schmidt RA, 1996. Status of cacao witches broom: biology, epidemiology, and management. *Annual Review Phytopathology* **34**, 573-94.
- Rincones J, Meinhardt LW, Vidal BC, Pereira GAG, 2003. Electrophoretic karyotype analysis of *Crinipellis pernicioso*, the causal agent of witches' broom disease of *Theobroma cacao*. *Mycological Research* **107**, 452-8.
- Rincones J, Mazotti GD, Griffith GW, Pomela A, Figueira A, Leal Jr. GA, Queiroz MV, Pereira JF, Azevedo RA, Pereira GA, Meinhardt LW, 2006. Genetic variability and chromosome-length polymorphisms of the witches' broom pathogen *Crinipellis pernicioso* from various hosts in South America. *Mycological Research* **110**, 821-32.
- Rorer JB, 1913. The Surinam witch-broom disease of cacao. Board of Agriculture. Trinidad & Tobago. Circular no. 10 (July1913) 13 pp.
- Santana MF, Araújo EF, Souza JT, Mizubuti ESG, Queiroz MV, 2012. Development of molecular markers based on retrotransposons for the analysis of genetic variability in *Moniliophthora pernicioso*. *European J Plant Pathology* **134**, 497-507.
- Sereno ML, Vencovsky R, Albuquerque PSB, Figueira A, 2006. Genetic diversity and natural population structure of cacao (*Theobroma cacao* L.) from the Brazilian Amazon evaluated by microsatellite markers. *Conservation Genetics* **7**, 13-24.
- Silva P, 1987, Cacao e lagartão e vassoura-de-bruxa: registros feitos por Alexandre Rodrigues Ferreira em 1785 e 1787 na Amazonia. *Boletim técnico CEPEC* (Itabuna, Brazil) **146**, 21 p.
- Silva JRQ, Figueira A, Pereira GAG, Albuquerque PSB, 2008. Development of novel microsatellites from *Moniliophthora pernicioso*, causal agent of the witches' broom disease of *Theobroma cacao*. *Molecular Ecology Resources* **8**, 783-5.

Stukenbrock EH, Bataillon T, 2010. A population genomics perspective on the emergence and adaptation of new plant pathogens in agro-ecosystems. *PLoS Pathogen* **8**(9), e1002893. doi:10.1371/journal.ppat.1002893.

Tarnowski TL, Ploetz RC, 2008. A multilocus phylogeny of the biotypes of *Moniliophthora perniciosa*, cause of witches' broom on cacao. *Phytopathology* **98**:S155.

Thomas E, van Zonneveld M, Loo J, Hodgkin T, Galluzzi G, van Etten J, 2012. Present spatial diversity patterns of *Theobroma cacao* L. in the Neotropics reflect genetic differentiation in Pleistocene refugia followed by human-influenced dispersal. *Plos One* **7**, e47676.

Wheeler BJE, Soares C, 1993. Disease gradients of *Crinipellis perniciosa* on cocoa seedlings. In: Rudgard SA, Maddison AC, Andebrhan T, eds. *Disease Management in Cocoa: Comparative epidemiology of witches' broom*. London, UK: Chapman & Hall, 157–64.

White TJ, Bruns T, Lee S, Taylor J, 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. *PCR Protocols: a guide to methods and applications*. New York, USA: Academic Press, 315-22.

Figure Legends

Figure 1. Approximate location of collection of the *Moniliophthora perniciosa* isolates from the C-, S- and L-biotypes.

Figure 2. Maximum likelihood tree of 33 *M. perniciosa* sequences (704 bp alignment), rooted with an unknown *Moniliophthora* species (MCA2500). Bootstrap values (% from 1000 replicates) are shown at salient nodes. Scalebar indicates substitutions per site. **B** indicates host Bignoniaceae and **M** indicates host in Malpigiaceae. **M*** was named *Moniliophthora brasiliensis* by Arruda *et al.* (2005).

Figure 3. Correlogram of pairwise co-ancestry coefficient (θ_{xy}) among the unique 56 C-biotype multilocus genotypes. Co-ancestry coefficient (θ_{xy}) was estimated as Loiselle *et al.* (1995) for the 56 C-biotype multilocus genotypes (one was excluded due to the lack of the spatial position) for each distance class within populations exclusively from the Amazon region, estimated using SPAGeDi 1. The solid line represents the average θ_{xy} value, whereas the dashed lines represent

the 95% (two-tailed) confidence interval of the average θ_{xy} distribution calculated by 1,000 permutations of individuals among of spatial distance.

Supporting Information

Figure S1

Analysis of identity of *Moniliophthora perniciosa* isolates by rDNA-ITS region. A and B. Amplification of the rDNA ITS1-5.8S-ITS2 region of 61 C-biotype isolates of *M. perniciosa*. C and D. Digestion of the amplified rDNA ITS1-5.8S-ITS2 region with *MspI*. E and F. Digestion of the amplified rDNA ITS1-5.8S-ITS2 region with *HinfI*.

Table S1

Description of isolates *Moniliophthora perniciosa* C-, S- and L- biotypes analyzed. Information about the code adopted (ID), previous identification, host and biotype, locale of isolate collection with state and country and year of collection; followed by geographical coordination of the site.

Table S2

Private alleles identified from microsatellite loci for isolates (coded as Table S1).

Table S3

Multilocus genotype with the respective identical individual and original location of collection.

Table S4

Assignment test of the 64 unique multilocus genotypes from the C-biotype. Results from individual test or as a group, in comparison with the S- and L- biotypes.

Table S5

Results of the assignment test of each of the seven unique multilocus genotypes from the C-biotype of Rondonia population; eight from Acre; and 11 from West Amazon in comparison with the other biotypes.

Table 1. Summary of sampled biotypes of *Moniliophthora perniciosa* (L-, S-, and C-biotypes, with seven geographic populations), with total number of isolates; locale of original collection with number of isolate identification (see Table S1); number of of multilocus genotypes; and number of identical individual for each multilocus genotype using the program Cervus 3.0.

Biotypes/ C-biotype populations	Locale of collection (number of isolate identification)	Total number of isolates	Multilocus genotypes	Number of Identical Individuals (clones)
L-Biotype ^x	Pichilingue (112, 113, 114, 115), Jauneche (116), San Carlos, Napo (117), Ecuador ^x	6	6	0
S-Biotype	Gandu, BA (126), Manaus, AM ^z (119, 120, 121, 128, 129, 130, 131), Rio Pomba, MG ^y (125), Viçosa, MG ^y (118, 123, 124, 127), unknown (122)	14	13	1
C-Biotype	--	111	64	47
Central Pará	Anapú (1); Altamira (2, 3); Baião (22); Brasil Novo (4, 5); Medicilândia (6, 7, 8,	20	12	8

(Transamazon region)	9, 10, 11); Mocajuba (23, 24); Trairão (25, 26, 27); Uruará (12, 13, 14)			
Pará (Lower Amazon)	Alenquer (15, 16, 107, 108, 109); Cametá (19, 20, 21); Óbidos (106); Santarém (17, 18)	11	9	2
East Amazonas	Augusto Montenegro (33, 34, 35, 36, 37, 38, 39); Boa Vista (32); Cepiac/ERNEG (45, 46); Costa da Conceição (43, 44); Manaus (28, 29,30, 31); Uricurituba (40, 41, 42)	19	10	9
West Amazonas (Upper Solimões)	Atalaia do Norte (105); Benjamim Constant (104); São Paulo de Olivença (102, 103); Tabatinga (92, 93, 94, 95, 96, 97, 98, 99, 100, 101); Agua Blanca, Ecuador ^x (110); San Carlos, Ecuador ^x (111)	16	11	5
Rondonia	Ariquemes (47, 48, 49, 50, 51); Cacaulândia (52, 53, 54); Cacoal (61); Jaru (55); Ji-Paraná (60, 62, 63); Ouro Preto D'Oeste (56, 57, 58, 59)	17	7	10
Acre	Assis Brasil (68, 69, 72, 73); Brasiléia (70, 71); Marechal Thaumaturgo (64, 65,	10	8	2

66); Plácido de Castro (67)

Bahia	Aurelino Leal (91); Camaçan (79, 87); Camamú (90); Canavieiras (78); Floresta Azul (85, 86); Gandú (84); Ilhéus (76); Itabuna (75); Itajuípe (74); Itapebi (77); Teolândia (89); Una (80); Uruçuca (81, 82, 83); Wenceslau Guimarães (88)	18	7	11
Overall Total		131	83	48

^x – isolates collected originally in Ecuador;

^y – MG = Brazilian state of Minas Gerais;

^z – AM = Brazilian state of Amazonas.

Table 2. Information about sample size (n), with results of allele richness (R); expected heterozygosity (H_e); genotypic diversity (D_g), effective number of alleles (A), and fixation index (F) for the L-, S- and C-biotypes (all genotypes) and the seven geographic populations of isolates from the C-biotype of *Moniliophthora perniciosa*, as well as for unique multilocus genotype. The parameter R was estimated for nine or six individuals, when all or unique multilocus genotypes, respectively, were used in the analyses. SE is the standard error. * $P < 0.05$.

All genotypes					Only unique multilocus genotypes				
Population	n	$R \pm SE$	$H_e \pm SE$	D_g	n	$A \pm SE$	$R \pm SE$	$H_e \pm SE$	$F \pm SE$
L-Biotype	6	3.82±0.05	0.69±0.06	1.00	6	3.82±0.50	3.82±0.50	0.66±0.04	0.74±0.12
S-Biotype	14	3.56±0.90	0.56±0.15	0.99	13	3.73±0.59	3.43±0.48	0.57±0.09	0.73±0.17
C-Biotype	111	1.90±0.12	0.29±0.04	-	64	2.73±0.24	2.37±0.16	0.43±0.06	0.92±0.05
(C) Central Pará (Transamazon region)	20	1.80±0.26	0.26±0.09	0.94	12	1.91±0.21	1.81±0.18	0.27±0.07	0.86±0.08
(C) Pará (Lower Amazon)	11	1.91±0.41	0.39±0.15	0.93	9	1.91±0.21	1.91±0.21	0.40±0.08	0.92±0.08

(C) East Amazonas	19	1.79±0.30	0.27±0.10	0.87	10	1.91±0.25	1.82±0.24	0.28±0.08	0.87±0.09
(C) West Amazonas (Upper Solimões)	16	1.71±0.37	0.24±0.13	0.95	11	1.73±0.24	1.70±0.23	0.25±0.08	1.00±0.00
(C) Rondonia	17	1.85±0.24	0.26±0.09	0.81	7	1.91±0.16	1.91±0.12	0.37±0.07	0.93±0.06
(C) Acre	10	2.09±0.43	0.31±0.13	0.96	8	2.09±0.21	2.09±0.21	0.35±0.07	0.77±0.13
(C) Bahia	18	2.14±0.38	0.32±0.12	0.77	7	2.27±0.27	2.27±0.27	0.45±0.09	1.00±0.00

Table 3. Results of the assignment test of each individual multilocus genotype from the L-biotype, S-biotype, C-biotype of East Amazonas Central, Pará of Transamazon region, Pará of lower Amazon and from Bahia of *Moniliophthora perniciosa* and as a group in comparison with genotypes from the other biotypes or populations (for the C-biotype).

Assigned L-biotype	1 st rank	Score (%)	Assigned S-biotype s	1 st rank	Score (%)	Assigned East Amazonas (C)	1 st rank	Score (%)
112 Pichilingue (L)	131 Manaus, AM (S)	99.5	118 Viçosa, MG	117 San Carlos (L)	100.0	28 Manaus, AM	92 Tabatinga, AM	71.6
113 Pichilingue (L)	131 Manaus, AM (S)	99.8	119 Manaus, AM	91 Aurelino Leal, BA (C)	50.1 ^y	29 Manaus, AM	63 Ji-Paraná, RO	88.0
114 Pichilingue (L)	131 Manaus, AM (S)	100.0	121 Manaus, AM	117 San Carlos (L)	100.0	30 Manaus, AM	63 Ji-Paraná, RO	89.7
115 Pichilingue (L)	63 Ji-Paraná, RO (C)	99.4	122 MG	117 San Carlos (L)	100.0	31 Manaus, AM	92 Tabatinga, PA	95.8

116 Jauneche (L)	131 Manaus, AM (S)	51.6 ^z	123 Viçosa, MG	117 San Carlos (L)	51.2 ^w	32 Boa Vista, AM	73 Assis Brasil, AC	65.8 ^x
117 San Carlos (L)	91 Aurelino Leal, BA (C)	92.7	124 Viçosa, MG	63 Ji-Paraná, RO (C)	99.9	33 A. Montenegro, AM	73 Assis Brasil, AC	81.0
L-Biotype isolates	131 Manaus, AM (S)	100.0	125 Rio Pomba, BA	117 San Carlos (L)	100.0	39 A. Montenegro, AM	63 Ji-Paraná, RO	45.3 ^u
			126 Gandu, BA	117 San Carlos (L)	100.0	43 Costa da Conceição, AM	73 Assis Brasil, AC	48.2 ^c
			127 Viçosa, MG	117 San Carlos (L)	100.0	44 Costa da Conceição, AM	73 Assis Brasil, AC	93.4
			128 Manaus, AM	117 San Carlos (L)	99.7	45 Ceplac/ERNEG, AM	73 Assis Brasil, AC	89.5
			129 Manaus, AM	117 San Carlos (L)	91.1	East Amazonas isolates	73 Assis Brasil, AC	94.2
			130 Manaus, AM	117 San Carlos (L)	100.0			
			131 Manaus, AM	117 San Carlos (L)	100.0			
			S-Biotype isolates	117 San Carlos (L)	100.0			
Assigned Central Pará (C)	1st rank	Score (%)	Assigned Lower Amazon (C)	1st rank	Score (%)	Assigned Bahia (C)	1st rank	Score (%)
1 Anapu, PA	73 Assis Brasil, AC	81.0	15 Alenquer, PA	73 Assis Brasil, AC	90.7	74 Itajuípe	92 Tabatinga, AM	99.2
2 Altamira, PA	73 Assis Brasil, AC	94.6	16 Alenquer, PA	73 Assis Brasil, AC	95.6	77 Itapebi	92 Tabatinga, AM	53.4 ^p
3 Altamira, PA	63 Ji-Paraná, RO	93.7	17 Santarém, PA	63 Ji-Paraná, RO	81.2	81 Uruçuca	63 Ji-Paraná, RO	99.2
4 Brasil Novo, PA	73 Assis Brasil, AC	50.9 ^v	18 Santarém, PA	45 CEPLAC Erneg, AM	91.6	84 Gandu	92 Tabatinga, AM	86.0
6 Medicilândia, PA	73 Assis Brasil, AC	69.6 ^t	19 Cametá, PA	73 Assis Brasil, AC	97.7	87 Camacan	63 Ji-Paraná, RO	78.5
7 Medicilândia, PA	45 Ceplac/ERNEG, AM	84.2	106 Óbidos, PA	110 A. Blanca, Manabi, EC	100.0	90 Camacan	63 Ji-Paraná, RO	71.0
10 Medicilândia, PA	45 Ceplac/ERNEG, AM	57.6 ^s	107 Alenquer, PA	110 A. Blanca, Manabi, EC	100.0	91 Aurelino Leal	92 Tabatinga, AM	100.0

11 Medicilândia, PA	63 Ji-Paraná, RO	73.9	108 Alenquer, PA	110 A. Blanca, Manabi, EC	100.0	Bahia group	92 Tabatinga, AM	100.0
12 Uruará, PA	45 Ceplac/ERNEG, AM	64.9 ^f	109 Duca Alenquer, PA	110 A. Blanca, Manabi, EC	100.0			
14 Uruará, PA	73 Assis Brasil, AC	50.8 ^q	Lower Amazon group	63 Ji-Paraná, RO	100.0			
26 Trairão, PA	73 Assis Brasil, AC	84.5						
27 Trairão, PA	45 Ceplac/ERNEG, AM	97.9						
Central Pará group	45 Ceplac/ERNEG, AM	99.1						

^z2nd score = 43.1% for 63 Ji-Paraná, RO (C);

^y2nd score = 49.1% for 117 San Carlos (L);

^w2nd score = 40.1% for 73 Assis Brasil, AC;

^x2nd score = 43.4% for 73 Assis Brasil, AC;

^u2nd score = 31.8% for 73 Assis Brasil, AC;

^v2nd score = 41.1% for 45 Ceplac/ERNEG, AM;

^t2nd score = 22.9% for 63 Ji-Paraná, RO;

^s2nd score = 21.7% for 92 Tabatinga, AM;

^r2nd score = 20.0% for 92 Tabatinga, AM;

^q2nd score = 31.4% for 45 Ceplac/ERNEG, AM;

^p2nd score = 45.9% for 73 Assis Brasil, AC.





