

## Aberystwyth University

### *Application of genomics-assisted breeding for generation of climate resilient crops: Progress and prospects*

Kole, Chittaranjan; Muthamilarasan, Mehanathan; Henry, Robert; Edwards, David; Sharma, Rishu; Abberton, Michael Terence; Batley, Jacqueline; Bentley, Alison; Blakeney, Michael; Bryant, John; Cai, Hongwei; Cakir, Mehmet; Cseke, Leland J.; Cockram, James; Oliveira, Antonio C.; Pace, Ciro D.; Dempewolf, Hannes; Ellison, Shelby; Gepts, Paul; Greenland, Andy J.

*Published in:*  
Frontiers in Plant Science

*DOI:*  
[10.3389/fpls.2015.00563](https://doi.org/10.3389/fpls.2015.00563)

*Publication date:*  
2015

*Citation for published version (APA):*

Kole, C., Muthamilarasan, M., Henry, R., Edwards, D., Sharma, R., Abberton, M. T., Batley, J., Bentley, A., Blakeney, M., Bryant, J., Cai, H., Cakir, M., Cseke, L. J., Cockram, J., Oliveira, A. C., Pace, C. D., Dempewolf, H., Ellison, S., Gepts, P., ... Prasad, M. (2015). Application of genomics-assisted breeding for generation of climate resilient crops: Progress and prospects. *Frontiers in Plant Science*, 6, Article 563.  
<https://doi.org/10.3389/fpls.2015.00563>

#### **Document License** CC BY

#### **General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

## Application of genomics-assisted breeding for generation of climate resilient crops: Progress and prospects

Chittaranjan Kole, Mehanathan Muthamilarasan, Robert Henry, David Edwards, Rishu Sharma, Michael Abberton, Jacqueline Batley, Alison Bentley, Michael Blakeney, John Bryant, Hongwei Cai, Mehmet Cakir, Leland J Cseke, James Cockram, Antonio Costa de Oliveira, Ciro De Pace, Hannes Dempewolf, Shelby Ellison, Paul Gepts, Andy Greenland, Anthony Hall, Kiyosumi Hori, Glenn Thomas Howe, Stephen Hughes, Mike W Humphreys, Massimo Iorizzo, Abdelbagi M. Ismail, Athole Marshall, Sean Mayes, Henry T Nguyen, Francis Chuks Ogbonnaya, Rodomiro Ortiz, Andrew H. Paterson, Philipp W. Simon, Joe Tohme, Roberto Tuberosa, Babu Valliyodan, Rajeev K Varshney, Stan D Wullschleger, Masahiro Yano and Manoj Prasad

Journal Name: Frontiers in Plant Science

ISSN: 1664-462X

Article type: Review Article

Received on: 25 May 2015


Accepted on: 08 Jul 2015

Provisional PDF published on: 08 Jul 2015

Frontiers website link: [www.frontiersin.org](http://www.frontiersin.org)

Citation: Kole C, Muthamilarasan M, Henry R, Edwards D, Sharma R, Abberton M, Batley J, Bentley A, Blakeney M, Bryant J, Cai H, Cakir M, Cseke LJ, Cockram J, Oliveira AC, Pace CD, Dempewolf H, Ellison S, Gepts P, Greenland A, Hall A, Hori K, Howe GT, Hughes S, Humphreys MW, Iorizzo M, Ismail AM, Marshall A, Mayes S, Nguyen HT, Ogbonnaya FC, Ortiz R, Paterson AH, Simon PW, Tohme J, Tuberosa R, Valliyodan B, Varshney RK, Wullschleger SD, Yano M and Prasad M(2015) Application of genomics-assisted breeding for generation of climate resilient crops: Progress and prospects. *Front. Plant Sci.* 6:563. doi:10.3389/fpls.2015.00563

Copyright statement: © 2015 Kole, Muthamilarasan, Henry, Edwards, Sharma, Abberton, Batley, Bentley, Blakeney, Bryant, Cai, Cakir, Cseke, Cockram, Oliveira, Pace, Dempewolf, Ellison, Gepts, Greenland, Hall, Hori, Howe, Hughes, Humphreys, Iorizzo, Ismail, Marshall, Mayes, Nguyen, Ogbonnaya, Ortiz, Paterson, Simon, Tohme, Tuberosa, Valliyodan, Varshney, Wullschleger, Yano and Prasad. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



# Application of genomics-assisted breeding for generation of climate resilient crops: Progress and prospects

Chittaranjan Kole<sup>1\*</sup>, Mehanathan Muthamilarasan<sup>2</sup>, Robert Henry<sup>3</sup>, David Edwards<sup>4</sup>, Rishu Sharma<sup>5</sup>, Michael Abberton<sup>6</sup>, Jacqueline Batley<sup>7</sup>, Alison Bentley<sup>8</sup>, Michael Blakeney<sup>9</sup>, John Bryant<sup>10</sup>, Hongwei Cai<sup>11</sup>, Mehmet Cakir<sup>12</sup>, Leland J. Cseke<sup>13</sup>, James Cockram<sup>14</sup>, Antonio de Costa Oliveira<sup>15</sup>, Ciro De Pace<sup>16</sup>, Hannes Dempewolf<sup>17</sup>, Shelby Ellison<sup>18</sup>, Paul Gepts<sup>19</sup>, Andy Greenland<sup>20</sup>, Anthony Hall<sup>21</sup>, Kiyosumi Hori<sup>22</sup>, Glenn T. Howe<sup>23</sup>, Stephen Hughes<sup>24</sup>, Mike W Humphreys<sup>25</sup>, Massimo Iorizzo<sup>26</sup>, Abdelbagi M. Ismail<sup>27</sup>, Athole Marshall<sup>28</sup>, Sean Mayes<sup>29</sup>, Henry T. Nguyen<sup>30</sup>, Francis C. Ogbonnaya<sup>31</sup>, Rodomiro Ortiz<sup>32</sup>, Andrew H. Paterson<sup>32</sup>, Phillip W. Simon<sup>34</sup>, Joe Tohme<sup>35</sup>, Roberto Tuberosa<sup>36</sup>, Babu Valliyodan<sup>37</sup>, Rajeev Varshney<sup>38</sup>, Stan D. Wulschleger<sup>39</sup>, Masahiro Yano<sup>40</sup> and Manoj Prasad<sup>41\*</sup>

<sup>1\*</sup>Chittaranjan Kole, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur 741252, Dist. Nadia, West Bengal, India; [ckole2012@gmail.com](mailto:ckole2012@gmail.com)

<sup>2</sup>Mehanathan Muthamilarasan, National Institute of Plant Genome Research, Aruna Asaf Ali Marg, JNU Campus, New Delhi 110067, India; [muthamilarasan@nipgr.ac.in](mailto:muthamilarasan@nipgr.ac.in)

<sup>3</sup>Robert Henry, Queensland Alliance for Agriculture and Food Innovation, John Hay Building, Queensland Biosciences Precinct [#80], University of Queensland, St Lucia, QLD 4072, Australia; [robert.henry@uq.edu.au](mailto:robert.henry@uq.edu.au)

<sup>4</sup>Dave Edwards, School of Agriculture and Food Sciences, University of Queensland, Brisbane, QLD 4072, Australia; [dave.edwards@uq.edu.au](mailto:dave.edwards@uq.edu.au)

<sup>5</sup>Rishu Sharma, Department of Plant Pathology, Faculty of Agriculture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur 741252, Dist. Nadia, West Bengal, India; [rrishu.sharma90@gmail.com](mailto:rrishu.sharma90@gmail.com)

<sup>6</sup>Michael Abberton, Genetic Resources Centre, International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria; [M.Abberton@cgiar.org](mailto:M.Abberton@cgiar.org)

<sup>7</sup>Jacqueline Batley, Centre for Integrated Legume Research, University of Queensland, Brisbane, QLD 4072, Australia; [J.Batley@uq.edu.au](mailto:J.Batley@uq.edu.au)

<sup>8</sup>Alison Bentley, The John Bingham Laboratory, NIAB, Huntingdon Road, Cambridge CB3 0LE, United Kingdom; [alison.bentley@niab.com](mailto:alison.bentley@niab.com)

<sup>9</sup>Michael Blakeney, Centre for Commercial Law Studies, Roy Goode House, 13-14 Charterhouse Square, London EC1M 6AX, UK; [m.blakeney@qmul.ac.uk](mailto:m.blakeney@qmul.ac.uk)

<sup>10</sup>John Bryant, CLES, Hatherly Laboratories, University of Exeter, Exeter EX4 4PS, UK; [J.A.Bryant@exeter.ac.uk](mailto:J.A.Bryant@exeter.ac.uk)

<sup>11</sup>Hongwei Cai, Forage Crop Research Institute, Japan Grassland Agriculture and Forage Seed Association, Nasushiobara, Japan; [hcai@jfsass.or.jp](mailto:hcai@jfsass.or.jp), and Department of Plant Genetics and Breeding, College of Agronomy and Biotechnology, China Agricultural University, Beijing, China; [caihw@cau.edu.cn](mailto:caihw@cau.edu.cn)

<sup>12</sup>Mehmet Cakir, School of Biological Sciences and Biotechnology, Faculty of Science and Engineering, Murdoch University, 90 South St., Murdoch, WA 6150, Australia; [M.Cakir@murdoch.edu.au](mailto:M.Cakir@murdoch.edu.au)

<sup>13</sup>Leland J. Cseke, The University of Alabama in Huntsville, Department of Biological Sciences, Huntsville, AL, USA; [csekel@uah.edu](mailto:csekel@uah.edu)

<sup>14</sup>James Cockram, The John Bingham Laboratory, NIAB, Huntingdon Road, Cambridge CB3 0LE, United Kingdom; [James.cockram@niab.com](mailto:James.cockram@niab.com)

<sup>15</sup>Antonio Costa de Oliveira, Plant Genomics Center, PO Box 354, FAEM, UFPEL, Brazil; [acostol@terra.com.br](mailto:acostol@terra.com.br)

<sup>16</sup>Ciro De Pace, Department of Agriculture, Forests, Nature and Energy (DAFNE), University of Tuscia, Viterbo, Italy; [depance@unitus.it](mailto:depance@unitus.it)

<sup>17</sup>Hannes Dempewolf, Global Crop Diversity Trust, C/O FAO, Viale delle Terme di Caracalla, 00153 Rome, Italy; [hannes.dempewolf@croptrust.org](mailto:hannes.dempewolf@croptrust.org)

<sup>18</sup>Shelby Ellison, Department of Horticulture, University of Wisconsin, 1575 Linden Drive, Madison, WI 53706, USA; [slrepinski@wisc.edu](mailto:slrepinski@wisc.edu)

<sup>19</sup>Paul Gepts, University of California, Department of Plant Sciences, Section of Crop and Ecosystem Sciences, 1 Shields Avenue, Davis, CA, USA; [plgepts@ucdavis.edu](mailto:plgepts@ucdavis.edu)

<sup>20</sup>Andy Greenland, The John Bingham Laboratory, NIAB, Huntingdon Road, Cambridge CB3 0LE, UK; [Andy.greenland@niab.com](mailto:Andy.greenland@niab.com)

<sup>21</sup>Anthony Hall, 2922 Lindsay Lane, Quincy, CA 95971-9602, USA; [anthony.hall@ucr.edu](mailto:anthony.hall@ucr.edu)

<sup>22</sup>Kiyosumi Hori, National Institute of Agrobiological Sciences, 2-1-2 Kannondai, Tsukuba, Ibaraki 305-8602, Japan; [horikiyo@affrc.go.jp](mailto:horikiyo@affrc.go.jp)

<sup>23</sup>Glenn Thomas Howe, Oregon State University, Department of Forest Ecosystems and Society, Corvallis, OR, USA; [glenn.howe@oregonstate.edu](mailto:glenn.howe@oregonstate.edu)

<sup>24</sup>Stephen Hughes, University of Exeter, C/O Byrne House, St Germans Road, Exeter, EX4 4PJ, UK; [S.G.Hughes@exeter.ac.uk](mailto:S.G.Hughes@exeter.ac.uk)

<sup>25</sup>Mike W. Humphreys, Institute of Biological, Environmental and Rural Sciences (IBERS), Aberystwyth University, Gogerddan, Aberystwyth, Ceredigion, Wales, UK, SY23 3EE; [mkh@aber.ac.uk](mailto:mkh@aber.ac.uk)

<sup>26</sup>Massimo Iorizzo, Department of Horticulture, University of Wisconsin, 1575 Linden Drive, Madison, WI 53706, USA; [iorizzo@wisc.edu](mailto:iorizzo@wisc.edu)

<sup>27</sup>Abdelbagi M. Ismail, International Rice Research Institute, Manila, Philippines; [a.ismail@irri.org](mailto:a.ismail@irri.org)

<sup>28</sup>Athole Marshall, Institute of Biological, Environmental and Rural Sciences (IBERS), Aberystwyth University, Reception, Penglais, Aberystwyth, Ceredigion, UK, SY23 3FL; [thm@aber.ac.uk](mailto:thm@aber.ac.uk)

<sup>29</sup>Sean Mayes, Crops for the Future, Jalan Broga, 43500 Semenyih, Malaysia; [Sean.Mayes@nottingham.ac.uk](mailto:Sean.Mayes@nottingham.ac.uk)

<sup>30</sup>Henry T. Nguyen, National Center for Soybean Biotechnology and Division of Plant Science, University of Missouri, Columbia, MO, USA; [nguyenhenry@missouri.edu](mailto:nguyenhenry@missouri.edu)

<sup>31</sup>Francis C. Ogonnaya, Grains Research and Development Corporation (GRDC), Barton ACT 2600 Australia, PO Box 5367, Kingston, ACT, 2604 Australia; [Francis.Ogonnaya@grdc.com.au](mailto:Francis.Ogonnaya@grdc.com.au)

<sup>32</sup>Rodomirol Ortiz, Swedish University of Agricultural Sciences, SLU, Department of Plant Breeding, Sundvagen 14 Box 101, SE 23053, Sweden; [rodomirol.ortiz@slu.se](mailto:rodomirol.ortiz@slu.se)

<sup>33</sup>Andrew H. Paterson, Plant Genome Mapping Laboratory, University of Georgia, Athens, GA, USA; [paterson@plantbio.uga.edu](mailto:paterson@plantbio.uga.edu)

<sup>34</sup>Philipp W. Simon, USDA-ARS, Department of Horticulture, University of Wisconsin, 1575 Linden Drive, Madison, WI 53706, USA; [Philipp.Simon@ars.usda.gov](mailto:Philipp.Simon@ars.usda.gov)

<sup>35</sup>Joe Tohme, Agrobiodiversity and Biotechnology Project, Centro Internacional de Agricultura Tropical, CIAT, AA 6713, Cali, Columbia; [J.Tohme@cgiar.org](mailto:J.Tohme@cgiar.org)

<sup>36</sup>Roberto Tuberosa, Department of Agroenvironmental Sciences and Technology Via Fanin, 44 40127-Bologna, Italy; [roberto.tuberosa@unibo.it](mailto:roberto.tuberosa@unibo.it)

<sup>37</sup>Babu Valliyodan, National Center for Soybean Biotechnology and Division of Plant Science, University of Missouri, Columbia, MO, USA; [valliyodanb@missouri.edu](mailto:valliyodanb@missouri.edu)

<sup>38</sup>Rajeev K. Varshney, Bldg # 300, Center of Excellence in Genomics (CEG), International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru 502 324, Greater Hyderabad, India; [r.k.varshney@cgiar.org](mailto:r.k.varshney@cgiar.org)

<sup>39</sup>Stan D. Wullschlegel, Environmental Sciences Division, Climate Change Science Institute (CCSI), Oak Ridge National Laboratory, Building 2040, Room E212, MS 6301, Oak Ridge, TN 37831-6301, USA; [wullschlegsd@ornl.gov](mailto:wullschlegsd@ornl.gov)

<sup>40</sup>Masahiro Yano, National Agriculture and Food Research Organization (NARO), Institute of Crop Science, Tsukuba, Japan; [myano@affrc.go.jp](mailto:myano@affrc.go.jp)

<sup>41</sup>Manoj Prasad, National Institute of Plant Genome Research, Aruna Asaf Ali Marg, JNU Campus, New Delhi 110067, India; [manoj\\_prasad@nipgr.ac.in](mailto:manoj_prasad@nipgr.ac.in)

**\*Correspondence**

Prof. Chittaranjan Kole  
Bidhan Chandra Krishi Viswavidyalaya  
Mohanpur 741252  
Dist. Nadia, West Bengal, India  
[ckole2012@gmail.com](mailto:ckole2012@gmail.com)

**Dr. Manoj Prasad**

National Institute of Plant Genome Research  
Aruna Asaf Ali Marg,  
New Delhi 110067, India  
[manoj\\_prasad@nipgr.ac.in](mailto:manoj_prasad@nipgr.ac.in)

## Abstract

Climate change affects agricultural productivity worldwide. Increased prices of food commodities are the initial indication of drastic edible yield loss, which is expected to surge further due to global warming. This situation has compelled plant scientists to develop climate change-resilient crops, which can withstand broad-spectrum stresses such as drought, heat, cold, salinity, flood and submergence, and pests along with increased productivity. Genomics appears to be a promising tool for deciphering the stress responsiveness of crop species with adaptation traits or in wild relatives towards identifying underlying genes, alleles or quantitative trait loci. Molecular breeding approaches have been proven helpful in enhancing the stress adaptation of crop plants, and recent advancement in next-generation sequencing along with high-throughput sequencing and phenotyping platforms have transformed molecular breeding to genomics-assisted breeding (GAB). In view of this, the present review elaborates the progress and prospects of GAB in improving climate change resilience in crop plants towards circumventing global food insecurity.

**Keywords:** Climate change, crop improvement, stress tolerance, breeding, genomics

## Introduction

Three major events in agricultural history, namely domestication, displacement of native crops by major crops along with genetically limited introductions of non-native species, and intensification of agricultural production through the Green Revolution have contributed significantly towards reduced genetic and trait diversity within major crop species. Despite this decrease in crop diversity, global production of the major staple crops was increased in the last century (Fischer et al., 2009). This increase in productivity has largely been driven by conventional plant breeding coupled to intensification and simplification of production systems. This includes selection for edible yield and adaptation, and against yield reducing factors such as susceptibility to pathogens as well as pests, and optimization of crop husbandry practices (through high inputs such as the use of fertilizers, herbicides, pesticides, and mechanization) to minimize the impact of environmental flux. However, selection under such ‘ideal’, high-input environments has led to the loss of certain genes which are responsible for efficiency or adaptation to stress(es) (Brown, 2003). This situation presents three potential challenges: (i) to modify the selection criteria to focus on efficiency or adaptation to stress(es) rather than total edible yield, (ii) to ensure the presence and efficiency of stress-tolerance genes and its exploitability in elite material and wider breeding germplasm, and (iii) to expand the use of minor crops, which may possess better nutrition quality, environmental sustainability or resilience and require lower inputs than major crops.

At present, global agriculture is facing a serious threat of climate change resulting in reduced productivity. Increasing food prices and greater global food insecurity are the outcomes of decreased productivity (FAO 2014) and this scenario, if persists, would lead to further increase in food prices in developed countries, and social unrest and famine in these regions. Climate change will affect food supply unless actions are taken to increase the resilience of crops as projections have shown a drastic decrease in the production of major cereals by 2020, including 9% for maize, 11% for rice and 14% for wheat (Hisas, 2011). Global warming, changes of

rainfall pattern and other extreme weather events may mostly contribute to this disaster, and the changing pattern of climate would result in increased attack of pathogens and pests. Moreover, the elevated CO<sub>2</sub> levels will reduce the nutritional quality of many crops, while some crops may become toxic due to changes in the chemical composition of their tissues (Dwivedi et al., 2013).

Therefore, increasing the resilience of crops to climate change is the prime need to ensure food and nutritional security, which could be achieved through genetic engineering-based approaches or molecular breeding strategies. Genetic engineering allows direct transfer of beneficial gene(s) or manipulation of existing gene(s) in the crop of interest for generating expected phenotype(s), whereas breeding approaches involve the improvement of germplasm through introduction of novel alleles into target crops by breeding. Since genetic modification remains controversial in a number of countries though it serves as an invaluable tool in tailoring modifications to produce alleles and phenotypes beyond the range available through exploitation of existing genetic variation, molecular breeding could offer an easy-to-accept approach for crop improvement.

### **Potential of genomics-assisted breeding in producing climate resilient crops**

Genomics offers tools to address the challenge of increasing food yield, quality and stability of production through advanced breeding techniques. Applications of DNA markers to facilitate marker-aided-selection (MAS) for crop improvement have been proved successful in crossbreeding. Advances in plant genomics provide further means to improve the understandings on crop diversity at species and gene levels, and offer DNA markers to accelerate the pace of genetic improvement (Muthamilarasan et al., 2013). A genomics-led breeding strategy for new cultivars commences by defining the stress(es) that will likely affect crop production and productivity under a certain climate change scenario. Data from multi-environment testing provide an opportunity for modeling “stress-impacts” on crops and target populations of environments. Plant breeders and genebank curators will search for morphological and physiological traits in available germplasm that could enhance crop adaptation under such climate variability. In this regard, crop physiology may help define the ideotypes to be pursued for enhancing such adaptation. Moreover, the use of geographic information systems and passport data can allow identification of accessions for stress-prone environments, whereas the available characterization, including DNA fingerprinting, and evaluation data as well as mapping of desired genes or quantitative trait loci (QTL) will assist in selecting promising accessions for further screening against specific stress(es). Similarly, precise phenotypic assessments and appropriate biometric analysis will assist in identifying unique responses of a set of genotypes in a given phenological stage influenced by variation of weather patterns. This information will be further used in genomics-aided breeding approaches such as genome-wide selection of promising germplasm for further use in crop breeding aiming at both population improvement and cultivar releases.

Genetic mapping and QTL analysis and association genetics (AG) have accelerated the dissection of genetic control of agricultural traits, potentially allowing MAS, QTL and AG studies or direct calculation and genomic selection (GS) of high value genotypes to be made in the context of breeding programs. Until recently, AG and GS were hampered by the need for very high marker density coverage of the genome. Advancement of next-generation sequencing (NGS) methods has facilitated the development of large-scale, genome-wide, high-throughput



markers such as single nucleotide polymorphisms (SNP), insertion-deletions (InDels), etc. even in relatively research-neglected crop species. Discovery of novel genes/alleles for any given trait could be then performed through genotyping-by-sequencing (GBS) approaches. Similarly, genome-wide association study (GWAS) identifies the genomic regions governing traits of interest by performing statistical associations between DNA polymorphisms and trait variations in diverse collection of germplasms that are genotyped and phenotyped for traits of interest. NGS coupled with GWAS increases the mapping resolution for precise location of genes/alleles/QTL (Ma et al., 2012; Liu et al., 2013; Varshney et al., 2014).

In course of evolution, nature has evolved new genes, and shuffled and selected these genes in a wide range of environments to produce the diversity evidenced in wild species. In contrast, the selection and domestication of crops by humans is relatively recent, having occurred over the last 10,000 years. During the domestication and breeding process, there has been a significant reduction of genetic diversity in major crops, alongside a selection for yield under highly managed agricultural environments. Currently, breeders are shuffling the combinations of relatively few alleles to produce enhanced combinations that provide increased yield and other attractive agronomic characteristics. In many large genome crop species, even this reshuffling process is limited by restricted recombination patterns within the species, leading to the consistent inheritance of blocks of genes, raising issues of linkage drag and fixed linkage blocks, which may not contain the best possible combination of alleles. Breaking down these linkage constraints will allow breeders to access novel combinations from within their current elite parents. The need to evaluate the genetics of the processes that allow genes to be recombined between parental genotypes in crops is a critical requisite. Genomics possesses the potential to increase the diversity of alleles available to breeders through mining of allied gene pools and genomes of crop wild relatives (CWRs). Genomics tools also enable rapid identification and selection of novel beneficial genes and their controlled incorporation into novel germplasm. In the next-generation genomics era, this technology will be used to safeguard the future through improved food security. Taken together, application of genomics for crop germplasm enhancement thus offers the greatest potential to increase food production in the coming decades. With continued rapid advances in genome technologies, the application of genomics to identify and transfer valuable agronomic genes from allied genepools and crop relatives to elite crops will increase in pace and assist in meeting the challenge of global food production.

## **Genomics of climate resilience in major crops**

The following section summarizes the state of knowledge of the genetic blueprints of many leading crops, together with information about breeding needs and priorities related to climate resilience. Genomic tools and resources are widely available and being employed in most of these plants and will soon be ubiquitous, aiding ‘MAS’ strategies that can be successful even based only on phenotypic information. Knowledge of gene functions is less consistent, leveraging to varying degrees of the accumulated information from botanical models. However, even in model crops, the exact functions of most genes remain unknown, and exploring the variations conferred during angiosperm diversification represents an opportunity to identify a host of solutions to agricultural challenges.

## **Cereals**

Cereals are a staple to billions and their production is increasingly threatened by the recent changes in weather patterns due to global warming, particularly in less-developed countries where the consequences of changing climate have devastating socio-economic impact. Reaching a level of cereal production sufficient to sustain an adequate level of global food security will require the effective integration of crossbreeding with ‘omics’ approaches that allow dissecting and more effectively manipulating the genetic make-up of adaptation to abiotic stresses (Langridge et al., 2011). In the past decade, genomics-based approaches have been extensively deployed to dissect the genetic make-up of abiotic stress adaptation and given the quantitative nature of abiotic stress tolerance, QTL have been the main target of research to identify the genetic loci regulating the adaptive response of cereal crops to unfavorable environmental conditions. This includes drought-adaptive traits (Serraj et al., 2009; Tuberosa 2012), root architecture (Wasson et al., 2012; Uga et al., 2013; Lynch et al., 2014), accumulation of water-soluble carbohydrates and their partitioning to storage organs (Landi et al., 2005; Salem et al., 2007; Snape et al., 2007; Rebetzke et al., 2008), abscisic acid concentration (Rebetzke et al., 2008; Rehman et al., 2011), stay-green (Yang et al., 2007; Borrell et al., 2014), canopy temperature (Lopes et al., 2014), and carbon isotope discrimination ( $\Delta^{13}\text{C}$ ) (Pinto et al., 2014).

Global warming is intimately associated with an increase in temperature that accelerates leaf senescence, disrupts starch accumulation and curtails yield, particularly when combined with drought. In wheat, a major QTL located on chromosome 4A explained 27% and 17% of phenotypic variance for reduction in yield under drought and heat stress, respectively (Pinto et al., 2014). The same study also identified common QTL for drought and heat stress adaptation on chromosomes 1B, 2B, 3B, 4B, and 7A. Yield QTL were shown to be associated with components of other traits, supporting the prospects for dissecting crop performance under abiotic stress conditions into physiological and genetic components in order to facilitate a strategic approach to breeding (Reynolds et al., 2008). Additional QTL with concurrent effects under both heat and drought conditions have been described by Wang et al. (2012).

In rice, the result of a study based on 227 intensively managed irrigated farms forecast a net negative impact on yield from the warming expected in the coming decades, and clearly show that diurnal temperature variation must be considered when investigating the impact of climate change (Welch et al., 2010). Higher temperatures are speculated to reduce rice grain yields through two main pathways: (i) high maximum temperatures that in combination with high humidity cause spikelet sterility, and (ii) increased nighttime temperatures, which may reduce assimilate accumulation (Wassmann et al., 2009).

Flooding is one of the abiotic stresses, whose frequency and intensity is increasing due to global warming and changes in rainfall patterns. Therefore, it is important to produce cereal crops with the ability to withstand the anoxic conditions associated with waterlogging and/or extended submergence. Among cereals, rice is more prone to submergence stress, which periodically affects approximately 15 million hectares of rain-fed lowland areas in Asia to cause annual losses in excess of US \$1 billion (Mackill et al. 2012). In rice, the *Sub1* QTL accounts for a major portion of variability for survival under prolonged submergence. Positional cloning of *Sub1* has revealed a cluster of three putative ethylene response factors (ERFs), namely *Sub1A*, *Sub1B*, and *Sub1C*. Further work unequivocally assigned the functional polymorphism to *Sub1A*

(Xu et al., 2006). Following the identification of *Sub1A* QTL, marker-aided backcrossing (MABC) was used to efficiently convert submergence-susceptible rice cultivars into tolerant cultivars in only three backcross generations. Accordingly, DNA markers were developed for introgressing *Sub1* into six popular cultivars to meet the needs of farmers in flood-prone regions (Bailey-Serres et al., 2010). This clearly demonstrate the effectiveness of MAS for introgressing agronomically beneficial QTL alleles into elite material. The success of this work is largely due to the major effect of *Sub1* QTL and the stability of its effect in different genetic backgrounds under submergence conditions. In maize, Mano et al., (2005a) identified QTL for adventitious root formation at the soil surface, one of the most important adaptations to soil waterlogging, which can severely impair root growth at an early stage, thus reducing the capacity of the plant to extract soil moisture at a later stage when water shortage is more likely to occur. Several QTL for adventitious root formation have been mapped, and a major QTL was mapped on chromosome 8 (Mano et al., 2005b).

Salinity is also an impact of global climate change, which affects over 20% of the world's agricultural soils and thereby affecting cultivation. In durum wheat (genome AABB), two major QTL have been shown to control Na<sup>+</sup> accumulation in shoot via Na<sup>+</sup> exclusion (James et al., 2006). Both exclusion genes represent introgressions from an accession of *Triticum monococcum* (genome AA). Remarkably, under standard conditions, durum wheat containing the salinity tolerant allele at *TmHKT1;5-A*, which is one of the two salt-tolerance loci showed the phenotype similar to durum wheat that lacked the beneficial allele at this locus. But under saline conditions, it outperformed its durum wheat parent, with increased yields of up to 25% (Munns et al., 2012). In barley, evaluation of a mapping population derived from a cross between a wild barley (*Hordeum vulgare* ssp. *spontaneum*) accession and cultivated barley (*H. vulgare*) allowed the identification of a major QTL capable of limiting Na<sup>+</sup> accumulation in the shoots under saline conditions (Shavrukov et al., 2010). In rice, several QTL for salinity tolerance have been identified (Wang et al., 2012) indicating that pyramiding by marker-assisted selection (MAS) of QTL can be applied to enhance salt tolerance of rice.

## Oilseeds and pulses

Oilseeds and pulses are major food crops, known for their unique protein and oil rich characteristics. Major biotic and abiotic stresses are the most serious production constraint for global oilseed and pulse production, and are predicted to worsen with anticipated climate change. Among the oilseeds, soybean has the highest protein content (40%) and the second highest oil content (20%). In spite of this importance, efforts are yet to be invested towards improving stress tolerance and other traits in soybean. *Phaseolus* beans are an essential part of the human diet and are a source of proteins, vitamins, and minerals (Gepts et al., 2008). Of the five domesticated *Phaseolus* species, common bean (*P. vulgaris* L.) is the economically important bean. Genetic studies and cultivar breeding in *P. vulgaris* have shown that heat and drought tolerance are under complex genetic control, although a single instance of a major gene has also been observed (Schneider et al., 1997; Asfaw et al., 2012). Selection of lines with improved drought adaptation has also been successful (Singh, 2007; Beebe et al., 2008; Urrea et al., 2009). Development of MAS methodology for drought adaptation has been initiated (Schneider et al., 1997; Asfaw et al., 2012) with the assistance of genomic resources developed through whole-genome sequencing of Andean (accession: G19833) (Schmutz et al., 2014) and Mesoamerican (accession: BAT93,

OAC Rex) bean genomes, and a bean breeder's genome toolbox and database (<http://phaseolusgenes.bioinformatics.ucdavis.edu/>).

In case of chickpea and pigeonpea, several abiotic and biotic stresses pose a threat to high and stable grain yields. To overcome these production constraints and meet the growing demand for these crops, efforts at national and international levels have led to the development of large-scale genetic and genomic resources (Varshney et al., 2013a). These resources have been used to understand the existing genetic diversity and exploit it in breeding programs. In chickpea, several intra- and inter-specific genetic maps have been developed (Gaur et al., 2011; Gujaria et al., 2011; Thudi et al., 2011; Hiremath et al., 2012) and genomic regions responsible for different biotic stresses (Anbessa et al., 2009; Kottapalli et al., 2009; Anuradha et al., 2011), abiotic stress (Rehman et al., 2011; Vadez et al., 2012) and agronomic traits (Cobos et al., 2009; Rehman et al., 2011; Bajaj et al., 2014, 2015; Kujur et al., 2015a, 2015b; Das et al., 2015) have been reported. In pigeonpea, more than 3000 SSR markers (Saxena et al., 2010; Bohra et al., 2011; Dutta et al., 2011), ESTs (Raju et al., 2010), 454/FLX transcript reads (Dubey et al., 2011; Dutta et al., 2011), transcriptome assemblies (Dubey et al., 2011; Kudapa et al., 2012) and SNPs (Saxena, 2008) have been developed for their use in genomics-assisted breeding for crop improvement.

The draft genome sequence of both Kabuli (<http://www.icrisat.org/gt-bt/ICGGC/GenomeSequencing.htm>) and Desi (<http://www.nipgr.res.in/CGWR/home.php>) chickpeas have recently been published (Varshney et al., 2013b; Jain et al., 2013). Similarly, International Initiative on Pigeonpea Genomics (IIPG, <http://www.icrisat.org/gt-bt/iipg/Home.html>) released the draft genome of pigeonpea (Varshney et al., 2012). These sequence data of chickpea and pigeonpea will assist in enhancing their crop productivity and lead to conserving food security in arid and semi-arid environments. Further, attempts have been made towards improvement of oilseed crops such as peanut (or groundnut) using genomics-assisted breeding. Large-scale genomic resources were developed during recent years to facilitate molecular breeding in peanut and QTL have been identified for stress adaptation related traits (Varshney et al., 2009; Gautami et al., 2012), rust and late leaf spot resistance (Khedikar et al., 2010; Sujay et al., 2012), and oil quality (Sarvamangala et al., 2011).

## **Millets**

Millets are small-grained graminaceous crops, well known for their water-use efficiency, excellent nutrient content, adaptation to a range of ecological conditions and ability to flourish in nutrient-poor soils. Foxtail millet, proso millet, pearl millet, barnyard millet, finger millet and kodo millet are few notable millet crops cultivated worldwide and of these, foxtail millet is considered as a C<sub>4</sub> crop model for studying the biology of other millets and biofuel grasses (Lata et al., 2013; Muthamilarasan and Prasad, 2015). Therefore, the Beijing Genomics Institute, China and the US Department of Energy - Joint Genome Institute have sequenced the foxtail millet genome (Zhang et al., 2012; Bennetzen et al., 2012). As foxtail millet serves as a rich source of genes, alleles, or QTL for genetic improvement of major cereals and bioenergy grasses, large-scale genomic resources were developed including simple sequence repeats (SSRs) (Pandey et al., 2013; Zhang et al., 2014), intron length polymorphisms (Muthamilarasan et al., 2014), eSSRs (Kumari et al., 2013), miRNA-based markers (Yadav et al., 2014) and transposable-elements

based markers (Yadav et al., 2015). Moreover, open access online databases such as foxtail millet marker database (FmMdb) (Suresh et al., 2014), foxtail millet miRNA database (FmMiRNADb) (Khan et al., 2014) and foxtail millet transposable elements-based marker database (FmTEMDb) (Yadav et al., 2015) have been constructed. In addition to development of these markers, their utility in population genetics, association mapping, comparative genomics and genomics-assisted breeding have also been demonstrated (Muthamilarasan and Prasad, 2015). An allele-specific marker developed from an SNP in *SiDREB2* gene linked to drought tolerance in foxtail millet (Lata et al., 2011) is being used in allele mining and MAS for drought tolerance (Lata and Prasad, 2012; Lata and Prasad, 2013).

In pearl millet, three major QTL for grain yield with low QTL  $\times$  environment interactions were identified across a range of post-flowering moisture environments (Bidingger et al., 2007). One of these major QTL accounted for up to 32% of the phenotypic variance of grain yield under drought. The effects of this QTL were validated in two independent MABC programs in which 30% improvement in general combining ability for grain yield expected from this QTL under terminal drought stress was recovered in introgression lines, based on the information provided by the markers flanking the QTL (Yadav et al., 2011). Compared to other crops, research on millets is at initial stage. Being predominantly climate resilient crops, millets could serve as valuable source of novel genes, alleles and QTL for stress tolerance, which needs to be identified and characterized. The close phylogenetic relationships between millets and other cereals could enable the introgression of novel alleles, genes or QTL identified in millets for better agronomic traits into other cereals towards ensuring food security under changing climate.

### **Forest and fruit tree crops**

Clones of trees, namely populus, pinus, abies, and eucalyptus are used in afforestation, as they are dedicated to the production of wood and other wood-derived products. Therefore, it is imperative to develop climate-change resilient clones or populations of these forest trees. Several procedures have been developed for high-throughput DNA genotyping and genome-wide marker identification in forest trees. The genome complexity reduction DArT (Alves-Freitas et al., 2011) and whole-exome capture using in-solution target enrichment (Neves et al., 2011) have been tested successfully for genome-wide marker identification needed for GS in *Pinus taeda*. Considering the importance of genome sequence for development of genetic markers, Conifer Genome Project (<http://www.pinegenome.org/cgp/>) has been launched with an aim of promoting advance genome research in loblolly pine (*P. taeda*; 21.7 Gbp/1C;  $n = 12$ ), white pines (*Pinus* subgenus *strobus*; 25.1 Gbp/1C;  $n = 12$ ), as well as *Sequoia sempervirens* (31.4 Gbp/1C;  $n = 3x = 33$ ) and Douglas-fir (*Pseudotsuga menziesii*; 18.6 Gbp/1C;  $n = 13$ ). An extensive genetic resources and gene catalog was developed for *P. taeda* and *Picea glauca* (white spruce; 19.7 Gbp/1C;  $n = 12$ ) (<http://www.pinegenome.org/cgp/>). The GENOAK project (<http://urgi.versailles.inra.fr/Projects/GenOak>) aims to establish a high quality reference genome sequence for pedunculate oak (*Quercus robur*; 905 Mbp/1C;  $n = 12$ ). The *Eucalyptus grandis* (640 Mbp/1C,  $n = 11$ ) genome has been deciphered ([http://phytozome.jgi.doe.gov/pz/portal.html#!info?alias=Org\\_Egrandis](http://phytozome.jgi.doe.gov/pz/portal.html#!info?alias=Org_Egrandis)) and will benefit agro-foresters utilizing this fast-growing hardwood tree to support industries based on Eucalypt fibre and hardwood products, and the production of Eucalypt feedstock for cellulosic biofuels.

Importantly, this would assist in accelerating forest tree breeding for fast response to the need of adapted populations facing environmental modifications induced by climate change.

Fruit trees are also as important as the pulse and cereal crops and climate-change resilient clones or populations of fruit tree crops are necessary to maintain the source of nutrients that help the daily intake of healthy food ingredients. Genomics-based breeding approaches, along with bioinformatics capability and other omics resources will be the essential components of perennial fruit crop breeding and particularly, to adapt their cropping to combat or mitigate climate change effects. Genome sequencing and annotation projects include perennial fruit crops such as apple (Velasco et al., 2010), banana (Velasco et al., 2007), cacao (D'Hont et al., 2012), grape (Argout et al., 2011), peach (Ahmed et al., 1992) and sweet orange (Xu et al., 2013). The advances in genome sequencing, along with high-resolution genetic mapping and precise phenotyping will accelerate the discovery of functional alleles and allelic variations that are associated with traits of interest for perennial fruit crop breeding. However, very less progress has been made in this aspect and particularly, enhancing climate resilience needs more attention. For achieving this, genetics and genomics methodologies could provide the toolbox for identifying genomic regions associated with the desired phenotype, and assist the selection from the wild genetic resources of the parental plants that will be intercrossed to provide the progenies for commencing breeding procedures for recurrent selection.

### **Genomics-assisted breeding strategies for climate resilient traits**

Genomics-based approaches and NGS have ushered in sequence-based breeding strategies that will expedite the dissection and cloning of the loci controlling abiotic stress tolerance, while providing unparalleled opportunities to tap into wild relatives of crops, hence expanding the reservoir of genetic diversity available to breeders (Tuberosa et al., 2011; Edwards et al., 2013) (Figure 1). In view of the complexity and low heritability of yield, particularly under drought and other abiotic stresses, GS will provide the most powerful approach to raise the yield potential to the levels required to meet the fast-increasing global demand in cereal grain. However, MAS will remain a valid option for major loci or QTL, while QTL cloning will become a more routine activity facilitated through a more widespread utilization of high-throughput, accurate phenotyping (Araus et al., 2014), sequencing (Imelfort et al., 2009; Edwards et al., 2012), and identification of suitable candidate genes through 'omics' profiling (Gupta et al., 2013). Cloned QTL will provide novel opportunities for genetic engineering for abiotic stress tolerance and for a more targeted search for novel alleles in wild germplasm (Salvi et al., 2007). Even with the application of advanced genomics technologies, mitigating the negative effects of climate change on crop productivity will remain a daunting undertaking. This requires a multidisciplinary and integrated approach, which will eventually allow plant breeders to more effectively select crops that are more resilient to climate change and ensure a sufficient level of food security for the decades to come.

### **Flowering time and drought adaptation**

Temperature influences crop development in concert with additional floral pathways such as day-length (photoperiod), which collectively control floral transition through interconnected genetic pathways. Global warming will result in increased ambient temperature with unchanged

photoperiods at given latitudes. Annual plants generally respond to increased temperatures with accelerated growth and development, having shortened lifecycles, less opportunity for photosynthesis (Reynolds et al., 2010), a shorter reproductive phase and lower yield potential (Ainsworth and Ort, 2010). There is also an increased risk of damage to reproductive tissue caused by the coincidence of high temperatures and sensitive developmental stages. Therefore, detailed knowledge of the interplay between genetic control of flowering, allelic variants, epistatic interactions and phenotypic variations in varied growth conditions is necessary in order to identify breeding targets for climate change scenarios.

There are increasing number of germplasm resources including precise near isogenic lines (NILs) (Bentley et al., 2011; Bentley et al., 2013) as well as next-generation populations such as multi-founder populations (e.g., multi-parent advanced generation intercross populations), which have been developed in wheat (Mackay et al., 2014) and other crops to facilitate further research and validation of climate-smart crops. New variation incorporated into elite backgrounds from landraces, ancestral or wild crop relatives (e.g., [www.wheatisp.org](http://www.wheatisp.org)) also offers potential for discovery of functional variation for manipulating flowering time to suit future climate permutations. However, initial work should focus on understanding the effect of flowering time on yield potential across environments and environmental stresses. Identifying the potential utility of loci of minor effect and/or which affect various stages of reproductive development could offer the ability to shorten or lengthen various phases of the flowering process, thereby enabling fine-tuning of flowering to suit particular regional climatic conditions, and to adapt to any changes in these conditions.

In case of drought tolerance, multi-disciplinary research is underway to improve plants' response to drought and water-use efficiency. With the advent of molecular breeding, QTL identification and QTL use in breeding programs assist in developing new cultivars with improved drought tolerance. In maize, extensive work has been carried out to investigate the role of root in mitigating the negative effects of drought. QTL for root traits have been described in a number of maize populations (Ruta et al., 2010; Tuberosa et al., 2011; Hund et al., 2011) in which some QTL showed a concurrent effect on grain yield performance under drought (Landi et al., 2005). Recently, Syngenta and Pioneer-DuPont deployed proprietary genomics-assisted approaches to select drought-tolerant maize hybrids (Agrisure Artesian™ and AQUAmax™, respectively) (Cooper et al., 2014). The superior performance of these maize hybrids in the severe drought that plagued the US Corn Belt in summer 2012 underlines their validity under dry soil conditions (Cooper et al., 2014). In wheat, yield QTL were shown to be associated with components of other traits, supporting the prospects for dissecting crop performance under abiotic stress conditions into its physiological and genetic components in order to facilitate a strategic approach to breeding (Reynolds et al., 2008). At least 15 different populations have been used to map drought adaptation in rice and four regions were identified as key for yield or yield components under stress, and drought-tolerant component traits were identified across populations with interval lengths of 35-64 cM (Kamoshita et al., 2008). The first region (on chromosome 1) was associated with grain yield drought-resistance traits, plant type traits (Zhang et al. 2001a), and QTL for cell-membrane stability (Tripathy et al., 2000) and osmotic adjustment (Lilley et al., 1992), and root traits (Robin et al., 2003). Second genomic region on chromosome 4 was rich in root trait QTL (Zheng et al. 2000; Hemamalini et al. 2000; Zhang et al. 2001b; Kamoshita et al. 2002; Nguyen et al. 2004; Boopathi et al. 2005) under well-watered

and drought conditions. The third region located on chromosome 8 contained QTL for plant water status, grain yield, cell membrane stability, osmotic adjustment, rate of non-stomatal water loss and deep and thick root traits (Zheng et al. 2000; Hemamalini et al. 2000; Zhang et al. 2001b; Nguyen et al. 2004; Boopathi et al. 2005). The fourth important region for drought was located in chromosome 9, which was characterized by QTL for root traits, cell membrane stability, plant water status, leaf rolling and leaf drying, biomass, number of grains per panicle, relative spikelet fertility and delay in flowering time (Hemamalini et al. 2000; Tripathy et al. 2000; Zhang et al. 2001a; Kamoshita et al. 2002; Price et al. 2002a, 2002b; Robin et al. 2003; Courtois et al. 2003; Babu et al. 2003; Zheng et al. 2003; Lafitte et al. 2004; Lanceras et al. 2004; Nguyen et al. 2004; Boopathi et al. 2005; Gomez et al. 2005; Jearakongman 2005; Li et al. 2005; Xu et al. 2005; Yue et al. 2006). Since these four regions are consistently reported to be associated with drought response and stood above the average, these regions should be part of marker assisted breeding program for drought tolerance in rice.

### **Cold and heat stress tolerance**

Tolerance to freezing temperatures is the most important component for winter survival, but also of considerable importance is the capability to withstand combinations of stresses due to desiccation, wind, ice-encasement, heaving, low light, snow cover, winter pathogens, and fluctuating temperatures. Resistance to desiccation through the maintenance of cell membrane integrity and retention of cellular water is essential, and it is unsurprising that the same genetic response to the onset of freezing temperatures is often observed with drought or salinity stress (Yue et al., 2006). Indeed, cold acclimation (CA) can frequently improve adaptation to a mild drought stress and *vice versa* (Seki et al., 2002). Major genes or gene clusters involved in the control of frost and drought adaptation are located on a region of the long arm of Triticeae group 5 chromosomes. Traits such as winter hardiness (Thomas et al., 1993), vernalization response and frost tolerance (Hayes et al., 1993; Galiba et al., 1995), cold- and drought-induced ABA production (Laurie et al., 1995), and osmotic stress-tolerance (Galiba et al., 1993), have all been mapped to this region. Across the grasses and cereals, this chromosome region has been a major focus for genome research and for plant breeding. It may well be as consequence of climate change from the perspective of future crop design that in many locations where winter temperatures are on the increase and favoring continued plant growth, and where this is accompanied by a decrease in winter rainfall, that unseasonal winter droughts will ensue, which will require a new breeding strategy for common stress tolerance to both stress factors.

The C-repeat binding factor (CBF) genes are key regulators of the expression of COR (cold regulated genes), which are conserved among diverse plant lineages such as eudicots and monocots. The CBF transcription factors recognize the cis-acting CRT/DRE (C-repeat/dehydration responsive element) element in the regulatory regions of COR genes (Stockinger et al., 1997). Twenty CBF genes have been identified in barley (*H. vulgare*), of which 11 are found in two tight tandem clusters on the long arm of chromosome 5H in the same region as the *Fr-H2* frost tolerance locus (Skinner et al., 2006; Francia et al., 2007). An orthologous genomic region in *T. monococcum* contains similar CBF gene clusters located at the *Fr-A<sup>m</sup>2* frost tolerance QTL (Miller et al., 2006; Vagujfalvi et al., 2003). Studies of the organization of CBF cluster in barley and wheat have shown that the number of CBF genes at *Fr-H2/Fr-A1* locus may vary among cultivars with winter forms having a higher copy number of



some CBFs (Francia et al., 2007; Knox et al., 2010). The co-segregation of CBF gene clusters with barley *Fr-H2* and wheat *Fr-A<sup>m</sup>2* frost tolerance loci, their role in cold acclimation (Stockinger et al., 1997), and the association of transcript levels of CBF genes with frost tolerance loci (Vagujfalvi et al., 2003) make them obvious candidates for one of the two major frost tolerance QTL on Triticeae group 5 chromosomes. The locations of two frost tolerance/winter survival QTL on the chromosome 5F of forage grass *Festuca pratensis* correspond most likely to the *Fr-A1* and *Fr-A2* loci on wheat homoeologous group 5A chromosomes. One of these QTL (*QFt5F-2/QWs5F-1*) has *FpCBF6* as a candidate gene shown to be rapidly up-regulated during CA (Alm et al., 2011).

Conversely, many crops are currently grown in places, where high temperature prevails and field studies have indicated that increase in temperature reduces grain yield of cereals and legumes by 4 to 14% per 1°C increase (Quarrie et al., 1997). Current projections indicate that both day and night temperatures are likely to increase during this century (Hall et al., 2000) and ideally, heat-resistant cultivars should not only have higher grain yields in hot environments but also similar grain yields as current cultivars in cool atmosphere. Public plant breeding programs have developed heat-resistant cultivars of cowpea, common bean, tomato and Pima cotton that are more productive in hot environments than standard cultivars. Commercial plant breeding companies rarely divulge their methods, but from the available heat-resistant commercial cultivars, it is clear that they have had some success in breeding for heat tolerance during reproductive development in tomato and upland cotton. In the past, very few public or commercial plant-breeding programs gave any emphasis to breeding heat-resistant cultivars. For crops that are sensitive to high temperatures during reproductive development the way forward is to give great emphasis to breeding and finding DNA markers for heat adaptation during flowering.

### **Submergence and salinity tolerance**

Waterlogging is a major problem for cereal production worldwide, as in sodic environments, soils are affected by seepage from irrigation canals, and excess wetting due to rainfall or floods, especially if it rains after irrigation. Genetic diversity in waterlogging tolerance was reported in various crops, including wheat, barley, maize and oats (Kerr, 1986), and diverse mechanisms have been associated with tolerance. They are associated with phenology and morphology, nutrition balances, metabolism, including anaerobic catabolism and anoxia tolerance, and post-anoxia damage and recovery (Setter et al., 2003). Tolerance of flooding during germination and early seedling growth is essential for direct seeding of rice, both in rainfed and irrigated areas, where even waterlogging is sufficient to cause considerable reduction in crop stand because of their high sensitivity to hypoxia at this stage (Setter et al., 2003). Substantial genetic variation was recently observed in the ability to germinate and establish in flooded soil. Tolerant genotypes are capable of catabolizing starch reserves in seeds germinating under hypoxia into simple sugars, and use them as substrates to generate energy via anaerobic pathways for the growing embryos (Miro et al., 2013; Septiningsih et al., 2013). Several QTL originating from a few rice landraces were identified, two of them with large effects; on chromosome 9 (*qAG-9-2*) (Setter et al., 2003) and chromosome 7 (*qAG-7-1*) (Septiningsih et al., 2013). These QTL are being targeted for cloning and for use through MAB, which could eventually result high yielding rice cultivars for deep-water areas. Recently, tolerant rice varieties carrying *SUB1* locus became

505 available. *SUB1* is a major QTL on chromosome 9 that has been cloned and the gene responsible  
506 for tolerance identified as *SUB1A-1*. This gene encodes an ERF that suppresses ethylene-  
507 mediated responses under submergence, and subsequently limits excessive elongation and halts  
508 chlorophyll degradation. Both processes are essential to prevent carbohydrate starvation of the  
509 submerged plants. These varieties can survive 4 to 18 days of complete submergence, with yield  
510 benefits of 1 to over 3.5 t ha<sup>-1</sup> (depending on flood duration and floodwater condition), compared  
511 to current farmers' varieties, and without any undesirable effects on the features of the original  
512 varieties (Singh et al., 2009; Bailey-Serres et al., 2010; Ismail et al., 2013; Mackill et al., 2012).  
513 Additional genes are being targeted for submergence tolerance, and once identified they could be  
514 combined with *SUB1* for higher tolerance during germination and stagnant flooding. Further, the  
515 progress made in rice could potentially be exploited to improve flood tolerance of other crop  
516 species and provide more resilient varieties for current and future flood-affected areas.

517  
518 Progressive salt accumulation due to excessive irrigation with poor water quality coupled with  
519 poor or improper drainage results in high salt levels (Tuberosa, 2012). Numerous studies have  
520 characterized responses mediated by salt stress in different plant species and highlighted the  
521 complexity of the mechanisms involved (Munns et al., 2008). Studies have shown that few major  
522 loci and many minor ones were associated with various aspects of salinity tolerance. The best  
523 known for rice is *Saltol* on chromosome 1 (Thomson et al., 2010), which possesses a major gene,  
524 *OsHKT1;5* (Ren et al., 2005). In wheat, two members of *HKT* gene family (including the wheat  
525 *HKT1;5* orthologue) have also been shown to co-localize with major QTL (Byrt et al., 2007).  
526 Apparently, many other QTL have been identified in rice and other cereals, and several of them  
527 are common across mapping populations. In addition, numerous genes have been identified  
528 through functional genomics studies of salt-stress responses, and many of them lead to improved  
529 tolerance when they are over- or under-expressed. Some even co-localize with QTL regions, but  
530 there has been no further success in using them for breeding tolerant cereal crops or in cloning  
531 additional QTL.

532  
533 Current approaches in this aspect involves using NGS to target major QTL for cloning, and to  
534 develop efficient SNP and InDel marker systems to manipulate these loci during MAB. The  
535 substantial genetic diversity in the tolerance of salt stress and mechanisms used by various crops  
536 to cope with increasing salt concentrations in soil and water provides opportunities to enhance  
537 salt-stress tolerance in cereals. However, this will require large investments to dissect and  
538 combine the genetic determinants of various traits. Developing such cultivars that are highly  
539 tolerant of salt stress is a requisite to cope with the current worsening climatic conditions and to  
540 meet the urgent need of producing more food from marginal land and limited water resources.

#### 541 542 **Host plant resistance to pathogens and pests**

543  
544 The climatic variables including changes in temperature, rainfall and atmospheric chemical  
545 composition along with predominantly elevated CO<sub>2</sub> levels would accelerate the reproduction  
546 time of many plant pathogens and pests, thereby increasing their infection pressure on crop  
547 plants (Boonekamp, 2012). Climate change may also affect the ability of plants to express  
548 resistance to pathogens and insects. Experiments conducted by Huang et al. (2009) indicated a  
549 45% increase in leaf lesions in oilseed rape, when the surrounding temperature was increased by  
550 5°C. This finding suggests that the expression and efficacy of R-genes in host plants may be

affected where both crop and associated pathogen or pest are affected by climatic variation. This may be influenced by different combinations of selective pressures, and each may respond to these pressures at different rates. Improved understanding on the host-pest/pathogen interactions and knowledge on different effects of climate change is a requisite for the development of climate-resilient crops. To date, research on the impact of climate change on plant diseases has been limited, with many studies focusing on the effects of a single atmospheric constituent or meteorological variable on the host, pathogen, or the interaction of the two, under controlled conditions. Whilst this work is a valuable base to start from, the combined effects of biotic and abiotic stresses must be studied (Ramegowda and Senthil-Kumar, 2015).

Recent advances in genome sequencing and genotyping assays allow for many strategies at the genomics level, which can be developed to understand the impact of climate change on plant diseases. The newly available genome sequences for plants, pathogens and pests provide the resources to study their co-evolution in response to climate change. An understanding of the co-evolution of genes responsible for virulence and resistance will lead to improved plant protection strategies and provide a model to understand plant-pathogen and plant-insect interactions in diverse species. Though it is important to understand the genomics of disease resistance in crop species, and how allelic differences are altering resistance, combining this with studies of CWR or germplasm collections further allows the identification of novel variants. These variants can be used for the introgression of novel resistance genes into cultivars, utilizing the germplasm for breeding and developing new cultivars, or genetic engineering with the advantageous genes. Taken together, it is obvious that the impact of climate change on disease resistance is difficult to predict and is likely to be variable depending on the crop and local environment. However, crop disease is an important factor when considering the impact of climate change on food production and intensive studies applying advanced genomics tools will be required to help ameliorate the impact of climate change on future cropping scenarios in relation to plant disease.

### **Genomic engineering tools for targeted mutagenesis by editing genes for adaptation**

Plant breeders have been applying mutagenesis to induce genetic variation for increasing crop yield and later, the strategy has been used for improving the adaptability of crop plants. Initially, X-ray radiation was used as a mutagen since it was readily available to researchers (Muller, 1927). Subsequently, gamma-ray radiation has been used to induce point mutations, although chromosomal mutation were also produced (Devreux and Mugnozza, 1964). From recent times, chemical mutagenesis is being practiced since they are easy to use, do not require any specialised equipment, and can provide a very high mutation frequency. Compared to radiation methods, chemical mutagens tend to induce SNPs rather than chromosomal mutations. Currently, chemical mutagens, such as Ethyl methanesulfonate (EMS) are being used to induce random mutations into the genome and have become a useful complement to the isolation of nuclear DNA from mutated lines by TILLING (Targeting Induced Local Lesions in Genomes) technology and screening of the M<sub>2</sub> population at the DNA level using advanced molecular techniques. Single mutations in specific genes for adaptation could be identified by cleavage of mismatched bases formed as a result of repeated melting and reannealing of PCR products amplified from a pool of alleles for the specific gene in a pool of DNA from a set (usually 8) of M<sub>2</sub> plants (McCallum et al., 2000; Caldwell et al., 2004). NGS can efficiently accelerate the identification of mutations at the whole-genome level. Promotor mutations and mutation in other regulatory elements

responsible for the downstream effect can be identified by qPCR, microarray and RNA-seq. Once a mutant allele is identified within gene(s) of interest, those mutations may be linked to a specific phenotype for stress resistance by backcrossing the mutant to the parental line. This TILLING approach is a reverse genetics procedure to associate a mutant allele to its phenotype. Of note, TILLING can also be used for a forward genetic approach by screening phenotypes for adaptation to stresses and then characterize the phenotype using a combination of whole-genome resequencing, linkage maps and microarrays, to gain a broad picture of gene expression changes due to the newly introduced SNPs compared to the original line.

Other molecular tools and resources are now available for genome engineering and reverse genetics experiments in crop plants in order to implement precise manipulation of genetic building blocks and regulatory machinery that underlie yield improvement under stress condition and directly correct harmful mutations by genome editing (Hsu et al., 2014). Targeted genome engineering has emerged as an alternative to classical plant breeding and transgenic methods to improve crop plants and ensure sustainable food production (Belhaj et al., 2013; Osakabe and Osakabe, 2015). Currently, four types of engineered nucleases are used for genome editing: engineered homing endonucleases/meganucleases (EMNs) (Silva et al., 2011), zinc finger nucleases (ZFNs) (Townsend et al., 2009), transcription activator-like effector nucleases (TALENs) (Cermak et al., 2011), and CRISPR (clustered regularly interspaced short palindromic repeats)/Cas (CRISPR-associated)9 (Cong et al 2013; Mali et al 2013).

Sequence-specific nucleases (SSN) enable precise genome engineering by introducing DNA double-strand breaks (DSB) that subsequently trigger endogenous DNA repair by non-homologous end joining (NHEJ) or homology directed repair (HDR) recombination mechanisms in different species. Site-directed mutagenesis mediated via NHEJ can be achieved while HDR cause directed gene knock-in/correction at specific locations in the genome (HDR uses a DNA template to replace the DNA sequence at the break point). NHEJ functions throughout the entire cell cycle whereas HR is restricted to late S/G2 phases in the cell cycle. Therefore, NHEJ is the major DSB repair pathway in eukaryotes. Belhaj et al. (2013) and Osakabe and Osakabe (2015) display a clear illustration of genome editing assays in model (*Arabidopsis thaliana* and *Nicotiana benthamiana*) and crop (*Oryza sativa*, *Triticum aestivum* and *Sorghum bicolor*) plant species. These SSN effects generate targeted genome modifications including mutations, insertions, replacements and chromosome rearrangements and have been induced in a variety of important crops, such as rice, maize, wheat, barley and soybean. Each technology has advantages and disadvantages with regard to cost, ease of construction, efficiency of targeting, and specificity (Chen and Gao, 2014; Gao, 2015). Major advantages of ZFNs are related to the acceptance of the technology as no transgenic is produced because viral vectors have been used for expressing transiently the nuclease, which do not integrate into the genome. However, it has disadvantages such as difficulties to design the experiments, limited number of target sites, and the regeneration of juvenile and chimeric mutated plants when custom-designed nucleases have been delivered in tree explants.

CRISPR was first discovered as an immune system of prokaryotes, which subsequently became a powerful tool for genome editing in eukaryotes (Gaj et al., 2013). It has emerged as an alternative to classical plant breeding and transgenic methods to improve crop plants. Plant transformation and co-expression of the Cas9 with a chimeric guide-RNA (gRNA) targeting a

GN19NGG motif in the gene of interest, results in a double-strand non-self DNA cleavage on both strands at a specific site near the protospacer adjacent motif (PAM) (Jinek et al., 2012; Gasiunas et al., 2012). The Cas9 from *Streptococcus pyogenes* (SpCas9) recognizes 5'-NGG-3' as the PAM sequence. PAM plays an important role in target binding and cleavage by the Cas9-gRNA complex. CRISPR/Cas9 has greater number of advantages, including the straightforward construct design and assembly and the achievement of high mutation rates, matching or exceeding those obtained with ZFNs and TALENs. Only 20 nucleotides in the gRNA need to be modified to recognize a different target making unnecessary the sophisticated protein engineering for each target that is crucial for the other SSN approaches.

So far, the CRISPR-Cas9 technology has been applied in *A. thaliana* (Feng et al., 2013, 2014; Jiang et al., 2013; Li et al., 2013; Mao et al., 2013) *Nicotiana benthamiana* (Jiang et al., 2013; Nekrasov et al., 2013; Li et al., 2013), *Oryza sativa* (Feng et al., 2013, Jiang et al., 2013, Mao et al., 2013, Shan et al., 2013; Feng et al., 2013; Xie and Yang, 2013; Miao et al., 2013; Zhang et al., 2014), *Solanum lycopersicum* (Brooks et al., 2014), *Sorghum bicolor* (Jiang et al., 2013), *Triticum aestivum* (Wang et al., 2014), *Citrus sinensis* (Jia and Wang, 2014) and *Populus tremula x alba* (Zhou et al., 2015). Genes controlling traits of importance for adaptation have also been edited by CRISPR-Cas9 technology. gRNAs were designed to target three specific sites of the rice *OsMPK5* gene which encodes a stress-responsive rice mitogen-activated protein kinase and the targeted mutation of *OsMPK5* enhanced rice disease resistance (Xie and Yang, 2013). Transgenic wheat plants carrying mutations in *TaMLO-A1* allele were susceptible to powdery mildew diseases (Wang et al., 2014). The bacterial blight susceptibility genes, *OsSWEET14* and *OsSWEET11*, were targeted for mutation at the promoter region in Arabidopsis, tobacco, sorghum and rice (Jiang et al., 2013). High CRISPR-Cas9 mutational efficiency was achieved for three 4-coumarate:CoA ligase (4CL) genes, *4CL1*, *4CL2* and *4CL5*, associated with lignin and flavonoid biosynthesis in *Populus tremula x alba* (Zhou et al., 2015).

Moreover, accelerated breeding of crop plants carrying targeted gene mutation(s) without foreign DNA is possible using CRISPR genome editing. In fact, although transgene Cas9 and selectable marker integration is hemizygous, CRISPR editing at the target loci is biallelic. Therefore, in autogamous plants, self-fertilization of T<sub>1</sub> plants will provide 25% of the T<sub>2</sub> plants without the transgene but maintaining the edited gene in homozygosity. In self-incompatible or dioecious perennial woody trees, biparental hemizygous Cas9/sgRNA transformation and biallelic-edited gene can be produced. Controlled crosses between male and female primary transformants with confirmed biallelic mutations should produce transgene-free, biallelic mutants in 25% of the progeny (Zhou et al., 2015). Taken together, genome engineering for targeted mutagenesis by editing genes serves as a potential strategy for generating elite cultivars of crop plants with durable climate resilience.

## Concluding remarks

It is realized that the global climate change is going to impose a severe threat on agricultural productivity worldwide, and thereby challenging food security and nutritional security. Advances in technology, particularly transgene-based and molecular breeding technologies have facilitated the development of elite genotypes with durable adaptation to climate change. Noteworthy, crossbreeding coupled with genomics forms genomics-assisted breeding, which is playing a

significant role for developing climate change resilient crops. Excellent model organisms for climate change such as foxtail millet and green foxtail (for C<sub>4</sub> photosynthesis), *Brachypodium* (grass model) have been identified for deciphering traits that need to be decoded and introgressed in the crop plants. Advances in DNA sequencing technologies and the sequencing of CWR, along with advanced genomics tools will expedite the identification of novel genes and key regulatory regions of stress tolerance towards the development of new cultivars with durable resistance. Although, the impact of climate change on crop's resistance is difficult to predict and is likely to be variable depending on the crop and environment, genomics-assisted breeding could contribute significantly to reduce the impact of climate change on future cropping scenarios.

### Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

### Author contributions

C.K. conceived and outlined the review; M.M., R.H., D.E., R.S., M.A., J.B., A.B., M.B., J.B., H.C., M.C., L.J.C., J.C., A.C.O., C.D.P., H.D., S.E., P.G., A.G., A.H., K.H., G.T.H., S.H., M.W.H., M.I., A.M.I., A.M., S.M., H.T.N., F.C.O., R.O., A.H.P., P.W.S., J.T., R.T., B.V., R.V., S.D.W., M.Y., M.P. wrote the manuscript. M.M., M.P. edited the manuscript and prepared the final version. M.P. and C.K approved the final version.

### Acknowledgements

This manuscript was researched and composed by the members of International Climate-Resilient Crop Genomics Consortium (ICRCGC). The white paper authored by the members is available at [http://www.climatechange-genomics.org/climate\\_change\\_version\\_1.1.pdf](http://www.climatechange-genomics.org/climate_change_version_1.1.pdf) and [http://www.icrcgc.org/ncc\\_manuscript.docx](http://www.icrcgc.org/ncc_manuscript.docx).

### References

- Ahmed, F. E., Hall, A. E., DeMason, D. A. (1992). Heat injury during floral development in cowpea (*Vigna unguiculata*, Fabaceae). *Am. J. Bot.* 79, 784-791.
- Ainsworth, E. A., and Ort, D. R. (2010). How do we improve crop production in a warming world? *Plant physiol.* 154, 526-530
- Alm, V., Busso, C. S., Ergon, A., Rudi, H., Larsen, A., Humphreys, M.W., and Rognli, O.A. (2011). QTL analyses and comparative genetic mapping of frost tolerance, winter survival and drought tolerance in meadow fescue (*Festuca pratensis* huds.). *Theor. Appl. Genet.* 123, 369-382.
- Alves-Freitas, D., Kilian, A., Grattapaglia, D. (2011). Development of DArT (Diversity Arrays Technology) for high-throughput genotyping of *Pinus taeda* and closely related species. *BMC Proceedings* 5, 22.

- Anbessa, Y., Taran, B., Warkentin, T. D., Tullu, A., and Vandenberg, A. (2009). Genetic analyses and conservation of QTL for ascochyta blight resistance in chickpea (*Cicer arietinum* L.). *Theor. Appl. Genet* 119, 757-765.
- Anuradha, C., Gaur, P. M., Pande, S., Gali, K. K., Ganesh, M., Kumar, J., and Varshney, R. K. (2011). Mapping QTL for resistance to botrytis grey mould in chickpea. *Euphytica* 182, 1-9.
- Araus, J. L., Cairns, J. E. (2014). Field high-throughput phenotyping: The new crop breeding frontier. *Trends Plant Sci.* 19, 52-61.
- Argout, X., Salse, J., Aury, J. M., Gaultier, M. J., Droc, G., Gouzy, J., et al. (2011). The genome of *Theobroma cacao*. *Nat. Genet.* 43, 101-108.
- Asfaw, A., Blair, M. W., Struik, P. C. (2012). Multi-environment quantitative trait loci analysis for photosynthate acquisition, accumulation, and remobilization traits in common bean under drought stress. *G3 (Bethesda)* 2, 579-595.
- Babu, R. C., Sarkarung, S., Wade, L. J., Nguyen, H. T., Nguyen, B. D., Chamarerk, V., et al. (2003). Genetic analysis of drought resistance in rice by molecular markers: Association between secondary traits and field performance. *Crop Sci.* 43, 1457-1469.
- Bailey-Serres, J., Fukao, T., Ismail, A., Heuer, S., Mackill, D. (2010). Submergence tolerant rice: SUB1's journey from landrace to modern cultivar. *Rice* 3, 138-144.
- Bailey-Serres, J., Fukao, T., Ronald P., Ismail A., Heuer, S., and Mackill, D. (2010). Submergence tolerant rice: *Sub1's* journey from landrace to modern cultivar. *Rice* 3, 138-147.
- Bajaj, D., Saxena, M. S., Kujur, A., Das, S., Badoni, S., Tripathi, S., et al. (2014). Genome-wide conserved non-coding microsatellite (CNMS) marker-based integrative genetical genomics for quantitative dissection of seed weight in chickpea. *J. Exp. Bot.* 66, 1271-1290.
- Bajaj, D., Upadhyaya, H. D., Khan, Y., Das S., Badoni S., Shree T., et al. (2015). A combinatorial approach of comprehensive QTL-based comparative genome mapping and transcript profiling identified a seed weight-regulating candidate gene in chickpea. *Sci. Rep.* 5, 9264.
- Beebe, S., Rao, I. M., Cajiao, C., Grajales, M. (2008). Selection for drought resistance in common bean also improves yield in phosphorus limited and favorable environments. *Crop Sci.* 48, 582-592.
- Belhaj, K., Chaparro-Garcia, A., Kamoun, S., and Nekrasov, V. (2013). Plant genome editing made easy: targeted mutagenesis in model and crop plants using the CRISPR/Cas system. *Plant Methods* 9, 39.
- Bennetzen, J. L., Schmutz, J., Wang, H., Percifield, R., Hawkins, J., Pontaroli, A. C., et al. (2012). Reference genome sequence of the model plant *Setaria*. *Nature Biotechnol.* 30, 555-561.

- Bentley, A. R., Horsnell, R., Werner, C. P., Turner, A. S., Rose, G. A., Bedard, C., et al. (2013). Short, natural, and extended photoperiod response in BC<sub>2</sub>F<sub>4</sub> lines of bread wheat with different *Photoperiod-1* (*Ppd-1*) alleles. *J. Exp. Bot.* 64, 1783-1793.
- Bentley, A. R., Turner, A. S., Gosman, N., Leigh, F. J., Maccaferri, M., Dreisigacker, S., et al. (2011). Frequency of photoperiod-insensitive *Ppd-A1a* alleles in tetraploid, hexaploid and synthetic hexaploid wheat germplasm. *Plant. Breed.* 130, 10-15.
- Bidinger, F. R., Napolean, T., Hash, C. T. (2007). Quantitative trait loci for grain yield in pearl millet under variable post flowering moisture conditions. *Crop Sci.* 47, 969-980.
- Bohra, A., Dubey, A., Saxena, R. K., Penmetsa, R. V., Poornima, K. N., Kumar, N., et al. (2011). Analysis of BAC-end sequences (BESs) and development of BES-SSR markers for genetic mapping and hybrid purity assessment in pigeonpea (*Cajanus* spp.). *BMC Plant Biol.* 11, 56.
- Boonekamp, P. M. (2012). Are plant diseases too much ignored in the climate change debate? *Eur. J. Plant Pathol.* 133, 291-294.
- Boopathi, N. M., Chezhaian, P., Jeyaprakash, P., Satheesh Kumar, S., Gomez, M., Suresh, R., et al. (2005). QTL mapping of drought-resistance traits using indica rice (*Oryza sativa* L.) lines adapted to target population of environment. In: *Proceedings of the 2nd International Conference on Integrated Approaches to Sustain and Improve Plant Production under Drought Stress*, Rome, Italy.
- Borrell, A. K., Mullet, J. E., George-Jaeggli, B., van Oosterom, E. J., Hammer, G. L., Klein, P. E., et al. (2014). Drought adaptation of stay-green sorghum is associated with canopy development, leaf anatomy, root growth, and water uptake. *J. Exp. Bot.* 65, 6251-63.
- Brooks, C., Nekrasov, V., Lippman, Z. B., and Van Eck, J. (2014). Efficient Gene Editing in Tomato in the First Generation Using the Clustered Regularly Interspaced Short Palindromic Repeats/CRISPR-Associated9 System. *Plant Physiol.* 166, 1292-1297.
- Byrt, C. S., Platten, J. D., Spielmeier, W., James, R. A., Lagudah, E. S., Dennis, E. S., et al. (2007). HKT1;5-like cation transporters linked to Na<sup>+</sup> exclusion loci in wheat, *Nax2* and *Kna1*. *Plant Physiol.* 143, 1918-1928.
- Caldwell, D. G., McCallum, N., Shaw, P., Muehlbauer, G. J., Marshall, D. F., and Waugh, R. (2004). A structured mutant population for forward and reverse genetics in Barley (*Hordeum vulgare* L.). *Plant J.* 40, 143-50.
- Cermak, T., Doyle, E. L., Christian, M., Wang, L., Zhang, Y., Schmidt, C., et al. (2011). Efficient design and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting. *Nucl. Acids Res.* 39, e82.
- Chen, K., and Gao, C. (2014). Targeted genome modification technologies and their applications in crop improvements *Plant Cell Rep.* 33, 575-583.



Cobos, M. J., Winter, P., Kharrat, M., Cubero, J. I., Gil, J., Milian, T., and Rubio, J. (2009). Genetic analysis of agronomic traits in a wide cross of chickpea. *Field Crops Res.* 111, 130-136.

Cong, L., Ran, F. A., Cox, D., Lin, S., Barretto, R., Habib, N., et al. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819-823.

Cooper, M., Gho, C., Leafgren, R., Tang, T., and Messina, C. (2014). Breeding drought-tolerant maize hybrids for the us corn-belt: Discovery to product. *J. Exp. Bot.* 65, 6191-6204.

Courtois, B., Shen, L., Petalcorin, W., Carandang, S., Mauleon, R., and Li, Z. (2003). Locating QTLs controlling constitutive root traits in the rice population IAC 165 × Co39. *Euphytica* 134, 335-345.

Das, S., Upadhyaya, H. D., Bajaj, D., Kujur, A., Badoni, S., Laxmi., et al. (2015). Deploying QTL-seq for rapid delineation of a potential candidate gene underlying major trait-associated QTL in chickpea. *DNA Res.* 22, 193-203.

Devreux, M., and Mugnozza, G. T. S. (1964). Effects of gamma radiation of the gametes, zygote and proembryo in *Nicotiana Tabacum* L. *Rad. Botany* 4, 373-386.

D'Hont, A., Denoeud, F., Aury, J. M., Baurens, F. C., Carreel, F., Garsmeur, O., et al. (2012). The banana (*Musa acuminata*) genome and the evolution of monocotyledonous plants. *Nature* 21, 213-217.

Dubey, A., Farmer, A., Schlueter, J., Cannon, S. B., Abernathy, B., Tuteja, R., et al. (2011). Defining the transcriptome assembly and its use for genome dynamics and transcriptome profiling studies in pigeonpea (*Cajanus cajan* L.). *DNA Res.* 18, 153-164.

Dutta, S., Kumawat, G., Singh, B., Gupta, D., Singh, S., Dogra, V., et al. (2011). Development of genic-SSR markers by deep transcriptome sequencing in pigeonpea [*Cajanus cajan* (L.) Millspaugh]. *BMC Plant Biol.* 11, 17.

Dwivedi, S. L., Sahrawat, K. L., Upadhyaya, H. D., Ortiz R. (2013). Food, nutrition and agrobiodiversity under global climate change. *Adv. Agron.* 120, 1-128.

Edwards, D., Batley, J., Snowdon, R. J. (2013). Accessing complex crop genomes with next-generation sequencing. *Theor. Appl. Genet.* 126, 1-11.

Edwards, D., Wang, X. (2012). "Genome sequencing initiatives" in *Genetics, Genomics and Breeding of Oilseed Brassicas*, eds. D. Edwards, J. Batley, I. A. P. Parkin, C. Kole (Enfield, NH: Science Publishers), 152-157.

Feng Z., Zhang B., Ding W., Liu X., Yang D-L., Wei P., et al. (2013). Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res.* 23, 1229-1232.

Feng, Z. Y., Mao, Y. F., Xu, N. F., Zhang, B. T., Wei, P. L., Wang, Z., et al. (2014). Multi-generation analysis reveals the inheritance, specificity and patterns of CRISPR/Cas induced gene modifications in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* 111, 4632-4637.

Fischer, R. A., Byerlee, D., Edmeades, G. O. (2009). *Can technology deliver on the yield challenge to 2050?* Rome: Food and Agriculture Organization of the United Nations.

Francia, E., Pecchioni, N., Barabaschi, D., Tondelli, A., Laidò, G., Rizza, F., et al. (2007). Fine mapping of a HvCBF gene cluster at the frost resistance locus Fr-H2 in barley.. *Theor. Appl. Genet.* 115, 1083-1091

Gaj, T., Gersbach, CA., and Barbas, CF. (2013). ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. *Trends Biotechnol.* 31, 397-405.

Galiba, G., Quarrie, S. A., Sutka, J., Morgounov, A., Snape, J. W. (1995). RFLP mapping of the vernalization (*VRN1*) and Frost-resistance (*FR1*) genes on chromosome 5A of wheat. *Theor. Appl. Genet.* 90, 1174-1179.

Galiba, G., Tuberosa, R., Kocsy, G., Sutka, J. (1993). Involvement of chromosomes 5A and 5D in coldinduced abscisic acid accumulation and in frost tolerance of wheat calli. *Plant breed.* 110, 237-242.

Gao, C. (2015). Developing genome editing technologies for crop improvement. In: *Meeting on Plant Genome Engineering*, Plant & Animal Genome XXIII, San Diego, CA, USA.

Gasiunas, G., Barrangou, R., Horvath, P., and Siksnys, V. (2012). Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. *Proc. Natl. Acad. Sci. U. S. A.* 109, E2579-E2586.

Gaur, R., Sethy, N. K., Choudhary, S., Shokeen, B., Gupta, V., and Bhatia, S. (2011). Advancing the stms genomic resources for defining new locations on the intraspecific genetic linkage map of chickpea (*Cicer arietinum* L.). *BMC Genomics* 12, 117.

Gautami, B., Foncéka, D., Pandey, M. K., Moretzsohn, M. C., Sujay, V., Qin, H., et al. (2012). An international reference consensus genetic map with 897 marker loci based on 11 mapping populations for tetraploid groundnut (*Arachis hypogaea* L.). *PLoS ONE* 7, e41213.

Gepts, P., Aragão, F. (2008). “Genomics of Phaseolus beans, a major source of dietary protein and micronutrients in the tropics,” in *Genomics of Tropical Crop Plants*, eds. P. Moore, R. Ming (Berlin: Springer), 113-143.

Gomez, M., Babu, R. C., Shanmugasundaram, P., Kumar, S. R., Suresh, R., Biji, K. R., et al. (2005). QTL mapping and marker assisted selection for drought tolerance in rice (*Oryza sativa* L.). in: *Proceedings of the 2nd International Conference on Integrated Approaches to Sustain and Improve Plant Production under Drought Stress* Rome, Italy

Gujaria, N., Kumar, A., Dauthal, P., Dubey, A., Hiremath, P., Prakash, A. B., et al. (2011). Development and use of genic molecular markers (GMMS) for construction of a transcript map of chickpea (*Cicer arietinum* L.). *Theor. Appl. Genet.* 122, 1577-1589.

Gupta, B., Sengupta, A., Saha, J., Gupta, K. (2013). Plant abiotic stress: ‘Omics’ approach. *J. Plant Biochem. Physiol.* 1, e108.

Hall, A. E., Ziska, L. H. (2000). "Crop breeding strategies for the 21st century," in *Climate Change and Global Crop Productivity*, eds. K. R. Reddy, H. F. Hodges (New York, USA: CAB International), 407-423.

Hayes, P. M., Blake, T., Chen, T. H., Tragoonrung, S., Chen, F., Pan, A., et al. (1998). Quantitative trait loci on barley (*Hordeum vulgare* L.) chromosome 7 associated with components of winter hardiness. *Genome* 36, 66-71.

Hemamalini, G. S., Shashidhar, H. E., and Hittalmani, S. (2000). Molecular marker assisted tagging of morphological and physiological traits under two contrasting moisture regimes at peak vegetative stage in rice (*Oryza sativa* L.). *Euphytica* 112, 69-78.

Hemamalini, G. S., Shashidhar, H. E., Hittalmani, S. (2000). Molecular marker assisted tagging of morphological and physiological traits under two contrasting moisture regimens at peak vegetative stage in rice (*Oryza sativa* L.). *Euphytica* 112, 69-78.

Hiremath, P. J., Kumar, A., Penmetsa, R. V., Farmer, A., Schlueter, J. A., Chamarthi, S. K., et al. (2012). Large-scale development of cost-effective snp marker assays for diversity assessment and genetic mapping in chickpea and comparative mapping in legumes. *Plant Biotechnol. J.* 10, 716-732.

Hisas, S. (2011). *The Food Gap. The Impacts of Climate Change in Food Production: A 2020 Perspective*. Alexandria, US: Universal Ecological Fund.

Hsu, P. D., Lander, E. S., and Zhang, F. (2014). Development and Applications of CRISPR-Cas9 for Genome Engineering. *Cell* 157, 1262-1278.

Hsu, P. D., Scott, D. A., Weinstein, J. A., Ran, F. A., Konermann, S., Agarwala, V., et al. (2013). DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat. Biotechnol.* 31, 827-832.

Huang, Y. J., Qi, A., King, G. J., Fitt, B. D. (2009). Quantitative resistance to symptomless growth of *Leptosphaeria maculans* (phoma stem canker) in *Brassica napus* (oilseed rape). *Plant Pathol.* 58, 314-323.

Hund, A., Reimer, R., Messmer, R. (2011). A consensus map of QTLs controlling the root length of maize. *Plant Soil* 344, 143-158.

Imelfort, M., Batley, J., Grimmond, S., Edwards, D. (2009). Genome sequencing approaches and successes. *Methods Mol. Biol.* 513, 345-358.

Ismail, A. M., Singh, U. S., Singh, S., Dar, M. H., and Mackill, D. J. (2013). The contribution of submergence-tolerant (*Sub1*) rice varieties to food security in flood-prone rainfed lowland areas in asia. *Field Crops Res.* 152, 83-93.

Jain, M., Misra, G., Patel, R. K., Priya, P., Jhanwar, S., Khan, A. W., et al. (2013). A draft genome sequence of the pulse crop chickpea (*Cicer arietinum* L.). *Plant J.* 74, 715-729.

Jearakongman, S. *Validation and discovery of quantitative trait loci for drought tolerance in backcross introgression lines in rice (Oryza sativa L.) cultivar IR64*. PhD Thesis Kasetsart University (2005).

- Jia, H., and Wang, N. (2014). Targeted Genome Editing of Sweet Orange Using Cas9/sgRNA. *PLoS ONE* 9, e93806.
- Jiang, W., Zhou, H., Bi, H., Fromm, M., Yang, B., and Weeks, D. P. (2013). Demonstration of CRISPR/Cas9/sgRNA-mediated targeted gene modification in Arabidopsis, tobacco, sorghum and rice. *Nucleic Acids Res.* 41, e188.
- Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816-821.
- Kamoshita, A., Chandra Babu, R., Boopathi, N. M., Fukai, S. (2008). Phenotypic and genotypic analysis of drought-resistance traits for development of rice cultivars adapted to rainfed environments. *Field Crops Res.* 109, 1-23.
- Kamoshita, A., Wade, L., Ali, M., Pathan, M., Zhang, J., Sarkarung, S., and Nguyen, H. (2002). Mapping QTLs for root morphology of a rice population adapted to rainfed lowland conditions. *Theor. Appl. Genet.* 104, 880-893.
- Kerr, RA. (1986). Greenhouse warming still coming. *Science* 232, 573-575.
- Khan, Y., Suresh, B. V., Muthamilarasan, M., Yadav, C. B., Prasad, M. (2014). Comprehensive genome-wide identification and expression profiling of foxtail millet [*Setaria italica* (L.)] miRNAs in response to abiotic stress and development of miRNA database. *Plant Cell Tiss. Organ Cult.* 118, 279-292.
- Khedikar, Y. P., Gowda, M. V., Sarvamangala, C., Patgar, K. V., Upadhyaya, H. D., Varshney, R. K. (2010). A QTL study on late leaf spot and rust revealed one major QTL for molecular breeding for rust resistance in groundnut (*Arachis hypogaea* L.). *Theor. Appl. Genet.* 121, 971-984.
- Knox, A. K., Dhillon, T., Cheng, H., Tondelli, A., Pecchioni, N., and Stockinger, E. J. (2010). Cbf gene copy number variation at frost resistance-2 is associated with levels of freezing tolerance in temperate-climate cereals. *Theor. Appl. Genet.* 121, 21-35.
- Kottapalli, P., Gaur, P. M., Katiyar, S. K., Crouch, J. H., Buhariwalla, H. K., Pande, S., and Gali, K. K. (2009). Mapping and validation of QTLs for resistance to an indian isolate of ascochyta blight pathogen in chickpea. *Euphytica* 165, 79-88.
- Kudapa, H., Bharti, A. K., Cannon, S. B., Farmer, A. D., Mulaosmanovic, B., Kramer, R., et al. (2012). A comprehensive transcriptome assembly of pigeonpea (*Cajanus cajan* L.) using sanger and second-generation sequencing platforms. *Mol. Plant* 5, 1020-1028.
- Kujur, A., Bajaj, D., Upadhyaya, H. D., Das, S., Ranjan, R., Shree, T., et al. (2015b). A genome-wide SNP scan accelerates trait-regulatory genomic loci identification in chickpea. *Sci. Rep.* 5, 11166.

949 Kujur, A., Upadhyaya, H. D., Shree, T., Bajaj, D., Das, S., Saxena, M., et al. (2015a). Ultra-  
950 high density intra-specific genetic linkage maps accelerate identification of functionally  
951 relevant molecular tags governing important agronomic traits in chickpea. *Sci. Rep.* 5, 9468.

952 Kulwal, P. L., Thudi, M., and Varshney, R. K. (2011). "Genomics interventions in crop  
953 breeding for sustainable agriculture," in *Encyclopedia of Sustainability Science and*  
954 *Technology*, ed. R. A. Meyers (New York, Springer), 2527-2540.

955 Kumari, K., Muthamilarasan, M., Misra, G., Gupta, S., Subramanian, A., Parida, S. K., et al.  
956 (2013). Development of eSSR-markers in *Setaria italica* and their applicability in studying  
957 genetic diversity, cross-transferability and comparative mapping in millet and non-millet  
958 species. *PLoS ONE* 8, e67742.

959 Lafitte, H. R., Price, A. H., and Courtois, B. (2004). Yield response to water deficit in an  
960 upland rice mapping population: Associations among traits and genetic markers. *Theor. Appl.*  
961 *Genet.* 109, 1237-1246.

962 Lanceras, J. C., Pantuwan, G., Jongdee, B., and Toojinda, T. (2004). Quantitative trait loci  
963 associated with drought tolerance at reproductive stage in rice. *Plant Physiol.* 135, 384-399.

964 Landi, P., Sanguineti, M. C., Salvi, S. (2005). Validation and characterization of a major QTL  
965 affecting leaf ABA concentration in maize. *Mol Breed.* 15, 291-303.

966 Langridge, P., Fleury, D. (2011). Making the most of 'omics' for crop breeding. *Trends*  
967 *Biotechnol.* 29, 33-40.

968 Lata, C., and Prasad, M. (2012). Validation of an allele-specific marker associated with  
969 dehydration stress tolerance in a core set of foxtail millet accessions. *Plant Breed.* 132, 496-  
970 499.

971 Lata, C., and Prasad, M. (2013b). Association of an allele-specific marker with dehydration  
972 stress tolerance in foxtail millet suggests *SiDREB2* to be an important QTL. *J. Plant Biochem.*  
973 *Biotechnol.* 23, 119-122.

974 Lata, C., Bhutty, S., Bahadur, R. P., Majee, M., and Prasad, M. (2011b). Association of an  
975 SNP in a novel DREB2-like gene *SiDREB2* with stress tolerance in foxtail millet [*Setaria*  
976 *italica* (L.)]. *J. Exp. Bot.* 62, 3387-3401.

977 Lata, C., Gupta, S., Prasad, M. (2013). Foxtail millet: a model crop for genetic and genomic  
978 studies in bioenergy grasses. *Crit. Rev. Biotechnol.* 33, 328-343.

979 Laurie, D. A., Pratchett, N., Snape, J. W., Bezant, J. H. (1995). RFLP mapping of five major  
980 genes and eight quantitative trait loci controlling flowering time in a winter x spring barley  
981 (*Hordeum vulgare* L.) cross. *Genome* 38, 575-585.

982 Lei, Y., Lu, L., Liu, H-Y., Li, S., Xing, F., and Chen, L-L. (2014). CRISPR-P: a web tool for  
983 synthetic single-guide RNA design of CRISPR-system in plants. *Mol. Plant* 7, 1494-1496.

- Li, J., Aach, J., Norville, J. E., McCormack, M., Zhang, D., Bush, J., et al. (2013). Multiplex and homologous recombination-mediated genome editing in *Arabidopsis* and *Nicotiana benthamiana* using guide RNA and Cas9. *Nat. Biotechnol.* 31, 688-691.
- Li, Z., Mu, P., Li, C., Zhang, H., Li, Z., Gao, Y., and Wang, X. (2005). QTL mapping of root traits in a doubled haploid population from a cross between upland and lowland japonica rice in three environments. *Theor. Appl. Genet.* 110, 1244-1252.
- Lilley, J. M., Ludlow, M. M., McCouch, S. R., O'Toole, J. C. (1996). Locating QTL for osmotic adjustment and dehydration tolerance in rice. *J. Exp. Bot.* 47, 1427-1436.
- Liu, S., Wang, X., Wang, H., Xin, H., Yang, X., Yan, J., et al. (2013). Genome-Wide Analysis of *ZmDREB* Genes and Their Association with Natural Variation in Drought Tolerance at Seedling Stage of *Zea mays* L. *PLoS Genet.* 9, e1003790.
- Lopes, M. S., Rebetzke, G. J., Reynolds, M. (2014). Integration of phenotyping and genetic platforms for a better understanding of wheat performance under drought. *J. Exp. Bot.* 65, 6167-77.
- Lynch, J. P., Chimungu, J. G., Brown, K. M. (2014). Root anatomical phenes associated with water acquisition from drying 1 soil: Targets for crop improvement. *J. Exp. Bot.* 65, 6155-66.
- Ma, Y., Qin, F., and Tran, L. S. (2012). Contribution of Genomics to Gene Discovery in Plant Abiotic Stress Responses. *Mol. Plant* 5, 1176-1178.
- Mackay, I. J., Bansept-Basler, P., Barber, T., Bentley, A. R., Cockram, J., Gosman, N., et al. (2014). An eight-parent multiparent advanced generation inter-cross population for winter-sown wheat: creation, properties and validation. *G3 (Bethesda)* 4, 1603-1610.
- Mackill, D. J., Ismail, A. M., Singh, U. S., Labios, R. V., and Paris, T. R. (2012). Development and rapid adoption of submergence-tolerant (Sub1) rice varieties. *Adv. Agron.* 115, 299-352.
- Mali, P., Yang, L., Esvelt, K. M., Aach, J., Guell, M., DiCarlo, J. E., et al. (2013). RNA-guided human genome engineering via Cas9. *Science* 339, 823-826.
- Mano, Y., Muraki, M., Fujimori, M., Takamizo, T., Kindiger, B. (2005a). Identification of QTL controlling adventitious root formation during flooding conditions in teosinte (*Zea mays* ssp. *huehuetenangensis*) seedlings. *Euphytica* 142, 33-42.
- Mano, Y., Omori, F., Muraki, M., Takamizo, T. (2005b). QTL mapping of adventitious root formation under flooding conditions in tropical maize (*Zea mays* L.) seedlings. *Breeding Sci.* 55, 343-347.
- Mao, Y., Zhang, H., Xu, N., Zhang, B., Gao, F., and Zhu, J. K. (2013). Application of the CRISPR/Cas system for efficient genome engineering in plants. *Mol. Plant* 6, 2008-2011.
- McCallum, C. M., Comai, L., Greene, E.A., and Henikoff, S. (2000). Targeted screening for induced mutations. *Nat. Biotechnol.* 18, 455-457.

1020 Miao, J., Guo, D., Zhang, J., Huang, Q., Qin, G., Zhang, X., et al. (2013). Targeted  
1021 mutagenesis in rice using CRISPR-Cas system. *Cell Res.* 23, 1233-1236.

1022 Miller, A. K., Galiba, G., and Dubcovsky, J. (2006). A cluster of 11 CBF transcription factors  
1023 is located at the frost tolerance locus Fr-Am2 in *Triticum monococcum*. *Mol. Genet. Genom.*  
1024 275, 193-203.

1025 Miro, B., Ismail, A. M. (2013). Tolerance of anaerobic conditions caused by flooding during  
1026 germination and early growth in rice (*Oryza sativa* L). *Front. Plant Sci.* 4, 269.

1027 Muller, H. J. (1927). Artificial transmutation of the gene. *Science* 66, 84-87.

1028 Munns, R., Tester, M. (2008). Mechanisms of salinity tolerance. *Ann. Rev. Plant Biol.* 59,  
1029 651-681.

1030 Muthamilarasan, M., Prasad, M. (2015). Advances in *Setaria* genomics for genetic  
1031 improvement of cereals and bioenergy grasses. *Theor. Appl. Genet.* 128, 1-14.

1032 Muthamilarasan, M., Theriappan, P., Prasad, M. (2013). Recent advances in crop genomics  
1033 for ensuring food security. *Curr Sci.* 105, 155-158.

1034 Muthamilarasan, M., Venkata Suresh, B., Pandey, G., Kumari, K., Parida, S. K., Prasad, M.  
1035 (2013). Development of 5123 intron-length polymorphic markers for large-scale genotyping  
1036 applications in foxtail millet. *DNA Res.* 21, 41-52.

1037 Nekrasov, V., Staskawicz, B., Weigel, D., Jones, J. D. G., and Kamoun, S. (2013). Targeted  
1038 mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided  
1039 endonuclease. *Nat. Biotechnol.* 31, 691-693.

1040 Neves, L., Davis, J., Barbazuk, B., Kirst, M. (2011). Targeted sequencing in the loblolly pine  
1041 (*Pinus taeda*) megagenome by exome capture. *BMC Proceedings* 7, O48-O48.

1042 Nguyen, T. T. T., Klueva, N., Chamareck, V., Aarti, A., Magpantay, G., Millena, A. C. M., et  
1043 al. (2004). Saturation mapping of QTL regions and identification of putative candidate genes  
1044 for drought tolerance in rice. *Mol. Genet. Genom.* 272, 35-46.

1045 Osakabe, Y., and Osakabe, K. (2015). Genome Editing with Engineered Nucleases in Plants.  
1046 *Plant Cell Physiol.* 56, 389-400.

1047 Paliwal, R., Röder, M. S., Kumar, U., Srivastava, J. P., Joshi, A. K. (2012). QTL mapping of  
1048 terminal heat tolerance in hexaploid wheat (*T. aestivum* L.). *Theor. Appl. Genet.* 125, 561-  
1049 575.

1050 Pandey, G., Misra, G., Kumari, K., Gupta, S., Parida, S. K., Chattopadhyay, D., et al. (2013).  
1051 Genome-wide development and use of microsatellite markers for large-scale genotyping  
1052 applications in foxtail millet [*Setaria italica* (L.)]. *DNA Res.* 20, 197-207.

1053 Pingali, P. L. (2012). Green Revolution: Impact, limits and the path ahead. *Proc. Natl. Acad.*  
1054 *Sci. U. S. A.* 109, 12302-1230.

1055 Pinto, R. S., Reynolds, M. P., Mathews, K. L., McIntyre, C. L., Olivares-Villegas, J. J.,  
1056 Chapman, S. C. (2010). Heat and drought adaptive QTL in a wheat population designed to  
1057 minimize confounding agronomic effects. *Theor. Appl. Genet.* 121, 1001-1021.

- Price, A. H., Cairns, J. E., Horton, P., Jones, H. G., and Griffiths, H. (2002a). Linking drought-resistance mechanisms to drought avoidance in upland rice using a QTL approach: Progress and new opportunities to integrate stomatal and mesophyll responses. *J. Exp. Bot.* 53, 989-1004.
- Price, A. H., Townend, J., Jones, M. P., Audebert, A., and Courtois, B. (2002b). Mapping QTLs associated with drought avoidance in upland rice grown in the philippines and west africa. *Plant Mol. Biol.* 48, 683-695.
- Quarrie, S. A., Laurie, D. A., Zhu, J., Lebreton, C., Semikhodskii, A., Steed, A., et al. (1997). QTL analysis to study the association between leaf size and abscisic acid accumulation in droughted rice leaves and comparisons across cereals. *Plant Mol. Biol.* 35, 155-165.
- Raju, N., Gnanesh, B., Lekha, P., Jayashree, B., Pande, S., Hiremath, P., et al. (2010). The first set of est resource for gene discovery and marker development in pigeonpea (*Cajanus cajan* L.). *BMC Plant Biol.* 10, 45.
- Ramegowda, V., and Senthil-Kumar, M. (2015). The interactive effects of simultaneous biotic and abiotic stresses on plants: Mechanistic understanding from drought and pathogen combination. *J. Plant Physiol.* 176, 47-54.
- Rebetzke, G. J., Herwaarden, A. F., Jenkins, C., Weiss, M., Lewis, D., Ruuska, S., et al. (2008). Quantitative trait loci for water-soluble carbohydrates and associations with agronomic traits in wheat. *Aust. J. Agric. Res.* 59, 891-905.
- Rehman, A. U., Malhotra, R. S., Bett, K., Tar'an, B., Bueckert, R., and Warkentin, T. D. (2011). Mapping QTL associated with traits affecting grain yield in chickpea (*Cicer arietinum* L.) under terminal drought stress. *Crop Sci.* 51, 450-463.
- Rehman, A., Malhotra, R., Bett, K., Tar'an, B., Bueckert, R., Warkentin, T. (2011). Mapping QTL associated with traits affecting grain yield in Chickpea under terminal drought stress. *Crop Sci.* 51, 450-463.
- Ren, Z., Gao, J., Li, L. G., Cai, X., Huang, W., Chao, D., et al. (2005). A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nat. Genet.* 37, 1141-1146.
- Reynolds, M. P., Hays, D., and Chapman, S. (2010). "Breeding for adaptation to heat and drought stress," in *Climate change and crop production*, ed. M. P. Reynolds (Oxfordshire, UK: CABI), 71-91.
- Reynolds, M., Tuberosa, R. (2008). Translational research impacting on crop productivity in drought-prone environments. *Curr. Opin. Plant Biol.* 11, 171-179.
- Robin, S., Pathan, M. S., Courtois, B., Lafitte, R., Carandang, S., Lanceras, S., et al. (2003). Mapping osmotic adjustment in an advanced back-cross inbred population of rice. *Theor. Appl. Genet.* 107, 1288-1296.
- Robin, S., Pathan, M. S., Courtois, B., Lafitte, R., Carandang, S., Lanceras, S., et al. (2003.) Mapping osmotic adjustment in an advanced back-cross inbred population of rice. *Theor. Appl. Genet.* 107, 1288-1296.



1096 Ruta, N., Liedgens, M., Fracheboud, Y., Stamp, P., Hund, A. (2010). QTLs for the elongation  
1097 of axile and lateral roots of maize in response to low water potential. *Theor. Appl. Genet.* 120,  
1098 621-631.

1099 Salem, K. F. M., Roder, M. S., Borner, A. (2007). Identification and mapping quantitative  
1100 trait loci for stem reserve mobilisation in wheat (*Triticum aestivum* L.). *Cereal Res. Commun.*  
1101 35, 1367-1374.

1102 Salvi, S., Sponza, G., Morgante, M., Tomes, D., Niu, X., Fengler, K. A., et al. (2007).  
1103 Conserved noncoding genomic sequences associated with a flowering-time quantitative trait  
1104 locus in maize. *Proc. Natl. Acad. Sci. U. S. A.* 104, 11376-11381.

1105 Sarvamangala, C., Gowda, M. V. C., Varshney, R. K. (2011). Identification of quantitative  
1106 trait loci for protein content oil content and oil quality for groundnut (*Arachis hypogaea* L.).  
1107 *Field Crops Res.* 122, 49-59.

1108 Saxena, K. (2008). Genetic improvement of pigeonpea-a review. *Trop. Plant Biol.* 1, 159-178.

1109 Saxena, R., Prathima, C., Saxena, K., Hoisington, D., Singh, N., and Varshney, R. (2010).  
1110 Novel SSR markers for polymorphism detection in pigeonpea (*Cajanus* spp.). *Plant Breed.*  
1111 129, 142-148.

1112 Schmutz, J., McClean, P. E., Mamidi, S., Wu, G. A., Cannon, S. B., Grimwood J., et al.  
1113 (2014). A reference genome for common bean and genome-wide analysis of dual  
1114 domestications. *Nat. Genet.* 46, 707-713.

1115 Schneider, K. A., Brothers, M. E., Kelly, J. D. (1997). Marker-assisted selection to improve  
1116 drought tolerance in common bean. *Crop Sci.* 37, 51-60.

1117 Seki, M., Narusaka, M., Ishida, J., Nanjo, T., Fujita, M., Oono, Y., et al. (2002). Monitoring  
1118 the expression profiles of 7000 Arabidopsis genes under drought, cold and high-salinity  
1119 stresses using a full-length cDNA microarray. *Plant J.* 31, 279-292.

1120 Septiningsih, E. M., Ignacio, J. C., Sendon, P. M., Sanchez, D. L., Ismail, A. M., Mackill, D.  
1121 J. (2013). QTL mapping and confirmation for tolerance of anaerobic conditions during  
1122 germination derived from the rice landrace Ma-Zhan Red. *Theor. Appl. Genet.* 126, 1357-  
1123 1366.

1124 Serraj, R., Kumar, A., McNally, K. L., Slamet-Loedin, I., Bruskiewich, R., Mauleon, R., et al.  
1125 (2009). Improvement of drought resistance in rice, *Adv. Agron.* 103, 41-99.

1126 Setter, T. M., Waters, I. (2003). Review of prospects for germplasm improvement for  
1127 waterlogging tolerance in wheat, barley and oats. *Plant Soil* 253, 1-34.

1128 Shan, Q., Wang, Y., Li, J., Zhang, Y., Chen, K., Liang, Z., et al. (2013). Targeted genome  
1129 modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* 31, 686-688.

1130 Silva, G., Poirot, L., Galetto, R., Smith, J., Montoya, G., Duchateau, P., and Pâques, F.  
1131 (2011). Meganucleases and Other Tools for Targeted Genome Engineering: Perspectives and  
1132 Challenges for Gene Therapy. *Curr. Gene Ther.* 11, 11-27.

1133 Singh, S. P. (2007). Drought resistance in the race durango dry bean landraces and cultivars.  
1134 *Agronomy J.* 99, 1219-1225.

- Singh, S., Mackill, D. J., and Ismail, A. M. (2009). Responses of Sub1 rice introgression lines to submergence in the field: Yield and grain quality. *Field Crops Res.* 113, 12-23.
- Skinner, J. S., Szucs, P., Zitzewitz, Jv., Marquez-Cedillo, L., Filichkin, T., Stockinger, E. J., et al. (2006). Mapping of barley homologs to genes that regulate low temperature tolerance in arabidopsis. *Theor. Appl. Genet.* 112, 832-842.
- Snape, J. W., Foulkes, J., Simmonds, J., Leverington, M., Fish, L. J., Wang, Y., et al. (2007). Dissecting gene  $\times$  environmental effects on wheat yields via QTL and physiological analysis. *Euphytica* 154, 401-408.
- Stockinger, E. J., Gilmour, S. J., and Thomashow, M. F. (1997). Arabidopsis thaliana CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc. Natl. Acad. Sci. U. S. A.* 94, 1035-1040.
- Sujay, V., Gowda, M. V., Pandey, M. K., Bhat, R. S., Khedikar, Y. P., Nadaf, H. L., et al. (2012). Quantitative trait locus analysis and construction of consensus genetic map for foliar disease resistance based on two recombinant inbred line populations in cultivated groundnut (*Arachis hypogaea* L.). *Mol Breed.* 30, 773-788.
- Suresh, B. V., Muthamilarasan, M., Misra, G., Prasad, M. (2013). FmMDb: A versatile database of foxtail millet markers for millets and bioenergy grasses research. *PLoS ONE* 8, e71418.
- Thomas, H., James, A. R. (1993). Freezing tolerance and solute changes in contrasting genotypes of *Lolium perenne* L. acclimated to cold and drought. *Ann. Bot.* 72, 249-254.
- Thomson, M. J., Ocampo, Md., Egdane, J., Rahman, M. A., Sajise, A. G., Adorada, D. L., et al. (2010). Characterizing the saltol quantitative trait locus for salinity tolerance in rice. *Rice* 3, 148-160.
- Thudi, M., Bohra, A., Nayak, S. N., Varghese, N., Shah, T. M., Penmetsa, R. V., et al. (2011). Novel SSR markers from bac-end sequences, dart arrays and a comprehensive genetic map with 1, 291 marker loci for chickpea (*Cicer arietinum* L.). *PLoS ONE* 6, e27275.
- Townsend, J. A., Wright, D. A., Winfrey, R. J., Fu, F., Maeder, M. L., Joung, J. K., and Voytas, D. F. (2009). High-frequency modification of plant genes using engineered zinc-finger nucleases. *Nature* 459, 442-445.
- Tripathy, J. N., Zhang, J., Robin, S., Nguyen, T. T., and Nguyen, H. T. (2000). QTLs for cell-membrane stability mapped in rice (*Oryza sativa* L.) under drought stress. *Theor. Appl. Genet.* 100, 1197-1202.
- Tripathy, J. N., Zhang, J., Robin, S., Nguyen, T. T., Nguyen, H. T. (2000). QTLs for cell-membrane stability mapped in rice (*Oryza sativa* L.) under drought stress. *Theor. Appl. Genet.* 100, 1197-1202.
- Tuberosa, R. (2012). Phenotyping for drought tolerance of crops in the genomics era. *Front. Physiol.* 3, 347.

1173 Tuberosa, R., Salvi, S., Sanguineti, M. C., Frascaroli, E., Conti, S., Landi, P. (2011).  
1174 “Genomics of root architecture and functions in maize,” in *Root genomics*, eds. In: A. Costa  
1175 de Oliveira, R. Varshney (Berlin: Springer), 179-204.

1176 Uga, Y., Sugimoto, K., Ogawa, S., Rane, J., Ishitani, M., Hara, N., et al. (2013). Control of  
1177 root system architecture by deeper rooting 1 increases rice yield under drought conditions.  
1178 *Nat. Genet.* 45, 1097-1102.

1179 Urrea, C. A., Yonts, C. D., Lyon, D. J., Koehler, A. E. (2009). Selection for drought tolerance  
1180 in dry bean derived from the Mesoamerican gene pool in western Nebraska. *Crop Sci.* 49,  
1181 2005-2010.

1182 Vadez, V., Krishnamurthy, L., Thudi, M., Anuradha, C., Colmer, T. D., Turner, N. C., et al.  
1183 (2012). Assessment of ICCV 2 × JG 62 chickpea progenies shows sensitivity of reproduction  
1184 to salt stress and reveals QTL for seed yield and yield components. *Mol. Breed.* 30, 9-21.

1185 Vagujfalvi, A., Galiba, G., Cattivelli, L., and Dubcovsky, J. (2003). The cold-regulated  
1186 transcriptional activator Cbf3 is linked to the frost-tolerance locus Fr-A2 on wheat  
1187 chromosome 5A. *Mol. Genet. Genom.* 269, 60-67.

1188 Varshney, R. K., Chen, W., Li, Y., Bharti, A. K., Saxena, R. K., Schlueter, J. A., et al. (2012).  
1189 Draft genome sequence of pigeonpea (*Cajanus cajan*), an orphan legume crop of resource-  
1190 poor farmers. *Nat. Biotechnol.* 30, 83-89.

1191 Varshney, R. K., Graner, A., and Sorrells, M. E. (2005). Genomics-assisted breeding for crop  
1192 improvement. *Trends Plant Sci.* 10, 621-630.

1193 Varshney, R. K., Mohan, S. M., Gaur, P. M., Gangarao, N. V., Pandey, M. K., Bohra, A., et  
1194 al. (2013a). Achievements and prospects of genomics-assisted breeding in three legume crops  
1195 of the semi-arid tropics. *Biotechnol. Adv.* 31, 1120-1134.

1196 Varshney, R. K., Penmetsa, R. V., Dutta, S., Kulwal, P. L., Saxena, R. K., Datta, S., et al.  
1197 (2009). Pigeonpea genomics initiative (PGI): an international effort to improve crop  
1198 productivity of pigeonpea (*Cajanus cajan* L.). *Mol. Breed.* 26, 393-408.

1199 Varshney, R. K., Song, C., Saxena, R. K., Azam, S., Yu, S., Sharpe, A. G., et al. (2013b).  
1200 Draft genome sequence of kabuli chickpea (*Cicer arietinum*): genetic structure and breeding  
1201 constraints for crop improvement. *Nat. Biotechnol.* 31, 240-246.

1202 Varshney, R. K., Terauchi, R., and McCouch, S. R. (2014). Harvesting the Promising Fruits of  
1203 Genomics: Applying Genome Sequencing Technologies to Crop Breeding. *PLoS Biol.* 12,  
1204 e1001883.

1205 Velasco, R., Zharkikh, A., Affourtit, J., Dhingra, A., Cestaro, A., Kalyanaraman, A., et al.  
1206 (2010). The genome of the domesticated apple (*Malus x domestica* Borkh.). *Nat. Genet.* 42,  
1207 833-839.

1208 Velasco, R., Zharkikh, A., Troggio, M., Cartwright, D. A., Cestaro, A., Pruss, D., et al.  
1209 (2007). A high quality draft consensus sequence of the genome of a heterozygous grapevine  
1210 variety. *PLoS ONE* 2, e1326.

- Wang, H., Penmetsa, R. V., Yuan, M., Gong, L., Zhao, Y., Guo, B., et al. (2012). Development and characterization of BAC-end sequence derived SSRs, and their incorporation into a new higher density genetic map for cultivated peanut (*Arachis hypogaea* L.). *BMC Plant Biol.* 12, 10.
- Wang, Y., Cheng, X., Shan, Q., Zhang, Y., Liu, J., Gao, C., and Qiu, J-L. (2014). Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* 32, 947-951.
- Wassmann, R., Jagadish, S. (2009). Regional vulnerability of climate change impacts on Asian rice production and scope for adaptation. *Adv. Agron.* 102, 91-133.
- Wasson, A. P., Richards, R. A., Chatrath, R., Misra, S. C., Prasad, S. V., Rebetzke, G. J., et al. (2012). Traits and selection strategies to improve root systems and water uptake in water-limited wheat crops. *J. Exp. Bot.* 63, 3485-3498.
- Welch, J. R., Vincent, J. R., Auffhammer, M., Moya, P. F., Dobermann, A., Dawe, D. (2010). Rice yields in tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures. *Proc. Natl. Acad. Sci. U. S. A.* 107, 14562-14567.
- Xie, K., and Yang, Y. (2013). RNA-Guided Genome Editing in Plants Using a CRISPR-Cas System. *Mol. Plant* 2013; 6, 1975-1980.
- Xu, J. L., Lafitte, H. R., Gao, Y. M., Fu, B. Y., Torres, R., and Li, Z. K. (2005). QTLs for drought escape and tolerance identified in a set of random introgression lines of rice. *Theor. Appl. Genet.* 111, 1642-1650.
- Xu, K., Xu, X., Fukao, T., Canlas, P., Maghirang-Rodriguez, R., Heuer, S., et al. (2006). *Sub1A* is an ethylene-response-factor-like gene that confers submergence tolerance to rice. *Nature* 442, 705-708.
- Xu, Q., Chen, L. L., Ruan, X., Chen, D., Zhu, A., Chen, C., et al. (2013). The draft genome of sweet orange (*Citrus sinensis*). *Nat. Genet.* 45, 59-U92.
- Yadav, C. B., Bonthala, V. S., Muthamilarasan, M., Pandey, G., Khan, Y., Prasad, M. (2015). Genome-wide development of transposable elements-based markers in foxtail millet and construction of an integrated database. *DNA Res.* 22, 79-90.
- Yadav, C. B., Muthamilarasan, M., Pandey, G., Khan, Y., Prasad, M. (2014). Development of novel microRNA-based genetic markers in foxtail millet for genotyping applications in related grass species. *Mol. Breed.* doi:10.1007/s11032-014-0137-9.
- Yadav, R. S., Sehgal, D., Vadez, V. (2011). Using genetic mapping and genomics approaches in understanding and improving drought tolerance in pearl millet. *J. Exp. Bot.* 62, 397-408.
- Yang, J., Zhang, J., Liu, K., Wang, Z., Liu, L. (2007). Absciscic acid and ethylene interact in rice spikelets in response to water stress during meiosis. *J. Plant Growth Regul.* 26, 318-328.
- Yue, B., Xue, W., Xiong, L., Yu, X., Luo, L., Cui, K., et al. (2006). Genetic basis of drought resistance at reproductive stage in rice: separation of drought tolerance from drought avoidance. *Genetics* 172, 1213-1228.

- Yue, B., Xue, W., Xiong, L., Yu, X., Luo, L., Cui, K., et al. (2006). Genetic basis of drought resistance at reproductive stage in rice: Separation of drought tolerance from drought avoidance. *Genetics* 172, 1213-1228.
- Zhang, G., Liu, X., Quan, Z., Cheng, S., Xu, X., Pan, S., et al. (2012). Genome sequence of foxtail millet (*Setaria italica*) provides insights into grass evolution and biofuel potential. *Nature Biotechnol.* 30, 549-554.
- Zhang, H., Zhang, J., Wei, P., Zhang, B., Gou, F., Feng, Z., et al. (2014). The CRISPR/Cas9 system produces specific and homozygous targeted gene editing in rice in one generation. *Plant Biotechnol. J.* 12, 797-807.
- Zhang, J., Nguyen, B. D., Sarkarung, S., Blum, A., Nguyen, H. T., Zheng, H. G., et al. (2001a). Locating genomic regions associated with components of drought resistance in rice: Comparative mapping within and across species. *Theor. Appl. Genet.* 103, 19-29.
- Zhang, S., Tang, C., Zhao, Q., Li, J., Yang, L., Qie, L., et al. (2014). Development of highly polymorphic simple sequence repeat markers using genome-wide microsatellite variant analysis in Foxtail millet [*Setaria italica* (L.) P. Beauv]. *BMC Genomics* 2, 15.
- Zhang, W. P., Shen, X. Y., Wu, P., Hu, B., and Liao, C. Y. (2001b). QTLs and epistasis for seminal root length under a different water supply in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 103, 118-123.
- Zheng, B. S., Yang, L., Zhang, W. P., Mao, C. Z., Wu, Y. R., Yi, K. K., et al. (2003). Mapping QTLs and candidate genes for rice root traits under different water-supply conditions and comparative analysis across three populations. *Theor. Appl. Genet.* 107, 1505-1515.
- Zheng, H. G., Babu, R. C., Pathan, M. S., Ali, L., Huang, N., Courtois, B., and Nguyen, H. T. (2000). Quantitative trait loci for root-penetration ability and root thickness in rice: Comparison of genetic backgrounds. *Genome / National Research Council Canada* 43, 53-61.
- Zhou, X., Jacobs, T. B., Xue, L.-J., Harding, S. A., and Tsai, C.-J. (2015). Exploiting SNPs for biallelic CRISPR mutations in the outcrossing woody perennial *Populus* reveals 4-coumarate:CoA ligase specificity and redundancy. *New Phytol.* doi: 10.1111/nph.13470.

1280 **Figure legends**

1281

1282 **Figure 1:** Flow-chart demonstrating the steps involved in generating climate resilient crops using  
1283 genomics and next-generation sequencing technology.

Figure 1.TIF

