

Aberystwyth University

Smell of Infection

Pawluk, Rebecca; Stuart, Rebekah; Garcia de Leaniz, Carlos; Cable, Jo; Morpew, Russ; Brophy, Peter; Consuegra, Sofia

Published in:

Journal of Proteome Research

DOI:

[10.1021/acs.jproteome.8b00953](https://doi.org/10.1021/acs.jproteome.8b00953)

Publication date:

2019

Citation for published version (APA):

Pawluk, R., Stuart, R., Garcia de Leaniz, C., Cable, J., Morpew, R., Brophy, P., & Consuegra, S. (2019). Smell of Infection: a novel, non-invasive method for detection of fish excretory- secretory proteins. *Journal of Proteome Research*, 18(3), 1371-1379. <https://doi.org/10.1021/acs.jproteome.8b00953>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Smell of Infection: a novel, non-invasive method for detection of fish excretory- secretory proteins

Rebecca J. Pawluk, Rebekah Stuart, Carlos Garcia de Leaniz, Joanne Cable, Russell M. Morphew, Peter M. Brophy, and Sofia Consuegra

J. Proteome Res., **Just Accepted Manuscript** • DOI: 10.1021/acs.jproteome.8b00953 • Publication Date (Web): 21 Dec 2018

Downloaded from <http://pubs.acs.org> on January 7, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3 **Smell of Infection: a novel, non-invasive method for detection of fish excretory-secretory**
4 **proteins**
5
6
7

8 Rebecca J. Pawluk^{1*}, Rebekah Stuart², Carlos Garcia de Leaniz¹, Joanne Cable⁴, Russell M.
9 Morphew³, Peter M. Brophy³, Sofia Consuegra¹
10
11
12
13
14
15
16

17 ¹Swansea University, College of Science, Biosciences, Swansea, SA2 8PP, UK
18
19

20 ²Wales Veterinary Science Centre, Buarth, Aberystwyth, Ceredigion, SY23 1ND, UK
21
22

23 ³Aberystwyth University, IBERS, Penglais, Aberystwyth, Ceredigion, SY23 3FL, UK
24
25

26 ⁴Cardiff University, School of Biosciences, Cardiff, CF10 3AX, UK
27
28

29 *Correspondence to rebeccajane93@hotmail.co.uk
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Chemical signals are produced by aquatic organisms following predatory attacks or perturbations such as parasitic infection. Ectoparasites feeding on fish hosts are likely to cause release of similar alarm cues into the environment due to the stress, wounding and immune response stimulated upon infection. Alarm cues are often released in the form of proteins, anti-microbial peptides and immunoglobulins that provide important insights into bodily function and infection status. Here we outline a non-invasive method to identify potential chemical cues associated with infection in fish by extracting, purifying and characterizing proteins from water samples from cultured fish. Gel free proteomic methods were deemed the most suitable for protein detection in saline water samples. It was confirmed that teleost proteins can be characterised from water and that variation in protein profiles could be detected between infected and uninfected individuals and fish and parasite only water samples. Our novel assay provides a non-invasive method for assessing the health condition of both wild and farmed aquatic organisms. Similar to environmental DNA monitoring methods, these proteomic techniques could provide an important tool in applied ecology and aquatic biology.

KEYWORDS: Alarm cues, gel free MS, *Kryptolebias marmoratus*, odour, parasitic infection

1. Introduction

Chemical cues released into the environment following natural perturbations, predator attacks or social threats allow animals to assess potential dangers and may lead to increased survival chances (1). In the aquatic environment, cues released in the form of soluble pheromones or as chemicals following physical trauma to the epidermis are described as alarm cues (2). These cues can signal the presence of danger, including the scent of predatory species in the environment (disturbance cues) (3, 4) or the odour of parasitized conspecifics (5). Additionally, innate immune responses activated upon pathogen infection can cause variation in body odour; this has been documented in humans for example, whereby endotoxin-exposed individuals smelt significantly more unpleasant than their control counterparts indicating a social cue of sickness (6). Changes in body odour have also been seen in infected mice, where females prefer the urinary odours of uninfected male individuals (7). Yet, studies that describe infection odours in aquatic environments are scarce (8, 9), despite the importance of chemical communication, particularly in potentially turbid conditions (10).

Chemical cues also allow individuals to recognise neighbours, assess social hierarchies, and determine mating ability by genotype matching, which may translate into attraction or conspecific avoidance (11, 12). Odour based mate choices have been linked to the Major Histocompatibility Complex (MHC) and its associated peptides (13) and allow individuals to discriminate between MHC genotypes on the basis of variation in immune-related peptides released into the water (13). This enables mate choice to maximise the immuno-competence of offspring (14), and MHC-related peptides are likely used for relatedness discrimination in several species e.g. *Salvelinus alpinus* (15) and *Gasterosteus aculeatus* (13). Analyses of social communication in mice has also provided evidence that a variety of polymorphic proteins in urine (major urinary proteins) provide a unique identity to the owner and are used as scent signals (16).

1
2
3 Many teleost fish can produce and perceive water-borne cues released by stressed conspecifics,
4 including free cortisol (17) and alarm cues released by specialized skin cells when individuals
5 are injured (18). Zebrafish (*Danio rerio*), for example, are able to detect cues in the water from
6 physically stressed individuals and avoid them (1) and Atlantic salmon can detect when a
7 predator has consumed conspecifics (19). Similarly, Nile tilapia (*Oreochromis niloticus*)
8 display anti-predatory behaviours (including freezing and dorsal fin erection) in response to
9 water conditioned with conspecific skin extract that mimics a predatory attack (20). Alarm cues
10 also arise from parasitic infections, particularly ectoparasites that cause trauma to the epidermis
11 initiating immune responses in a similar way to predator attacks (21, 22). Argulid parasites
12 attach to host fish by large maxillary suckers and feed on host blood and skin (23), this type of
13 parasite feeding causes skin wounding, ulcers and increases the susceptibility of the host to
14 secondary infections (24). Immune responses in fish skin, largely identified in the mucosal
15 layer include several components that reduce pathogen and allergen entry (25). Such
16 components consist of antimicrobial peptides and enzymes (such as lysozymes, phosphatases
17 and proteases) for enhanced protection and wound healing (26-28), lectins and proteins that
18 enhance pathogen expulsion and immune activation (25, 29, 30) and immunoglobulin
19 antibodies for protection against surface infections (31). It is on this basis that the current study
20 aimed to investigate excretory-secretory proteins released by hosts and parasites upon
21 infection.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46
47 Gel based and gel free proteomic techniques are often used to detect proteins from body tissues
48 of an organism (32, 33) and are also useful for studying the excretory-secretory proteins
49 produced by endo-parasitic helminths (34). For example, *Fasciola hepatica*, an economically
50 important helminth of livestock is well studied, partly as it can easily be cultured in media,
51 providing a clearer understanding of host-parasite communication and host manipulation (35-
52 37). Fewer studies use *in vitro* culturing techniques to obtain proteins from ectoparasitic
53
54
55
56
57
58
59
60

1
2
3 organisms; culturing of such parasites commonly involves rearing host teleost species under
4 laboratory conditions followed by subsequent infection of these hosts via anaesthesia (46).
5
6 Despite difficulties associated with culturing these parasites (38), such systems provide a
7
8 unique chance to collect and analyse protein data from host-parasite interactions. Furthermore,
9
10 there is a distinct lack of knowledge surrounding environmental proteins from fresh and marine
11
12 water bodies and the potential use of proteins as a non-invasive detection method for infection.
13
14 Research into meta-proteomics of water samples to date has focused on methods for marine
15
16 sediments (e.g. 39). Yet, these methods have never been applied to teleost fish, particularly in
17
18 brackish water samples or in relation to infection or parasite presence, which can be critical the
19
20 early detection of infection in aquatic species.
21
22
23
24
25
26

27 *Kryptolebias marmoratus* (the mangrove killifish) is a small, naturally inbred, cyprinid fish
28
29 species that is easily maintained under laboratory conditions. Isogeneity in this species via
30
31 constant selfing makes it a useful model organism for the study of physiological plasticity and
32
33 response to environmental stressors (40, 41). *Argulus foliaceus*, is a generalist fish parasite
34
35 found throughout Europe that is known to cause epizootics and problems in both brackish and
36
37 freshwater fish farms (42-45). Here we assessed the use of a novel non-invasive proteomic
38
39 approach to identify fish infection by comparing proteins released into water by infected and
40
41 uninfected *K. marmoratus*. Additionally, we used this study system to develop a proteomic
42
43 method of potential application to assess health status and stress response in fish populations
44
45
46
47

48 **2. Experimental Procedures**

49 *2.1 Water sample collection, preparation and visualisation*

50
51 The study species, *K. marmoratus*, originated in Belize, from two self-fertilising lines (DAN
52
53 and R), that have undergone >30 generations of selfing (46). Eighty size- matched *K.*
54
55 *marmoratus* (12 to 14 months old) reared individually from hatching were selected from the
56
57
58
59
60

1
2
3 two lines (40 DAN and 40 R) and kept in individual aquaria (12 L x 8 W x 8.5 H cm) containing
4
5 750 ml of brackish water (15 ppt salinity, constituted from dechlorinated water and marine
6
7 filtered water) under controlled conditions (12L:12D photoperiod, $24 \pm 1^{\circ}\text{C}$).
8
9

10
11 Twenty fish from each line were individually infected with one adult individual of the
12
13 ectoparasitic crustacean *Argulus foliaceus*. The culture of *A. foliaceus* originated from carp
14
15 (*Cyprinus carpio*) caught in a still water fishery in North Lincolnshire, July 2014, and thereafter
16
17 was maintained on *Gasterosteus aculeatus* (three-spined sticklebacks) at Cardiff University as
18
19 detailed in Stewart *et al.* (2017) (47). The other twenty fish from each line were kept as control
20
21 individuals, as described in Pawluk (48). Two water samples for proteomic analyses (50 ml
22
23 each) were collected from each aquarium (80 aquaria total) prior to filtration (Minisart 25 mm
24
25 Pore size 0.2 μm filter) and storage at -80°C , the second sample acting as a spare. Twenty water
26
27 samples (50 ml each) were also taken from containers of the same size containing a single *A.*
28
29 *foliaceus* as a control for parasite proteins released into the water. Amicon filter units were
30
31 used to reduce water samples from 36 ml (3 subsamples of 12ml) to approximately 2 ml
32
33 (10KDa cut off, 5000 x g for 10 minutes, with two filter washes); subsequent quantification
34
35 (Bradford, 49) using Sigma Bradford Reagent (according to the manufacturer's instructions)
36
37 and a Cary 50 Bio UV-visible spectrophotometer at 595 nm was completed. All samples were
38
39 precipitated using the addition of 4 volumes of ice cold acetone followed by 1-hour incubation
40
41 at -20°C . Proteins were then pelleted by centrifugation at 4°C and 21000 x g for 15 mins. Pellets
42
43 were then dried for 30 mins before being resuspended in 30 μl of Buffer Z (8 M Urea, 2% w/v
44
45 CHAPS, 33 mM DTT, 0.5% ampholytes pH range 3-10). Subsequent 1D SDS-page was
46
47 performed using polyacrylamide gels (12.5%); they were then fixed overnight (in 10% ethanol,
48
49 40% acetic acid) and silver stained (50). The molecular weight of protein bands identified on
50
51 1D gels were calculated using a standard curve of $\text{Log}(\text{Mw})$ versus the mobility shift (R_f).
52
53
54
55
56
57
58
59
60

2.3 Trypsin digestion and gel free mass spectrophotometry

Fifteen water samples were prepared using Amicon filtration, as described previously, prior to trypsin digest and mass spectrometry analysis (3 individual fish replicates per group: DAN infected, DAN control, R infected, R control and *Argulus* only). Sample aliquots of 100 μl each were added to 6 M urea, in 100 mM tris buffer, and then reduced using 5 μl reducing agent containing DTT and Tris stock (200 mM DTT, 100 mM Tris) for 1 hour. Subsequently, 20 μl alkylating agent (200 mM iodoacetamide and 100 mM Tris) was added to each sample and vortexed before 1 hour incubation at room temperature. A further 20 μl of reducing agent was added per sample and incubated at room temperature for 1 hour. Sample urea concentration was reduced (to approx. 0.6 M) by diluting each sample with 75 μl of water before mixing. Trypsin digestion began following the addition of trypsin solution to a final concentration of 50 ng/ μl , which was subsequently mixed and then incubated overnight at 37°C. The reaction was stopped by adjusting the pH to >6 by adding concentrated acetic acid. All 15 samples were reduced to 100 μl aliquots using a speed vacuum prior to analysis liquid Chromatography tandem mass spectrometry using the Agilent 6550 iFunnel Q-TOF mass spectrometer with Dual AJS ESI source coupled to a 1200 series HPLC-Chip system (Agilent, Cheshire, UK).

The HPLC-Chip/Q-TOF system was equipped with a capillary loading pump (1200 series, Agilent Technologies) and a nano pump (1200 series, Agilent Technologies). Sample injection was conducted with a micro auto sampler (1100 series, Agilent Technologies), where 1 μl of sample in 0.1% formic acid was loaded on to the enrichment column at a flow of 2.5 $\mu\text{L}/\text{min}$ followed by separation at a flow of 300 nL/min. A Polaris Chip was used (G4240-62030, Agilent Technologies), comprising a C18 enrichment/trap column (360 nl) and a C18 separation column (150 mm x 75 $\hat{\text{A}}\mu\text{m}$), where ions were generated at a capillary voltage of 1950 V. The solvent system was: solvent A (ultra-pure water with 0.1% formic acid), and solvent B (90% acetonitrile with 0.1% formic acid). The liquid chromatography was performed

1
2
3 with a piece-linear gradient using 3-8% of solvent B over 0.1 minutes, 8-35% solvent B over
4
5 14.9 minutes, 35-90% solvent B over 5 minutes and hold at 90% solvent B for 2 minutes.
6
7 Tandem mass spectrometry was performed in AutoMS2 mode in the 300-1700 Da range, at a
8
9 rate of 5 spectra per second, performing MS2 on the 5 most intense ions in the precursor scan.
10
11 Masses were excluded for 0.1 min after MS/MS was performed. Reference mass locking was
12
13 used for internal calibration using the mass of 391.2843 Da. Peak lists were generated with
14
15 Mass Hunter Qualitative Analysis software (V B.06, Agilent Technologies) and exported as
16
17 Mascot Generic Files.
18
19
20
21

22 *2.4 Data submission and assessment based on MASCOT scoring*

23
24

25 Mass spectral data from all 15 samples were submitted for database searching using the
26
27 MASCOT program (Matrix Science Ltd., version 2.1). Search parameters allowed a maximum
28
29 of one missed cleavage. Additionally, a fixed modification was set for cysteine at 161 Da and
30
31 variable modifications tested for matches 0, 1, 2, or 3 oxidised methionine residues. Lastly, a
32
33 set peptide tolerance of 1.2 Da and MS/MS tolerance of 0.6 Da were stipulated. Spectra were
34
35 searched against the *K. marmoratus* transcriptome (48) as well as a recent version of the NCBI
36
37 non-redundant protein database sequences. Both analyses resulted in a total of 30 protein lists
38
39 that were subsequently sorted into treatment groups; each sample from each group was filtered
40
41 for protein number above the MASCOT significance threshold at $p = 0.05$ (Transcriptome data
42
43 ≥ 50 , NCBI data ≥ 59), protein identified in all three replicates and proteins identified in two of
44
45 three individuals for each treatment (Table S2 & S3). The mass spectrometry proteomics data
46
47 have been deposited to the ProteomeXchange Consortium via the PRIDE (51) partner
48
49 repository with the dataset identifier PXD010987.
50
51
52
53
54
55

56 *2.5 GO annotation, pathway analysis and statistical testing*


57
58
59
60

1
2
3 All proteins for each sample and treatment were considered to compare data searches and to
4 complete a functional analysis (GO annotation, level GO ALL) of the results using the
5 Database for Annotation, Visualization and Integrated Discovery, version 6.8 (DAVID) (52).
6
7 Pathway analysis was completed using REACTOME pathways (53) for the three biological
8 replicates of each group. Lastly, protein release for treatment groups was compared statistically
9 using t-tests. MASCOT scores for samples from the two respective groups (Infected and
10 control) were compared for specific proteins (OSER1, KRT94). Additionally the number of
11 proteins related to specific functional groups (identified from DAVID) were compared for
12 methylation and immune functions. All figures produced for data visualisation were created
13 using R version 3.4.0 (54).
14
15
16
17
18
19
20
21
22
23
24
25
26

27 3. Results

28 3.1 Protein visualisation and functional analyses

29
30 Results from quantification and one dimensional SDS-page indicated the presence of proteins
31 at approximately 61.6, 47.9, 42.3 and 24.8 kDa (Figure 1).
32
33
34
35
36
37
38



57 **Figure 1.** SDS-PAGE gel of the expression of proteins concentrated using Amicon filtration
58 steps, from two strains (D: DAN and R: R) for unwashed (UW) and washed (W) filters.
59
60

1
2
3 As 1D gel electrophoresis successfully indicated purified proteins a gel free approach was taken
4 to identify all proteins within these samples. Soluble proteins from water samples were
5 identified using both the *K. marmoratus* transcriptome (48) (MASCOT score cut of ≥ 50) and
6 from a search of the NCBI database (MASCOT score cut of > 59). The total number of proteins
7 identified for the *K. marmoratus* transcriptome and NCBI database ranged from 28 to 53 per
8 sample. The percentage of identified proteins over the MASCOT cut off score for each data set
9 ranged from 2.9 – 16.3% from transcriptome sourced data and 7.7-20.9% from NCBI sourced
10 data (Table S1). Trypsin (used for digestion) and keratin, mostly deemed to be from human
11 origin and thus contaminants were highly abundant (had a MASCOT score on or above the cut
12 off) and overlapped between search methods (Transcriptome and NCBI) (Table S2 and S3).
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 Gene ontology analysis indicated that functional proteins involved with immune response and
28 development (Figure S2 and S3) were identified in both data sets however on inspection of
29 these proteins there was little overlap between search methods. Pathway analysis (Table 1,
30 Table S5 and Figure S3) was completed for a comparison of enriched protein groups for all 5
31 groups (DAN control and infected, R control and infected, and *Argulus* only water). For all
32 fish water samples (DAN and R, infected and control) proteins were mainly involved in
33 metabolism, in three of these groups these proteins had the highest number of hits.
34 Proportionally fewer proteins were reported in *Argulus* only samples when compared with the
35 others and proteins from this group primarily had roles in transcription and homeostasis.
36 Proteins with roles in cytokine signalling were shown to be in infected fish water samples but
37 not in DAN.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. The top five entities identified from the REACTOME pathway analysis of 5 groups of water samples (from DAN infected, DAN control, R infected and R control *Kryptolebias marmoratus* as well as *Argulus* only samples).

Group	Pathway name	Entities found	Entities Total	Entities pvalue	Entities FDR	Reactions found
<i>Dan infected</i>	Metabolism	10	2,116	0.84	8.35E-01	34
	Immune System	6	2,226	1.00	9.95E-01	17
	Adaptive Immune System	3	944	0.93	9.28E-01	6
	Generic Transcription Pathway	3	1,152	0.97	9.73E-01	7
	RNA Polymerase II Transcription	3	1,274	0.99	9.86E-01	18
<i>Dan control</i>	Metabolism	6	2,116	1.00	9.97E-01	9
	Disease	4	1,148	0.95	9.52E-01	25
	Innate Immune System	4	1,180	0.96	9.59E-01	12
	Post-translational protein modification	3	1,415	1.00	9.97E-01	21
	Signaling by Receptor Tyrosine Kinases	2	471	0.82	8.22E-01	2
<i>R infected</i>	Metabolism	8	2,116	0.99	9.92E-01	23
	Immune System	8	2,226	1.00	9.96E-01	17
	Metabolism of proteins	6	2,111	1.00	9.99E-01	11
	Signal Transduction	6	2,738	1.00	1.00E+00	8
	Innate Immune System	5	1,180	0.94	9.41E-01	8
<i>R control</i>	Signal Transduction	10	2,738	0.98	9.84E-01	40
	Metabolism	8	2,116	0.96	9.62E-01	24
	Metabolism of proteins	5	2,111	1.00	9.98E-01	4
	Immune System	4	2,226	1.00	1.00E+00	15
	GPCR downstream signalling	3	1,146	0.98	9.77E-01	4
<i>Argulus</i>	Signal Transduction	7	2,738	1.00	9.99E-01	27
	Metabolism of proteins	6	2,111	0.99	9.91E-01	31
	Metabolism	5	2,116	1.00	9.97E-01	7
	Immune System	5	2,226	1.00	9.98E-01	21
	Gene expression (Transcription)	3	1,416	0.99	9.92E-01	5

3.2 Protein identification and comparisons

Analysis of proteins identified using the killifish transcriptome identified a similar number of proteins released (MASCOT ≥ 50) by all experimental groups (Table S1). Actin family beta (for cell motility, structure and integrity) and fish keratin (from host tissue) proteins were expressed in all treatment groups. OSER1, a protein with roles in oxidative stress response, was abundant in all fish water samples, but not in *Argulus* only samples. Despite this, no significant variations in MASCOT scores between control and infected water samples were identified for OSER1 ($t_{6,96}=1.521$, $P=0.172$). In addition to proteins detected from the killifish transcriptome, results were searched against the NCBI database to identify proteins from fish and parasite origins. Of the forty-seven proteins with a MASCOT score on or above the 59 cut off, 49% were deemed as human contamination (Table S3, in red) but 13% of proteins included claw keratin and proteins associated with teleost fish (polyserase-2-like proteins and Chymotrypsin-like protease proteins).

Variations between infected and uninfected fish samples were identified for both transcriptome and NCBI data sets. Proteins identified from the transcriptome indicated that Keratin 94 (of fish origin), associated with host tissues, was only identified in infected fish water samples. However, these samples did not differ significantly when MASCOT scores from each group were compared ($t_{6,9}=0.88$, $P=0.396$). Zinc finger proteins (a class of protein involved with nucleic acid binding and protein dimerization) were highly abundant in DAN infected and *Argulus* only samples.

Gene ontology analysis of proteins from the transcriptome was conducted and compared between treatments (Figure 2). This analysis indicated that proteins involved in pigmentation (TSPAN36 and SLC45A2) and binding were only detected among infected fish and no protein group was identified solely in the control groups. Previous work has demonstrated links

1
2
3 between infection and pigmentation (55, 56), therefore, proteins related to colour in *K.*
4 *marmoratus* are likely promising indicators of immune response. All groups contained proteins
5 related to metabolism, proteolysis and circulation. Finally, proteins for sound perception,
6 cilium movement and mucus production (Figure 2A) were only identified in water from fish
7 origin. Statistical analysis of gene ontology data from NCBI indicated that a significantly
8 greater abundance of proteins for methylation were identified in infected fish water samples
9 when compared with controls ($t_5 = 2.397$, $P=0.042$) as can be seen in figure 2. Conversely
10 despite variation indicated in figure 2, the number of proteins with roles in immune response
11 did not differ significantly between groups ($t_5 = -0.682$, $P=0.522$). For control groups,
12 functional proteins identified were involved in apoptosis and autophagy. Lastly protein families
13 observed in all groups included extracellular proteins, as well as proteins involved in the
14 immune response, and in carbohydrate binding (for all see Figure 2A).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Go annotation of all proteins identified from infected (with *Argulus foliaceus*), uninfected and parasite only water samples for two lines (DAN and R) of *Kryptolebias marmoratus* transcriptome (A) and NCBI data (B).

4. Discussion

Current, non-lethal fish health monitoring involves collection and subsequent analysis of the external surface mucus of teleost species (57, 58). Fish skin mucosa acts as an important barrier against pathogens (59, 60) and within this barrier proteins may be secreted, synthesized and released (in the form of dead cells) leading to the production of identifiable immune related molecules (61). Release of molecules into the surrounding aquatic medium is thought to act as chemical stimulus that enable teleost fish to assess the health, genetic background and identity of conspecifics (11, 15). The methods used in the current study indicate that these proteins can be extracted, purified and characterized from teleost water samples and have the potential to inform infection status non-invasively, without the need for invasive sampling. In the current study, initial gel based methods demonstrated a number of observable protein bands that may be indicative of heat stress and heat shock proteins (62, 63), binding proteins (64) and keratins (65). Furthermore, gel-free methods following trypsin digestion of samples greatly increased the number of proteins identified and confirmed the presence of fish keratins, binding proteins and immune related molecules.

Skin extracts are thought to be present in the mucosa following skin rupture, leading to the release of alarm cues, identifiable by conspecifics (59, 66). Fish keratin proteins (and many actin proteins below the MASCOT cut-off) were identified in samples from infected fish only; these proteins are likely to originate from the skin surface (due to the structural function of keratin) in the form of ruptured or dead cells (67). Additionally, keratins from trout have been shown to display antibacterial activity (68) and these proteins will likely indicate danger to surrounding conspecifics. In addition to keratins, binding proteins were highly expressed in the current water samples from infected R line fish. Binding proteins, identified in the mucosa of fish have been shown to serve many functions including immune response and lipid metabolism (69, 70) and it is plausible that these proteins from the mucus are being detected in

1
2
3 current samples. Importantly, fish mucus has been shown to contain a plethora of immune
4 related proteins including those from the complement (71), immunoglobulins (72) and
5 antimicrobial peptides (73). Both NCBI and transcriptome based databases indicated immune
6 related proteins in current fish water samples; these were identified for individual characterized
7 proteins (e.g. cytokines) but also at the level of go annotation and pathway analyses whereby
8 immune annotations and pathways were highly enriched. One current example of an immune
9 protein identified by MASCOT and go annotation scoring, from infected R line individuals, is
10 interleukin 6 signal transducer (IL6st). Interleukin 6 receptor was previously identified in the
11 mucus of gilthead sea bream (*Sparus aurata*) that were chronically stressed (74) and several
12 other interleukin molecules with well-known roles in inflammatory response and protection
13 from pathogens have been described in the skin of Nile tilapia (*Oreochromis niloticus*) (75,
14 76). As fish mucus is continually secreted and replaced (75) it is likely that immune molecules,
15 such as IL6, are transferred to the media on a constant basis. As demonstrated in the current
16 study, the regeneration and movement of mucosal molecules into the aquatic media allows for
17 collection and analysis of these proteins in low water volumes.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38 Further to specific immune or structural proteins, the fish mucosa of chronically stressed has
39 allowed for research into a variety of stress related proteins thought to influence the response
40 of fish to environmental fluctuations (67, 74). Proteins associated with oxidative stress
41 response are commonly studied in fish exposed to toxic environmental pollutants, for example
42 metals (77) and associated biomarkers have been suggested as a potential target for
43 environmental monitoring and pollutant detection (78, 79). Additionally, these proteins have
44 been identified in fish skin mucus and have roles in regulation of cell death (74, 80). Here we
45 provide evidence that the OSER1 protein, of fish origin, was detectable from all samples of
46 aquaria water and has the potential indicate environmental stress response in this species. In
47 this instance OSER1 could be linked with stress related to experimental testing such as sham
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 infection (for control fish) or louse attachment for infected individuals. This warrants further
4 investigation but confirms the ability of current methods to detect a well-known marker.
5
6
7

8 Many more proteins of interest appeared in two or three individual fish below the MASCOT
9 cut-off. Presence of these proteins might have increased with a larger number of samples at a
10 higher concentration and/or larger volumes of water. Briefly, infected fish samples and *Argulus*
11 only samples contained proteins (below the MASCOT significance cut off) from mite and lice
12 species; these included PHUM400690 (for immune response to ticks), capicua protein (a
13 transcriptional repressor) and IscW (an immune suppressor). As only one parasite and a small
14 host were tested in the current study it is likely that larger fish, with several parasites would
15 reveal more clues about these proteins. If these proteins were discovered to increase with
16 parasite number they would likely be a strong target for environmental protein detection
17 methods. Proteins with roles in oxidative stress response, response to lice, pigmentation and
18 immune response identified here have the potential to reveal infection and/or stress status in
19 fish. Proteins associated with *Argulus foliaceus* were observed in water from infected fish, yet
20 several proteins were only identified in the *Argulus* control group. These proteins (zinc finger
21 protein and proteins involved in the immune response to lice and ticks) demonstrated the
22 possibility of using proteomic techniques to detect the presence of parasitic organisms within
23 the environment, as with environmental DNA (81-83). Lastly proteins with roles in oxidative
24 stress response, response to lice, pigmentation and immune response identified here have the
25 potential to indicate infection and/or stress status in fish, an area that requires further
26 investigation. We propose that the proteomic methods outlined here have the potential to
27 provide a similar tool for evaluating stress and health status in individuals, with the added
28 benefit of functional analysis of expressed proteins. Lastly, we suggest that proteomic and
29 environmental DNA methods could be used in tandem to gain non-invasive data on farmed and
30 wild fish populations.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

5. Concluding remarks

This is the first study outlining protein characterisation from brackish water samples without the need for gel electrophoresis and represents a major step forward in the field of metaproteomic techniques. Our novel assay provides a non-invasive method for detecting infection status in fish, which will be useful for assessing the health condition of both wild and farmed aquatic organisms. With further investigation and optimisation, protein detection methods represent a promising new method for environmental monitoring and could become an important tool in applied and aquatic biology.

Acknowledgments

This study was supported by a Natural Environment Research Council Industrial CASE studentship (NE/L00948X) and the BCHC (Aberystwyth University) Access Fellowship to RJP.

Author's contributions

RJP, SC & RS designed the study with help from PMB and RMM. RJP, SC & RMM wrote the manuscript. PMB provided the lab for experimental procedures. JC provided expertise and facilities for all experimental infections. CGL and JC provided manuscript corrections and advise.

Conflict of interest statement

The authors have declared no conflict of interest

Ethics approval

All the experiments in this study have been conducted following Home Office regulations, approved by both Swansea and Cardiff University Ethics Committees and under Home Office licence number PPL 302357.

SUPPORTING INFORMATION

Supplementary material_Proteomics.docx

1. **Details S1.** Extra details for methods and materials – sample processing, 1D SDS page, TCA methods.
2. **Table S1.** Protein number, percentage above mascot cut off, unique proteins and unidentified proteins from all groups and all fish replicates from both Transcriptome and NCBI data sets.
3. **Table S2.** Proteins identified on or equal to mascot score cut off, proteins found in all replicates and proteins for in two replicates from data obtain from the Mangrove Killifish Transcriptome
4. **Table S3.** Proteins identified on or equal to mascot score cut off, proteins found in all replicates and proteins for in two replicates from data obtain from the NCBI database
5. **Table S4.** Proteins of interest identified from Mangrove killifish transcriptome data and the NCBI database
6. **Table S5.** REACTOME pathway analysis for 5 groups of water samples from *Kryptolebias marmoratus*
7. **Figure S1:** Go annotations from transcriptome data (Level Go All) for all 5 groups of *Kryptolebias.marmoratus* (Dan infected, Dan control, R infected, R control and *Argulus* only) and all three replicates per group.
8. **Figure S2:** Go annotations from NCBI data (Level Go All) for all 5 groups of *Kryptolebias marmoratus* (Dan infected, Dan control, R infected, R control and *Argulus* only) and all three replicates from each group.
9. **Figure S3.** Pathway analysis using REACTOME for 3 biological replicates of *Kryptolebias marmoratus* water samples from 5 groups.

References

1. Abreu, M. S.; Giacomini, A. C. V.; Gusso, D.; Koakoski, G.; Oliveira, T. A.; Marqueze, A.; Barreto, R. E.; Barcellos, L. J., Behavioral responses of zebrafish depend on the type of threatening chemical cues. *Journal of Comparative Physiology A* **2016**, 202, (12), 895-901.
2. Wyatt, T. D., Proteins and peptides as pheromone signals and chemical signatures. *Animal Behaviour* **2014**, 97, 273-280.
3. Brown, G. E., Learning about danger: chemical alarm cues and local risk assessment in prey fishes. *Fish and Fisheries* **2003**, 4, (3), 227-234.
4. Chivers, D. P.; Mirza, R. S.; Johnston, J. G., Learned recognition of heterospecific alarm cues enhances survival during encounters with predators. *Behaviour* **2002**, 139, (7), 929-938.
5. Sharp, J. G.; Garnick, S.; Elgar, M. A.; Coulson, G. In *Parasite and predator risk assessment: nuanced use of olfactory cues*, Proc. R. Soc. B, 2015; *The Royal Society*: **2015**; p 20151941.
6. Olsson, M. J.; Lundström, J. N.; Kimball, B. A.; Gordon, A. R.; Karshikoff, B.; Hosseini, N.; Sorjonen, K.; Olgart Höglund, C.; Solares, C.; Soop, A., The scent of disease: human body odor contains an early chemosensory cue of sickness. *Psychological science* **2014**, 25, (3), 817-823.
7. Ehman, K.; Scott, M., Urinary odour preferences of MHC congenic female mice, *Mus domesticus*: implications for kin recognition and detection of parasitized males. *Animal Behaviour* **2001**, 62, (4), 781-789.
8. Lehtonen, T. K.; Kvarnemo, C., Odour cues from suitors' nests determine mating success in a fish. *Biology letters* **2015**, 11, (5), 20150021.
9. Stone, C. F.; Moore, J., Parasite-induced alteration of odour responses in an amphipod–acanthocephalan system. *International journal for parasitology* **2014**, 44, (13), 969-975.

- 1
2
3 10. Webster, M.; Atton, N.; Ward, A.; Hart, P., Turbidity and foraging rate in threespine
4 sticklebacks: the importance of visual and chemical prey cues. *Behaviour* **2007**, 144, (11),
5 1347-1360.
6
7
8
9
10 11. Boehm, T.; Zufall, F., MHC peptides and the sensory evaluation of genotype. *Trends*
11 *in neurosciences* **2006**, 29, (2), 100-107.
12
13
14 12. Charpentier, M.; Prugnolle, F.; Gimenez, O.; Widdig, A., Genetic heterozygosity and
15 sociality in a primate species. *Behavior Genetics* **2008**, 38, (2), 151-158.
16
17
18 13. Milinski, M.; Griffiths, S.; Wegner, K. M.; Reusch, T. B. H.; Haas-Assenbaum, A.;
19 Boehm, T., Mate choice decisions of stickleback females predictably modified by MHC
20 peptide ligands. *Proceedings of the National Academy of Sciences of the United States of*
21 *America* **2005**, 102, (12), 4414-4418.
22
23
24
25
26
27 14. Consuegra, S.; Garcia de Leaniz, C., MHC-mediated mate choice increases parasite
28 resistance in salmon. *Proceedings of the Royal Society B: Biological Sciences* **2008**, 275,
29 (1641), 1397-1403.
30
31
32
33
34
35 15. Olsén, K. H.; Grahn, M.; Lohm, J., Influence of MHC on sibling discrimination in
36 Arctic char, *Salvelinus alpinus* (L.). *Journal of chemical ecology* **2002**, 28, (4), 783-795.
37
38
39
40 16. Beynon, R. J.; Hurst, J., Multiple roles of major urinary proteins in the house mouse,
41 *Mus domesticus*. *Biochemical Society Transactions*: **2003**, 142-146.
42
43
44 17. Fischer, E. K.; Harris, R. M.; Hofmann, H. A.; Hoke, K. L., Predator exposure alters
45 stress physiology in guppies across timescales. *Hormones and behavior* **2014**, 65, (2), 165-172.
46
47
48 18. Mathuru, A. S.; Kibat, C.; Cheong, W. F.; Shui, G.; Wenk, M. R.; Friedrich, R. W.;
49 Jesuthasan, S., Chondroitin fragments are odorants that trigger fear behavior in fish. *Current*
50 *Biology* **2012**, 22, (6), 538-544.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 19. Roberts, L. J.; Garcia de Leaniz, C., Something smells fishy: predator-naïve salmon use
4 diet cues, not kairomones, to recognize a sympatric mammalian predator. *Animal Behaviour*
5 **2011**, 82, (4), 619-625.
6
7
8
9
10 20. Barreto, R. E.; Júnior, A. B.; Giassi, A. C. C.; Hoffmann, A., The ‘club’ cell and
11 behavioural and physiological responses to chemical alarm cues in the Nile tilapia. *Marine and*
12 *Freshwater Behaviour and Physiology* **2010**, 43, (1), 75-81.
13
14
15
16
17 21. Forlenza, M.; Walker, P. D.; De Vries, B. J.; Bonga, S. E. W.; Wiegertjes, G. F.,
18 Transcriptional analysis of the common carp (*Cyprinus carpio* L.) immune response to the fish
19 louse *Argulus japonicus* Thiele (Crustacea: Branchiura). *Fish & shellfish immunology* **2008**,
20 25, (1-2), 76-83.
21
22
23
24
25
26 22. Fast, M. D., Fish immune responses to parasitic copepod (namely sea lice) infection.
27 *Developmental & Comparative Immunology* **2014**, 43, (2), 300-312.
28
29
30
31 23. Walker, P. D.; Flik, G.; Bonga, S. W., The biology of parasites from the genus *Argulus*
32 and a review of the interactions with its host. *Host-parasite interactions* **2004**, 55, 107-129.
33
34
35
36 24. Bandilla, M.; Valtonen, E.; Suomalainen, L.-R.; Aphalo, P.; Hakalahti, T., A link
37 between ectoparasite infection and susceptibility to bacterial disease in rainbow trout.
38 *International journal for parasitology* **2006**, 36, (9), 987-991.
39
40
41
42 25. Ángeles Esteban, M., An overview of the immunological defenses in fish skin. *ISRN*
43 *immunology* **2012**, 1-29..
44
45
46
47 26. McGuckin, M. A.; Lindén, S. K.; Sutton, P.; Florin, T. H., Mucin dynamics and enteric
48 pathogens. *Nature Reviews Microbiology* **2011**, 9, (4), 265.
49
50
51
52 27. Ellis, A., Immunity to bacteria in fish. *Fish & shellfish immunology* **1999**, 9, (4), 291-
53 308.
54
55
56 28. Sarmaşık, A., Antimicrobial peptides: a potential therapeutic alternative for the
57 treatment of fish diseases. *Turkish Journal of Biology* **2002**, 26, (4), 201-207.
58
59
60

- 1
2
3 29. Ingram, G., Substances involved in the natural resistance of fish to infection—a review.
4
5 *Journal of Fish Biology* **1980**, 16, (1), 23-60.
6
7
8 30. González-Chávez, S. A.; Arévalo-Gallegos, S.; Rascón-Cruz, Q., Lactoferrin: structure,
9
10 function and applications. *International journal of antimicrobial agents* **2009**, 33, (4), 301. e1-
11
12 301. e8.
13
14 31. Zhang, Y.-A.; Salinas, I.; Li, J.; Parra, D.; Bjork, S.; Xu, Z.; LaPatra, S. E.;
15
16 Bartholomew, J.; Sunyer, J. O., IgT, a primitive immunoglobulin class specialized in mucosal
17
18 immunity. *Nature immunology* **2010**, 11, (9), 827.
19
20
21 32. D'Ambrosio, C.; Arena, S.; Talamo, F.; Ledda, L.; Renzone, G.; Ferrara, L.; Scaloni,
22
23 A., Comparative proteomic analysis of mammalian animal tissues and body fluids: bovine
24
25 proteome database. *Journal of chromatography B* **2005**, 815, (1-2), 157-168.
26
27
28 33. Zargar, S.; Gupta, N.; Mir, R.; Rai, V., Shift from gel based to gel free proteomics to
29
30 unlock unknown regulatory network in plants: a comprehensive review. *Journal of*
31
32 *Advanced Research Biotech* **2016**, 1, 19.
33
34
35 34. Brophy, P. M.; Mackintosh, N.; Morpew, R. M., Anthelmintic metabolism in parasitic
36
37 helminths: proteomic insights. *Parasitology* **2012**, 139, (9), 1205-1217.
38
39
40 35. Marcilla, A.; Trelis, M.; Cortés, A.; Sotillo, J.; Cantalapiedra, F.; Minguéz, M. T.;
41
42 Valero, M. L.; Del Pino, M. M. S.; Muñoz-Antoli, C.; Toledo, R., Extracellular vesicles from
43
44 parasitic helminths contain specific excretory/secretory proteins and are internalized in
45
46 intestinal host cells. *PloS one* **2012**, 7, (9), e45974.
47
48
49 36. Robinson, M. W.; Menon, R.; Donnelly, S. M.; Dalton, J. P.; Ranganathan, S., An
50
51 integrated transcriptomics and proteomics analysis of the secretome of the helminth pathogen
52
53 *Fasciola hepatica* proteins associated with invasion and infection of the mammalian host.
54
55 *Molecular & Cellular Proteomics* **2009**, 8, (8), 1891-1907.
56
57
58
59
60

- 1
2
3 37. Morphey, R. M.; Wright, H. A.; LaCourse, E. J.; Woods, D. J.; Brophy, P. M.,
4 Comparative proteomics of excretory-secretory proteins released by the liver fluke *Fasciola*
5 *hepatica* in sheep host bile and during in vitro culture ex host. *Molecular & Cellular*
6 *Proteomics* **2007**, 6, (6), 963-972.
7
8
9
10
11
12 38. Hutson, K. S.; Cable, J.; Grutter, A. S.; Paziewska-Harris, A.; Barber, I., Aquatic
13 Parasite Cultures and Their Applications. *Trends in parasitology* **2018**.
14
15
16
17 39. Wöhlbrand, L.; Feenders, C.; Nachbaur, J.; Freund, H.; Engelen, B.; Wilkes, H.;
18 Brumsack, H. J.; Rabus, R., Impact of Extraction Methods on the Detectable Protein
19 Complement of Metaproteomic Analyses of Marine Sediments. *Proteomics* **2017**, 17, (22),
20 1700241.
21
22
23
24
25
26 40. Lee, J. S.; Raisuddin, S.; Schlenk, D., *Kryptolebias marmoratus* (Poey, 1880): a
27 potential model species for molecular carcinogenesis and ecotoxicogenomics. *Journal of Fish*
28 *Biology* **2008**, 72, (8), 1871-1889.
29
30
31
32
33 41. Harrington Jr, R. W.; Kallman, K. D., The homozygosity of clones of the self-fertilizing
34 hermaphroditic fish *Rivulus marmoratus* Poey (Cyprinodontidae, Atheriniformes). *The*
35 *American Naturalist* **1968**, 102, (926), 337-343.
36
37
38
39
40 42. Pasternak, A. F.; Mikheev, V. N.; Valtonen, E. T. In *Life history characteristics of*
41 *Argulus foliaceus* L.(Crustacea: Branchiura) populations in Central Finland, *Annales*
42 *Zoologici Fennici*, 2000; JSTOR: 2000; pp 25-35.
43
44
45
46
47 43. Bower-Shore, C., An investigation of the common fish louse, *Argulus foliaceus* (Linn.).
48 *Parasitology* **1940**, 32, (4), 361-371.
49
50
51
52 44. Mirzaei, M.; Khovand, H., Prevalence of *Argulus foliaceus* in ornamental fishes
53 [goldfish (*Carassius auratus*) and Koi (*Cyprinus carpio*)] in Kerman, southeast of Iran. *Journal*
54 *of parasitic diseases* **2015**, 39, (4), 780-782.
55
56
57
58
59
60

- 1
2
3 45. Aalberg, K.; Koščová, L.; Šmiga, L.; Košuth, P.; Koščo, J.; Oros, M.; Barčák, D.; Lazar,
4
5 P., A study of fish lice (*Argulus* sp.) infection in freshwater food fish. *Folia Veterinaria* **2016**,
6
7 60, (3), 54-59.
8
9
10 46. Ellison, A.; Jones, J.; Inchley, C.; Consuegra, S., Choosy males could help explaining
11
12 androdioecy in a selfing fish. *American Naturalist* **2013**, 181, 855-862.
13
14
15 47. Stewart, A.; Jackson, J.; Barber, I.; Eizaguirre, C.; Paterson, R.; van West, P.; Williams,
16
17 C.; Cable, J., Hook, line and infection: a guide to culturing parasites, establishing infections
18
19 and assessing immune responses in the three-spined stickleback. In *Advances in parasitology*,
20
21 Elsevier: 2017; Vol. 98, pp 39-109.
22
23
24 48. Pawluk, R. J.; Uren Webster, T. M.; Cable, J.; Garcia de Leaniz, C.; Consuegra, S.,
25
26 Immune-related transcriptional responses to parasitic infection in a naturally inbred fish: roles
27
28 of genotype and individual variation. *Genome Biology and Evolution* **2018**, 319-327.
29
30
31 49. Bradford, M. M., A rapid and sensitive method for the quantitation of microgram
32
33 quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*
34
35 **1976**, 72, (1-2), 248-254.
36
37
38 50. Shevchenko, A.; Wilm, M.; Vorm, O.; Mann, M., Mass spectrometric sequencing of
39
40 proteins from silver-stained polyacrylamide gels. *Analytical chemistry* **1996**, 68, (5), 850-858.
41
42
43 51. Vizcaíno, J. A.; Csordas, A.; Del-Toro, N.; Dianes, J. A.; Griss, J.; Lavidas, I.; Mayer,
44
45 G.; Perez-Riverol, Y.; Reisinger, F.; Ternent, T., 2016 update of the PRIDE database and its
46
47 related tools. *Nucleic acids research* **2015**, 44, (D1), D447-D456.
48
49
50 52. Dennis, G.; Sherman, B. T.; Hosack, D. A.; Yang, J.; Gao, W.; Lane, H. C.; Lempicki,
51
52 R. A., DAVID: database for annotation, visualization, and integrated discovery. *Genome*
53
54 *biology* **2003**, 4, (9), R60.
55
56
57
58
59
60


- 1
2
3 53. Fabregat, A.; Sidiropoulos, K.; Garapati, P.; Gillespie, M.; Hausmann, K.; Haw, R.;
4 Jassal, B.; Jupe, S.; Korninger, F.; McKay, S., The reactome pathway knowledgebase. *Nucleic*
5 *acids research* **2015**, 44, (D1), D481-D487.
6
7
8
9
10 54. Team, R. C., R: A language and environment for statistical computing. **2013**, 201.
11
12 55. Price, A. C.; Weadick, C. J.; Shim, J.; Rodd, F. H., Pigments, patterns, and fish
13 behavior. *Zebrafish* **2008**, 5, (4), 297-307.
14
15
16 56. Seppälä, O.; Karvonen, A.; Valtonen, E. T., Impaired crypsis of fish infected with a
17 trophically transmitted parasite. *Animal Behaviour* **2005**, 70, (4), 895-900.
18
19
20 57. Bergsson, G.; Agerberth, B.; Jörnvall, H.; Gudmundsson, G. H., Isolation and
21 identification of antimicrobial components from the epidermal mucus of Atlantic cod (*Gadus*
22 *morhua*). *The FEBS journal* **2005**, 272, (19), 4960-4969.
23
24
25 58. Salles, C.; Gagliano, P.; Leitão, S.; Salles, J.; Guedes, H.; Cassano, V.; De-Simone, S.
26 G., Identification and characterization of proteases from skin mucus of tambacu, a Neotropical
27 hybrid fish. *Fish physiology and biochemistry* **2007**, 33, (2), 173.
28
29
30 59. Reverter, M.; Tapissier-Bontemps, N.; Lecchini, D.; Banaigs, B.; Sasal, P., Biological
31 and Ecological Roles of External Fish Mucus: A Review. *Fishes* **2018**, 3, (4), 41.
32
33
34 60. Gomez, D.; Sunyer, J. O.; Salinas, I., The mucosal immune system of fish: the evolution
35 of tolerating commensals while fighting pathogens. *Fish & shellfish immunology* **2013**, 35, (6),
36 1729-1739.
37
38
39 61. Brinchmann, M. F., Immune relevant molecules identified in the skin mucus of fish
40 using-omics technologies. *Molecular BioSystems* **2016**, 12, (7), 2056-2063.
41
42
43 62. Iwama, G. K.; Vijayan, M. M.; Forsyth, R. B.; Ackerman, P. A., Heat shock proteins
44 and physiological stress in fish. *American Zoologist* **1999**, 39, (6), 901-909.
45
46
47 63. Kiang, J. G.; Tsokos, G. C., Heat shock protein 70 kDa: molecular biology,
48 biochemistry, and physiology. *Pharmacology & therapeutics* **1998**, 80, (2), 183-201.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 64. Siharath, K.; Kelley, K. M.; Bern, H. A., A low-molecular-weight (25-kDa) IGF-
4 binding protein is increased with growth inhibition in the fasting striped bass, *Morone saxatilis*.
5
6
7
8 *General and comparative endocrinology* **1996**, 102, (3), 307-316.
9
- 10 65. Conrad, M.; Lemb, K.; Schubert, T.; Markl, J., Biochemical identification and tissue-
11 specific expression patterns of keratins in the zebrafish *Danio rerio*. *Cell and tissue research*
12
13 **1998**, 293, (2), 195-205.
14
- 15 66. Chivers, D. P.; Zhao, X.; Ferrari, M. C., Linking morphological and behavioural
16 defences: prey fish detect the morphology of conspecifics in the odour signature of their
17 predators. *Ethology* **2007**, 113, (8), 733-739.
18
- 19 67. Jurado, J.; Fuentes-Almagro, C. A.; Guardiola, F. A.; Cuesta, A.; Esteban, M. Á.;
20 Prieto-Álamo, M.-J., Proteomic profile of the skin mucus of farmed gilthead seabream (*Sparus*
21 *aurata*). *Journal of proteomics* **2015**, 120, 21-34.
22
- 23 68. Molle, V.; Campagna, S.; Bessin, Y.; Ebran, N.; Saint, N.; Molle, G., First evidence of
24 the pore-forming properties of a keratin from skin mucus of rainbow trout (*Oncorhynchus*
25 *mykiss*, formerly *Salmo gairdneri*). *Biochemical Journal* **2008**, 411, (1), 33-40.
26
- 27 69. Rajan, B.; Fernandes, J. M.; Caipang, C. M.; Kiron, V.; Rombout, J. H.; Brinchmann,
28 M. F., Proteome reference map of the skin mucus of Atlantic cod (*Gadus morhua*) revealing
29 immune competent molecules. *Fish & shellfish immunology* **2011**, 31, (2), 224-231.
30
- 31 70. Cordero, H.; Brinchmann, M. F.; Cuesta, A.; Meseguer, J.; Esteban, M. A., Skin mucus
32 proteome map of European sea bass (*Dicentrarchus labrax*). *Proteomics* **2015**, 15, (23-24),
33 4007-4020.
34
- 35 71. Shen, Y.; Zhang, J.; Xu, X.; Fu, J.; Li, J., Expression of complement component C7
36 and involvement in innate immune responses to bacteria in grass carp. *Fish & shellfish*
37 *immunology* **2012**, 33, (2), 448-454.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 72. Salinas, I.; Zhang, Y.-A.; Sunyer, J. O., Mucosal immunoglobulins and B cells of
4 teleost fish. *Developmental & Comparative Immunology* **2011**, 35, (12), 1346-1365.
5
6
7
8 73. Zhang, X.-J.; Wang, P.; Zhang, N.; Chen, D.-D.; Nie, P.; Li, J.-L.; Zhang, Y.-A., B cell
9 Functions can Be Modulated by antimicrobial Peptides in rainbow Trout *Oncorhynchus*
10 *mykiss*: novel insights into the innate nature of B cells in Fish. *Frontiers in immunology* **2017**,
11 8, 388.
12
13
14
15
16
17 74. Pérez-Sánchez, J.; Terova, G.; Simó-Mirabet, P.; Rimoldi, S.; Folkedal, O.; Caldach-
18 Giner, J. A.; Olsen, R. E.; Sitjà-Bobadilla, A., Skin mucus of gilthead sea bream (*Sparus aurata*
19 L.). Protein mapping and regulation in chronically stressed fish. *Frontiers in physiology* **2017**,
20 8, 34.
21
22
23
24
25
26 75. Esteban, M. Á.; Cerezuela, R., Fish mucosal immunity: skin. In *Mucosal Health in*
27 *Aquaculture*, Elsevier: **2015**; pp 67-92.
28
29
30
31 76. Wei, X.; Li, B.; Wu, L.; Yin, X.; Zhong, X.; Li, Y.; Wang, Y.; Guo, Z.; Ye, J.,
32 Interleukin-6 gets involved in response to bacterial infection and promotes antibody production
33 in Nile tilapia (*Oreochromis niloticus*). *Developmental & Comparative Immunology* **2018**, 89,
34 141-151.
35
36
37
38
39
40 77. Valavanidis, A.; Vlahogianni, T.; Dassenakis, M.; Scoullou, M., Molecular biomarkers
41 of oxidative stress in aquatic organisms in relation to toxic environmental pollutants.
42 *Ecotoxicology and environmental safety* **2006**, 64, (2), 178-189.
43
44
45
46
47 78. Pandey, S.; Parvez, S.; Sayeed, I.; Haque, R.; Bin-Hafeez, B.; Raisuddin, S.,
48 Biomarkers of oxidative stress: a comparative study of river Yamuna fish *Wallago attu* (Bl. &
49 Schn.). *Science of the total environment* **2003**, 309, (1-3), 105-115.
50
51
52
53
54 79. McCarthy, J. F.; Shugart, L. R., *Biomarkers of environmental contamination*. United
55 States: N. p., **1990**.
56
57
58
59
60

- 1
2
3 80. Dzul-Caamal, R.; Salazar-Coria, L.; Olivares-Rubio, H. F.; Rocha-Gómez, M. A.;
4
5 Girón-Pérez, M. I.; Vega-López, A., Oxidative stress response in the skin mucus layer of
6
7 *Goodea gracilis* (Hubbs and Turner, 1939) exposed to crude oil: A non-invasive approach.
8
9 *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **2016**,
10
11 200, 9-20.
12
13
14 81. Thomsen, P. F.; Kielgast, J. O. S.; Iversen, L. L.; Wiuf, C.; Rasmussen, M.; Gilbert, M.
15
16 T. P.; Orlando, L.; Willerslev, E., Monitoring endangered freshwater biodiversity using
17
18 environmental DNA. *Molecular Ecology* **2012**, 21, (11), 2565-2573.
19
20
21 82. Ficetola, G. F.; Miaud, C.; Pompanon, F.; Taberlet, P., Species detection using
22
23 environmental DNA from water samples. *Biology Letters* **2008**, 4, (4), 423-425.
24
25
26 83. Robinson, C. V.; Uren Webster, T. M.; Cable, J.; James, J.; Consuegra, S.,
27
28 Simultaneous detection of invasive signal crayfish, endangered white-clawed crayfish and the
29
30 crayfish plague pathogen using environmental DNA. *Biological Conservation* **2018**, (222),
31
32 241-252.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For TOC only



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60