

Aberystwyth University

Effects of rumen bacterial lipases on ruminal lipid metabolism

Williams, Cate

Publication date:
2016

Citation for published version (APA):

Williams, C. (2016). *Effects of rumen bacterial lipases on ruminal lipid metabolism*. Poster session presented at INRA - ROWETT 2016 , Clermont-Ferrand , France.

General rights


Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk



Effects of rumen bacterial lipases on ruminal lipid metabolism

CL Williams^a, ND Scollan^a, JKS Tweed^a, H Vallin^a, L Davies^b and SA Huws^a.

^aInstitute of Biological, Environmental and Rural Sciences, Aberystwyth University, Wales, UK, SY23 3DA
^bHybu Cig Cymru – Meat Promotion Wales, Tŷ Rheidol, Parc Merlin, Aberystwyth, SY23 3FF.
E-mail: clw10@aber.ac.uk

Introduction

The burden of chronic, fat-related diseases upon the NHS is reaching a critical level and can be linked to an increased intake of saturated fats (SFA). SFAs are found in many products, including red meat and milk despite the ruminant diet being rich in health-beneficial poly-unsaturated fatty acids^{1,2}. This is due to lipolysis and biohydrogenation (fig 1), processes that saturate PUFA in the rumen to protect bacteria from PUFA's toxic double bonds^{1,2,3}. Enhancing lipolysis could decrease biohydrogenation due to the presence of higher levels of PUFA.

Study Aims

To determine the level of PUFA required to inhibit biohydrogenation *in vitro*.




Fig.1: The biohydrogenation process in the rumen, LA-c9c12-18:2; LNA-c9c12c15-18:3; VA-t11-18:1. CLA-conjugated linoleic acid; LA- linoleic acid; LNA- linolenic acid; VA-vaccenic acid².

Materials and Methods

A batch culture experiment with 6 treatments was used, each with a basal level of freeze-dried silage and: control (No LA/LNA) or LA/LNA at 50µM and 250µM, 500µM, 750µM or 1µM concentrations. These were incubated under rumen like conditions⁴ and destructively harvested in triplicate at four time points (0, 4, 8 and 24h). Fatty acid profiles were assessed using FAME analysis and gas chromatography. RNA was extracted, reverse transcribed and a restriction digest carried out using HaeIII before TRFLP to assess the effects on the rumen microbiota.

Acknowledgements

This research is funded by the European Social Fund (ESF) through the Welsh government's European Union Convergence programme. The study was also funded in part by Hybu Cig Cymru.

Results

- LNA and LA at 250µM caused a significant decrease in 18:0 at 4 to 8 h ($P=0.013$) of 16.7 and 11.5mg/g-1 ($P=0.067$) respectively (Table 1 & 2).
- LA and LNA also caused significant increases in 18:3n-3 at 4 hours (increase of 1.98mg/g⁻¹), and changes in 18:2n-6 were significant ($P=>0.05$) at 4-8 hours (up to 2.12mg/g⁻¹) (Table 1 & 2).
- Concentrations of 250µM and 500µM LA/LNA were most effective in reducing 18:0 and increasing PUFA content (Table 1 & 2).
- TRFLP dendrograms showed no significant changes in the bacterial population relative to time or LA/LNA concentration (data not shown).

Table 1. Effects of linoleic acid addition on fatty acid concentrations *in vitro*

Time	Fatty Acid	linoleic acid Concentration (µM)					SED	P
		0	50	250	500	750		
0 hrs	18:0	41.1 ^a	54.8 ^a	52.3 ^a	48.1 ^a	37.5 ^a	41.9 ^a	0.240
	18:1 trans-11	5.96 ^b	4.84 ^a	4.95 ^a	4.77 ^a	4.13 ^a	4.48 ^a	0.037
	18:2 trans-9, trans-11	0.75 ^a	0.64 ^a	0.69 ^a	0.51 ^a	1.37 ^a	0.88 ^a	0.451
	18:2 cis-9, cis-12	0.01 ^a	0.02 ^a	0.01 ^a	0.01 ^a	0.02 ^a	0.00 ^a	0.537
	18:3 n-3	0.29 ^a	0.02 ^a	0.00 ^a	0.03 ^a	0.44 ^a	0.07 ^a	0.273
4 hrs	18:0	54.7 ^b	49.9 ^b	40.8 ^a	43.0 ^a	50.6 ^b	49.7 ^b	0.067
	18:1 trans-11	3.94 ^b	3.91 ^b	3.65 ^b	3.36 ^a	3.75 ^b	4.16 ^b	0.087
	18:2 trans-9, trans-11	0.09 ^b	0.08 ^b	0.06 ^a	0.07 ^b	0.08 ^b	0.09 ^b	0.133
	18:2 cis-9, cis-12	0.96 ^a	0.95 ^a	2.13 ^c	1.64 ^c	1.15 ^b	0.77 ^a	0.001
	18:3 n-3	1.35 ^b	1.19 ^a	2.27 ^c	1.73 ^b	1.41 ^b	1.06 ^a	<.001
8 hrs	18:0	51.1 ^a	50.8 ^a	44.8 ^a	49.0 ^a	43.9 ^a	52.1 ^a	0.723
	18:1 trans-11	4.22 ^b	4.43 ^b	3.75 ^b	3.51 ^b	4.17 ^b	4.81 ^b	0.190
	18:2 trans-9, trans-11	0.07 ^a	0.07 ^a	0.05 ^a	0.05 ^a	0.04 ^a	0.03 ^a	0.630
	18:2 cis-9, cis-12	0.67 ^a	0.68 ^a	0.72 ^a	0.46 ^a	0.68 ^a	0.49 ^a	0.198
	18:3 n-3	1.01 ^b	0.97 ^b	0.96 ^b	0.67 ^a	0.84 ^b	0.73 ^a	0.015
24 hrs	18:0	58.2 ^a	55.0 ^a	56.0 ^a	55.8 ^a	50.7 ^a	61.0 ^a	0.656
	18:1 trans-11	4.20 ^a	5.16 ^a	4.87 ^a	5.15 ^a	4.90 ^a	5.94 ^a	0.723
	18:2 trans-9, trans-11	0.05 ^b	0.07 ^b	0.05 ^b	0.03 ^a	0.03 ^a	0.06 ^b	0.022
	18:2 cis-9, cis-12	0.61 ^a	0.52 ^a	1.04 ^b	0.53 ^a	0.55 ^a	0.37 ^a	0.005
	18:3 n-3	0.78 ^c	0.72 ^c	0.75 ^c	0.60 ^a	0.58 ^a	0.63 ^b	0.006

Table 2. Effects of linolenic acid addition on fatty acid concentrations *in vitro*

Time	Fatty Acid	linolenic acid Concentration (µM)					SED	P
		0	50	250	500	750		
0 hrs	18:0	54.9 ^a	59.0 ^a	60.8 ^a	51.6 ^a	52.5 ^a	60.9 ^a	0.335
	18:1 trans-11	4.16 ^a	3.86 ^a	4.13 ^a	6.65 ^b	4.08 ^a	4.56 ^a	0.001
	18:2 trans-9, trans-11	0.06 ^a	0.06 ^a	0.05 ^a	0.07 ^a	0.06 ^a	0.07 ^a	0.418
	18:2 cis-9, cis-12	0.61 ^{ab}	0.38 ^a	0.83 ^{bc}	1.05 ^c	0.72 ^{abc}	0.52 ^{ab}	0.047
	18:3 n-3	0.66 ^a	0.48 ^a	0.67 ^a	1.45 ^b	0.74 ^a	0.70 ^a	0.123
4 hrs	18:0	58.9 ^b	49.8 ^{ab}	44.1 ^a	41.6 ^a	48.7 ^{ab}	51.9 ^{ab}	0.162
	18:1 trans-11	5.48 ^{ab}	5.52 ^{ab}	7.61 ^b	6.33 ^{ab}	5.55 ^{ab}	3.64 ^a	0.151
	18:2 trans-9, trans-11	0.06 ^{bc}	0.08 ^c	0.07 ^{bc}	0.02 ^a	0.06 ^{bc}	0.03 ^{ab}	0.021
	18:2 cis-9, cis-12	0.53 ^a	1.580 ^b	0.76 ^a	0.80 ^a	1.28a	0.57 ^a	0.360
	18:3 n-3	0.70 ^a	1.51 ^b	0.91 ^{abc}	0.84 ^a	0.90 ^{ab}	0.64 ^a	0.075
8 hrs	18:0	57.0 ^b	53.8 ^b	42.9 ^a	50.1 ^{ab}	55.6 ^b	57.4 ^b	0.013
	18:1 trans-11	3.54 ^a	3.37 ^a	3.89 ^a	3.98 ^a	4.84 ^a	3.59 ^a	0.402
	18:2 trans-9, trans-11	0.05 ^b	0.05 ^b	0.00 ^a	0.07 ^c	0.08 ^c	0.08 ^c	<.001
	18:2 cis-9, cis-12	0.44 ^{ab}	0.42 ^a	0.64 ^{abc}	1.13 ^{bc}	0.44 ^{ab}	0.43 ^a	0.185
	18:3 n-3	0.76 ^a	0.71 ^a	0.80 ^a	1.46 ^b	0.67 ^a	0.67 ^a	0.116
24 hrs	18:0	54.9 ^a	59.2 ^a	57.8 ^a	55.2 ^a	49.4 ^a	49.4 ^a	0.263
	18:1 trans-11	3.43 ^a	3.65 ^a	4.05 ^a	4.47 ^a	4.56 ^a	4.12 ^a	0.531
	18:2 trans-9, trans-11	0.06 ^a	0.05 ^a	0.05 ^a	0.06 ^a	0.04 ^a	0.06 ^a	0.761
	18:2 cis-9, cis-12	0.41 ^a	0.46 ^a	0.49 ^a	1.26 ^b	0.50 ^a	0.47 ^a	<.001
	18:3 n-3	0.55 ^{ab}	0.52 ^{ab}	0.57 ^{ab}	0.64 ^b	0.50 ^{ab}	0.46 ^a	0.243

Discussion

Observed increases in PUFA, and decreases in 18:0 suggest that biohydrogenation was successfully inhibited, particularly when using 250-500µM concentrations. However, after 24 hours biohydrogenating bacteria appear to have recovered, and were biohydrogenating PUFA. RNA analysis showed no changes in the bacterial community over time or concentration, suggesting that rumen function would be unaffected. The next step is to evaluate whether such concentrations are achievable by enhancing lipolysis.

