

Aberystwyth University

Identifying drivers of changes in relative abundances in agroecosystems

Brophy, C.; Finn, J. A.; Luscher, A.; Suter, M.; Kirwan, L.; Sebastià, M. T.; Helgadóttir, Á.; Baadshaug, O. H.; Belanger, G.; Black, A.; Collins, R. P.; Jure, C.; Dalmannsdóttir, S.; Delgado, I.; Elgersma, A.; Fothergill, M.; Frankow-Lindberg, B. E.; Ghesquiere, A.; Golinska, B.; Grieu, P.

Published in:

Proceedings of the 27th General Meeting of the European Grassland Federation

Publication date:

2018

Citation for published version (APA):

Brophy, C., Finn, J. A., Luscher, A., Suter, M., Kirwan, L., Sebastià, M. T., Helgadóttir, Á., Baadshaug, O. H., Belanger, G., Black, A., Collins, R. P., Jure, C., Dalmannsdóttir, S., Delgado, I., Elgersma, A., Fothergill, M., Frankow-Lindberg, B. E., Ghesquiere, A., Golinska, B., ... Connolly, J. (2018). Identifying drivers of changes in relative abundances in agroecosystems. In B. Horan, D. Hennessy, M. O'Donovan, E. Kennedy, B. McCarthy, J. A. Finn, & B. O'Brien (Eds.), *Proceedings of the 27th General Meeting of the European Grassland Federation: Sustainable meat and milk production from grasslands* (Vol. 23, pp. 586-588). Teagasc.

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Identifying the drivers of changes in the relative abundances of species in agroecosystems

Brophy C.¹, Finn J.A.², Lüscher A.³, Suter M.³, Kirwan L.⁴, Sebastià M.T.^{5,6}, Helgadóttir Á.⁷, Baadshaug O.H.⁸, Bélanger G.⁹, Black A.^{10,26}, Collins R.P.¹¹, Čop J.¹², Dalmannsdóttir S.⁷, Delgado I.¹³, Elgersma A.^{14,27}, Fothergill M.¹¹, Frankow-Lindberg B.E.¹⁵, Ghesquiere A.¹⁶, Golińska B.¹⁷, Goliński P.¹⁷, Grieu P.¹⁸, Gustavsson A.M.¹⁹, Höglind M.²⁰, Huguenin-Elie O.³, Jørgensen M.²⁰, Kadziulienė Z.²¹, Kurki P.²², Llurba R.^{5,6}, Lunnan T.²⁰, Porqueddu C.²³, Thumm U.²⁴ and Connolly J.²⁵

¹*Department of Mathematics and Statistics, Maynooth University, Maynooth, Co Kildare, Ireland;*

²*Teagasc, Environment Research Centre, Johnstown Castle, Wexford, Ireland;* ³*Agroscope, Forage Production and Grassland Systems, Reckenholzstrasse 191, 8046 Zurich, Switzerland;* ⁴*UCD Institute of Food and Health, UCD, Belfield, Dublin 4, Ireland;* ⁵*Group GAMES & Dept HBJ, ETSEA, Universitat de Lleida, Av. Rovira Roure 191, 25198 Lleida, Spain;* ⁶*Laboratory ECOFUN, Centre Tecnologic Forestal de Catalunya, Ctra Sant Llorenç km 2, 25280 Solsona, Spain;* ⁷*Agricultural University of Iceland, Árleyni 22, 112 Reykjavík, Iceland;* ⁸*Department of Plant Sciences, Norwegian University of Life Sciences, P.O. Box 5003, 1432, Ås, Norway;* ⁹*Agriculture and Agri-Food Canada, 2560, Hochelaga Blvd, Québec G1V 2J3, Canada;* ¹⁰*Teagasc, Beef Research Centre, Grange, Dunsany, Co. Meath, Ireland;* ¹¹*IBERS, Aberystwyth University, Plas Gogerddan, Aberystwyth, SY23 3EB, Wales, United Kingdom;* ¹²*Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, 1000 Ljubljana, Slovenia;* ¹³*CITA-DGA, Av. Montañana 930, 50059 Zaragoza, Spain;* ¹⁴*Plant Sciences Group, Wageningen University, P.O. Box 16, 6700, AA Wageningen, the Netherlands;* ¹⁵*Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Box 7043, 750 07 Uppsala, Sweden;* ¹⁶*ILVO, Plant Science, Applied Genetics and Breeding, Caritasstraat 39, 9090, Melle, Belgium;* ¹⁷*Department of Grassland and Natural Landscape Sciences, Poznan University of Life Sciences, Dojazd 11, 60-632, Poznan, Poland;* ¹⁸*UMR AGIR, INP-ENSAT, University of Toulouse, 31326, Castanet Tolosan, France;* ¹⁹*Department of Agricultural Research for Northern Sweden, Swedish University of Agricultural Sciences, 901 83, Umeå, Sweden;* ²⁰*NIBIO, Norwegian Institute of Bioeconomy Research, P.O. Box 115, 1431 Ås, Norway;* ²¹*Institute of Agriculture, Lithuanian Research Centre for Agriculture and Forestry, Akademija, 58344, Kedainiai, Lithuania;* ²²*Natural Resources Institute (Luke), Management and Production of Renewable Resources, Lönnrotinkatu 5, 50100, Mikkeli, Finland;* ²³*CNR-ISPAAM, Traversa la Crucca 3, località Balduca, 07100, Sassari, Italy;* ²⁴*Department of Crop Science, University of Hohenheim, 70593, Stuttgart, Germany;* ²⁵*School of Mathematics and Statistics, University College Dublin, Dublin 4, Ireland;* *Present addresses:* ²⁶*Faculty of Agriculture and Life Sciences, P.O. Box 85084, Lincoln University, Lincoln 7647, Canterbury, New Zealand* ²⁷*Independent scientist, P.O. Box 323, 6700 AH, Wageningen, the Netherlands*

Abstract

Increasing species diversity often promotes ecosystem functions in grasslands, but sward diversity may be reduced over time through competitive interactions among species. We investigated the development of species' relative abundances in intensively managed agricultural grassland mixtures over three years to identify the drivers of diversity change. A continental-scale field experiment was conducted at 31 sites using 11 different four-species mixtures each sown at two seed abundances. The four species consisted of two grasses and two legumes, of which one was fast establishing and the other temporally persistent. We modelled the dynamics of the four-species mixtures over the three-year period. The relative abundances shifted substantially over time; in particular, the relative abundance of legumes declined over time but stayed above 15% in year three at many sites. We found that species' dynamics were primarily driven by differences in the relative growth rates of competing species and secondarily by density dependence and climate. Alongside this, positive diversity effects in yield were found in all years at many sites.

Keywords: biodiversity, dynamics, grass, legume, multispecies mixtures, relative growth rate

Introduction

The common practice of managing highly fertilised grassland monocultures has often been critiqued and there is a need for productive grass-legume systems that require less fertiliser and lead to improved environmental outcomes (Lüscher *et al.*, 2014, Suter *et al.*, 2015). There is wide consensus that increasing species diversity promotes many ecosystem functions. Over time, however, some species in a mixture may become dominant at the expense of others and sward diversity may decline, thus reducing the benefits to ecosystem function (Carroll *et al.*, 2011). Here, we examine the dynamics of the relative abundances of multiple species in agronomic grassland mixtures and identify reasons why changes occur at the species level across 31 coordinated multi-year experimental sites.

Materials and methods

A common experiment was carried out at 30 sites across Europe and one site in Canada. At each site 22 four-species mixture plots were established. The four species comprised two grasses and two legumes, of which one was fast establishing and the other temporally persistent. Thus, there were four functional groups: grass (G) / legume (L) by fast establishing (F) / temporally persistent (P) which were denoted G_F , G_P , L_F and L_P . The identity of the species within functional groups varied across the sites; yet, there was a total of 11 unique species used across the experiment. At each site, the relative abundances of the four species were varied systematically across 11 mixture plots ranging from each species equally present (25% of each) to one species dominant (70%, 10%, 10%, 10%) and each of the 11 mixtures was sown at two seed abundance levels. Monocultures of each species were also established at each seed abundance level, giving an additional eight plots at each site. N fertiliser was applied at most sites (maximum rate of 150 kg N ha⁻¹ annum) and plots were harvested between two and seven times per annum depending on local practice. The annual plot-level biomass of each of the four species was recorded for three years following the year of establishment. Further experimental details are available in Kirwan *et al.* (2007, 2014).

We analysed relative growth rates (Connolly and Wayne, 2005) for each species in mixture to explain changes in relative abundances for sown-year 1, years 1 - 2 and years 2 - 3.

Results and discussion

Across all sites, we found significant changes in the relative abundances of our four-species mixtures over the three years. The main driver of those changes was differences in the relative growth rates of species. On average across all sites, the temporally persistent grass (G_P) become dominant by year 3 (Figure 1) but the relative abundance of G_P in year 3 varied substantially across sites, ranging from 5% at one site to 100% at another (Brophy *et al.*, 2017). The relative abundance of legumes ($L_F + L_P$) was generally high in year 1, and while it declined over time, there were 12 sites that still had average legume abundance above 15% in year three. Legume persistence was positively related to sites' annual minimum temperature (computed as the average of the lowest five annual values) in years 2 ($P = 0.002$) and 3 ($P = 0.003$). Overall, we found several intra- and inter-specific density-dependent dynamics in our multi-species communities, which gave evidence for stabilising processes acting on the system (Brophy *et al.*, 2017). Alongside the substantial shifts in dynamics, Brophy *et al.* (2017) and Finn *et al.* (2013) showed significant positive diversity effects at many sites in all three years, the strengths of which were positively related to legume abundance.




Figure 1. The average relative abundance of each functional group across all sites in each (post-seeding) year. The species are classified according to functional traits: fast establishing grass (G_f) and legume (L_f) and temporally persistent grass (G_p) and legume (L_p).

Conclusion

This continental-scale field experiment showed the importance of the relative growth rates of competing species for community dynamics and species shift over time. Alongside this, significant positive diversity effects were evident across the three experimental years at many sites. Diversity effects in multi-species mixtures can be further enhanced through the inclusion of legumes and strategic selection of the species and their cultivars, paying particular attention to their traits and competitive abilities relative to each other.

Acknowledgements

Support was received from Science Foundation Ireland, grant number 09/RFP/EOB2546, the European's Seventh Framework Programme (FP7/2007-2013; grant agreement no. 266018) and the EU Commission through COST Action 852.

References

- Brophy C. *et al.* (2017) Major shifts in species' relative abundance in grassland mixtures alongside positive effects of species diversity in yield: a continental-scale experiment. *Journal of Ecology* 105, 1210-1222.
- Carroll I.T., Cardinale B.J. and Nisbet R.M. (2011) Niche and fitness differences relate the maintenance of diversity to ecosystem function. *Ecology* 92, 1157-1165.
- Connolly J. and Wayne P. (2005) Assessing determinants of community biomass composition in two-species plant competition studies. *Oecologia* 142, 450-457.
- Finn J.A. *et al.* (2013) Ecosystem function enhanced by combining four functional types of plant species in intensively managed grassland mixtures: a 3-year continental-scale field experiment. *Journal of Applied Ecology* 50, 365-375.
- Kirwan L. *et al.* (2014) The Agrodiversity Experiment: three years of data from a multisite study in intensively managed grasslands. *Ecology* 95, 2680.
- Kirwan L. *et al.* (2007) Evenness drives consistent diversity effects in intensive grassland systems across 28 European sites. *Journal of Ecology* 95, 530-539.
- Lüscher A., Mueller-Harvey I., Soussana J.F., Rees R.M. and Peyraud J.L. (2014) Potential of legume-based grassland-livestock systems in Europe: a review. *Grass and Forage Science* 69, 206-228.
- Suter M. *et al.* (2015) Nitrogen yield advantage from grass-legume mixtures is robust over a wide range of legume proportions and environmental conditions. *Global Change Biology* 21, 2424-2438.