

Aberystwyth University

The role of nitrite and nitric oxide under low oxygen conditions in plants

Gupta, Kapuganti Jagadis; Mur, Luis A. J.; Wany, Aakanksha; Kumari, Aprajita; Fernie, Alisdair R.; Ratcliffe, R. George

Published in:
New Phytologist

DOI:
[10.1111/nph.15969](https://doi.org/10.1111/nph.15969)

Publication date:
2020

Citation for published version (APA):

Gupta, K. J., Mur, L. A. J., Wany, A., Kumari, A., Fernie, A. R., & Ratcliffe, R. G. (2020). The role of nitrite and nitric oxide under low oxygen conditions in plants. *New Phytologist*, 225(3), 1143-1151.
<https://doi.org/10.1111/nph.15969>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

DR JAGADIS GUPTA KAPUGANTI (Orcid ID : 0000-0002-7090-5097)

DR LUIS ALEJANDRO MUR (Orcid ID : 0000-0002-0961-9817)

DR ALISDAIR FERNIE (Orcid ID : 0000-0001-9000-335X)

Article type : Research Review

The role of nitrite and nitric oxide under low oxygen conditions in plants

Kapuganti Jagadis Gupta^{1*}, Luis A.J. Mur², Aakanksha Wany¹, Aprajita Kumari¹, Alisdair R. Fernie³, R. George Ratcliffe^{4*}

¹National Institute of Plant Genome Research, Aruna Asaf Ali Marg, 110067, New Delhi, Delhi, India.

²Institute of Environmental and Rural Science, Aberystwyth University, Edward Llwyd Building, Aberystwyth SY23 3DA, UK

³Max-Planck-Institute of Molecular Plant Physiology, Am Mühlenberg 1, D-14476 Potsdam-Golm, Germany

⁴Department of Plant Sciences, University of Oxford, South Parks Road, Oxford OX1 3RB, UK

*To whom correspondence should be addressed

Email: jgk@nipgr.ac.in

Tel: +91-11-26735111

ORCID Id: <http://orcid.org/0000-0002-7090-5097>

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/nph.15969

This article is protected by copyright. All rights reserved.

Email: george.ratcliffe@plants.ox.ac.uk

ORCID Id: 0000-0001-8394-157

Tel: +44 (0) 1865 275001

Received: 25 April 2019

Accepted: 24 May 2019

Summary

Plant tissues, particularly roots, can be subjected to periods of hypoxia due to environmental circumstances. Plants have developed various adaptations in response to hypoxic stress and these have been extensively described. Less well-appreciated is the body of evidence demonstrating that scavenging of nitric oxide (NO) and the reduction of nitrate/nitrite regulate important mechanisms that contribute to tolerance to hypoxia. Whilst ethylene controls hyponasty and aerenchyma formation, NO production apparently regulates hypoxic ethylene biosynthesis. In the hypoxic mitochondrion, cytochrome c oxidase, which is a major source of NO, is also inhibited by NO, thereby reducing the respiratory rate and enhancing local oxygen concentrations. Nitrite can maintain ATP generation under hypoxia by coupling its reduction to the translocation of protons from the inner side of mitochondria and generating an electrochemical gradient. This reaction can be further coupled to a reaction whereby non-symbiotic haemoglobin oxidizes NO to nitrate. In addition to these functions, nitrite has been reported to influence mitochondrial structure and supercomplex formation, as well as playing a role in oxygen sensing via the N-end rule pathway. These studies establish that nitrite and NO perform multiple functions during plant hypoxia and suggest that further research into the underlying mechanisms is warranted.

Keywords: cytochrome c oxidase, hyponasty, hypoxia, mitochondria, nitric oxide (NO), nitrite.

Introduction

Molecular oxygen facilitates the efficient production of ATP in all aerobic eukaryotic organisms by providing the terminal electron acceptor for the mitochondrial electron transport chain. Oxygen deprivation, leading to a state of hypoxia (low oxygen) or anoxia (no oxygen), compromises the process of oxidative phosphorylation. For plants, this problem typically arises during flooding (Bailey-Serres *et al.*, 2012) as a result of the 10^4 -fold reduction in gaseous diffusion in flood water (Armstrong, 1979). Moreover, even under optimal growth conditions, certain dense tissues such as seeds (Borisjuk *et al.*, 2007) and tubers (Geigenberger *et al.*, 2000) are hypoxic, with O_2 concentrations in the range 1 - 50 μM . While the latter observations indicate that plants can cope with low levels of O_2 during normal development, hypoxia inevitably restricts the availability of oxygen for oxidative phosphorylation and increases the importance of fermentation as a source of ATP (Ricard *et al.*, 1994). As a result, plant metabolism must adapt to lower ATP production, including the induction of energy-conserving pathways of sucrose degradation that lead to improved plant performance under hypoxia (Geigenberger *et al.*, 2000; Bologna *et al.*, 2003).

As well as metabolic adaptations to hypoxia, plants have also developed anatomical and morphological adaptations, including the formation of aerenchyma, aerial adventitious roots, and leaf gas films (Bailey-Serres *et al.*, 2012). Responses to hypoxia have been extensively studied in plants and many transcriptional, post-translational and metabolic events that regulate these responses have been identified (Geigenberger *et al.*, 2000; Licausi *et al.*, 2011; Narsai *et al.*, 2017; Fukao *et al.*, 2019). One emerging theme is the involvement of reactive oxygen species (ROS) and nitric oxide (NO) signalling under low oxygen in plants (Pucciariello and Perata, 2017), and in this update the intention is to focus on the multiple roles of NO and nitrite in the hypoxic response.

Hypoxic synthesis and turnover of NO

NO is a free radical signalling molecule that is produced by several oxidative and reductive pathways (Gupta *et al.*, 2011; Astier *et al.*, 2017). The reductive pathways are active under hypoxic conditions, with mitochondria playing a major role in NO production through the action of cytochrome oxidase (COX) and other deoxyhemeproteins (Figure 1). Isolated plant mitochondria typically produce NO at a rate of 1-20 nmol.mg protein⁻¹.h⁻¹ within a few minutes of adding NADH and nitrite to a hypoxic incubation medium, and the K_i for oxygen, which inhibits the process, is 0.05% (Gupta *et al.*, 2005). High levels of NO lead to cell death (Wang *et al.* 2013), so if NO is to have other signalling functions under hypoxia it is necessary to have mechanisms for preventing its excessive accumulation.

NO production is countered by NO degradation, with several haem proteins such as flavohaemoglobin, haemoglobin (Hb), myoglobin and their associated reductases fulfilling this role in animal cells (Gardner, 2005) and flavoglobin scavenging NO in yeast (Liu *et al.*, 2000; Cassanova *et al.*, 2005). In plants, Class 1 phytooglobins (Pgb) are efficient NO scavengers (Hebelstrup *et al.*, 2008) and their very high affinity for oxygen ($K_m \sim 2$ nM) allows them to function under hypoxia (Figure 1). Overexpression of the Pgb gene in barley decreased NO release under hypoxia, while knockdown of the gene increased it, confirming that Pgb makes a significant contribution to the regulation of NO levels (Cochrane *et al.*, 2017). The inhibition of COX by NO (Brown and Cooper, 1994; Cleeter *et al.*, 1994) also facilitates the operation of the oxygen-requiring Pgb-NO cycle under hypoxia by inhibiting respiration at low oxygen concentrations. Thus under hypoxia oxygenated Pgb converts NO to nitrate, and the resulting metphytooglobin is converted back to Pgb by monodehydroascorbate reductase-mediated ascorbate reduction (Igamberdiev *et al.*, 2006; Gupta and Igamberdiev, 2011). The Pgb-NO cycle also regenerates NAD⁺ and may be considered an alternative to the usual pathways of fermentation (Igamberdiev and Hill, 2004), although like lactate fermentation, the Pgb-NO cycle is acidifying (Libourel *et al.*, 2006) and thus a potential contributor to acidosis under hypoxia.

S-nitrosoglutathione reductase (GSNOR) is another enzyme that contributes to the regulation of NO levels in plants (Leterrier *et al.*, 2011). GSNOR converts the NO derivative S-nitrosoglutathione (GSNO) to oxidised glutathione (GSSG) and ammonia, and it was recently shown that the inhibitory NO-dependent S-nitrosation of GSNOR1 (Frunghillo *et al.*, 2014) leads to degradation of the enzyme by selective autophagy (Zhan *et al.*, 2018). Elimination of the S-nitrosation site in GSNOR abolished the positive effect of NO on the hypoxic germination of Arabidopsis seeds, indicating that the NO-dependent post-translational modification of GSNOR is a physiologically relevant process that contributes to the hypoxic response (Zhan *et al.*, 2018). Elevated GSNO was shown to increase the expression of both alcohol dehydrogenase and pyruvate decarboxylase in germinating Arabidopsis seeds (Zhan *et al.*, 2018) emphasising the importance of GSNOR regulation by NO under hypoxia.

NDB-type dehydrogenases also play a role in NO degradation by forming superoxide anions that convert NO to peroxynitrite (ONOO^-) (de Oliveira *et al.*, 2008). This route of NO degradation is stimulated by calcium, and abolished by superoxide dismutase and complete anoxia. These observations indicate that NDB dehydrogenases actively generate superoxide, and are involved in superoxide-dependent NO degradation. NDB-type dehydrogenases were also found to be induced in transgenic Arabidopsis plants with downregulated expression of GSNOR (Frunghillo *et al.*, 2013), providing further correlative evidence for their role in NO homeostasis.

The net result of these biosynthetic and degradative pathways is a marked increase in NO production in response to hypoxia. This increase is in turn increasingly implicated in a range of adaptive responses to oxygen deprivation, including hyponasty, aerenchyma formation, oxygen homeostasis, mitochondrial activity and oxygen sensing

Role of NO in hyponasty under hypoxia

The hyponastic response is a highly effective escape strategy employed by most plant species following submergence, shade or elevated ambient temperatures. It is accompanied by a strong directional growth mediated by reversible turgor reactions and changes in the osmotic state of the cells. It mainly depends on the unequal growth rates of two anatomically different sides of the organ in question. Hyponasty can be defined as a type of asymmetric growth, whereby abaxial tissue displays a higher growth rate than the adaxial cells. It is a common feature in leaf blades and petioles of many monocots and dicots (Polko *et al.*, 2011).

During flooding, ethylene plays a major role in the hyponastic response due to its reduced diffusion in submerged tissues (Voesenek *et al.*, 1993). Vreeburg *et al.* (2005) showed that ethylene-mediated hyponastic signaling is characterized by acidification of the apoplast and a higher expression of expansin proteins, both of which play important roles in modifying cell wall structure. Ultimately, ethylene responsive factors (ERFs) mediate hyponasty, however, the function of these proteins is gibberellin (GA) dependent (Polko *et al.*, 2011). Other hormones are also implicated in the process, with auxins such as indole-3-acetic acid (IAA) positively regulating stage-specific submergence-induced hyponasty, whilst abscisic acid (ABA) acts as a negative regulator of this process (Cox *et al.*, 2006).

NO is also involved in the complex signalling network leading to hyponasty. Arabidopsis seedlings produce increased ethylene and NO under hypoxia, and Pgb gene expression (*AtGLB1*) correlated with hyponastic growth (Hebelstrup *et al.*, 2012). These observations led to the proposal that NO likely acts as a regulator of hyponasty via induction of ethylene synthesis. Plants also experience shading and reduced light levels during flooding, and since shading alone can induce an ethylene-dependent hyponastic response (Pierik *et al.*, 2009) it is possible that shade also has an impact on NO metabolism. It is known that light-dark dynamics can influence both NO and nitrite levels (Planchet *et al.*, 2005), but the relevance of this observation under more natural conditions remains to be evaluated.

Role of NO in the formation of ethylene-induced aerenchyma

Another adaptive response of plants to hypoxia is the formation of aerenchyma, the gas-filled tissue that allows the exchange of gases between shoot and root under conditions of flooding and waterlogging (Drew *et al.*, 2000). Schizogenous aerenchyma is formed by a process of cell separation at the middle lamella during cell development, whilst lysigenous aerenchyma is formed as a consequence of the random death of cortical cells. It was previously shown that hypoxia is an inducer of aerenchyma formation (Drew *et al.*, 2000), and that ethylene plays a role in cortical cell death (Yamauchi *et al.* 2014). NO also plays a role in programmed cell death (Delledonne *et al.*, 1998; Chen *et al.*, 2009; Wang *et al.*, 2013), and so the possibility arises that NO could play a role in aerenchyma formation in hypoxic tissues given that NO production takes place within a few minutes of the onset of hypoxia (Gupta *et al.*, 2005).

Recently Wany *et al.* (2017) investigated whether ethylene-induced aerenchyma formation in wheat roots required hypoxia-induced NO. Wheat roots produced NO under hypoxia as expected, and scavenging of NO by 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (cPTIO) led to a marked reduction in aerenchyma formation following 24 or 48 hours of hypoxia. Interestingly, it was found that hypoxically-induced NO was important for the induction of the genes encoding 1-aminocyclopropane-1-carboxylic acid (ACC) synthase and ACC oxidase, both of which are required for ethylene biosynthesis, as well as the ethylene-responsive genes *ERF1* and *PDF13*. Cell death events such as increased electrolyte leakage, increased cellulase activity, DNA fragmentation, and cytoplasmic streaming were all inhibited under hypoxia in the presence of the NO scavenger, reinforcing the conclusion that NO is essential for the development of ethylene-induced aerenchyma (Wany *et al.*, 2017). Moreover, ethylene inhibitors and NO scavengers, either alone or in combination, suppressed the genes involved in signal transduction leading to aerenchyma development. These findings suggest that NO plays a role in aerenchyma formation in wheat, acting either upstream of ethylene, or in parallel with it. The involvement of NO in aerenchyma formation was also indicated by the observation that the *respiratory burst oxidative homolog/NADPH oxidase*

(*RBOH/NOX*) gene, which is known to have a role in aerenchyma formation (Yun *et al.*, 2011), is also induced by NO (Wany *et al.*, 2017). *RBOH/NOX* plays a role in superoxide production, hence the induction of *RBOH/NOX* by NO, correlated with the generation of ROS in wheat cortical cells. Since superoxide reacts with NO to form peroxynitrite, a strong nitrating agent, increased tyrosine nitration was observed during aerenchyma formation. It will be interesting to see whether these results can be replicated under conditions more akin to natural flooding events, since ethylene is known to accumulate to saturating levels rapidly and the possibility of induction by NO has not been considered hitherto (Sasidharan *et al.*, 2018).

Plants are well equipped with detoxification systems to counter the deleterious effects of ROS and NO. In particular, there are many regulatory mechanisms that influence the levels of ROS and NO, thus protecting plants from severe damage under stress conditions. However, during processes such as programmed cell death and aerenchyma formation, plant cells need to maintain high ROS levels in the tissues where limited cell death is advantageous. One strategy to achieve this is to lower antioxidant gene expression (Liu *et al.*, 2018). Wany and Gupta (2018) found an inverse correlation between antioxidant gene expression and increased ROS following 24 h of hypoxia in wheat roots during aerenchyma formation. NO is known to increase antioxidant gene expression during stress (Tossi *et al.*, 2011), but during cell death it seems that suppression of the antioxidant mechanism is required. A prominent example is the inhibition of glycine decarboxylase by S-nitrosation, which alters cellular redox status and promotes cell death (Palmieri *et al.*, 2010). A survey of S-nitrosylated proteins in cells undergoing aerenchyma formation could potentially afford new insights into the role of NO in the promotion or limitation of cell progression. Such insights could be based on transgenic manipulation of root NO levels, for example by over-expression of *Pgb*, during aerenchyma formation, although this approach might also affect oxygen homeostasis and root development (Gupta *et al.*, 2014). An alternative approach might be to test the relationship between the nitrogen status of the soil and aerenchyma formation, given that the synthesis of NO ultimately depends on the availability of nitrate (Planchet *et al.*, 2005; Gupta *et al.*, 2013). Preliminary evidence suggests that nitrate nutrition, as

opposed to ammonium nutrition, favours aerenchyma formation and this needs further investigation (Wany and Gupta, 2018).

While the recent evidence suggests that NO is essential for the development of aerenchyma via cortical cell death under flooding stress (Wany *et al.*, 2017), it is necessary to regulate the levels of NO in other root zones to avoid cell death in tissues required for continued growth (Mira *et al.*, 2016a). The root apical meristem (RAM) contains stem cells which are important for root growth and it has been shown that transgenic suppression of the hypoxically-induced phytoglobins ZmPgb1.1 or ZmPgb1.2. led to structural abnormalities in RAM (Mira *et al.*, 2016b). Suppression of Pgb also enhanced expression of ethylene biosynthetic and responsive genes, providing further support for the role of Pgb in regulating NO levels under hypoxia. In contrast, overexpression of Pgb improved hypoxic root growth by alleviating apical meristem cell death again emphasising the role of Pgb as NO scavengers. These observations highlight the need for differential regulation of Pgb expression in different root zones to ensure that hypoxically-induced NO can promote cortical cell death and aerenchyma formation without damage to the stem cells in the RAM. The mechanism by which this is achieved is not fully understood, but it is relevant that exposure of the root apex to hypoxia has been shown to lead to increased hypoxic acclimation of the entire root (Mugnai *et al.*, 2012), emphasising the existence of systemic signalling pathways that coordinate cell-type specific responses to hypoxia.

Role of NO in oxygen homeostasis

Oxygen homeostasis is important for maintaining an appropriate internal oxygen level in tissues during normal development. This phenomenon is crucial when environmental effects, such as flooding and waterlogging, reduce the oxygen supply and drive the tissues towards anoxia. Recently it was shown that NO has a potential role in oxygen homeostasis under normoxia via the regulation of respiration (Gupta *et al.*, 2014). It is well known that NO inhibits respiration by inhibiting cytochrome c oxidase in isolated mitochondria (Millar and Day, 1998). NO binds to the Fe²⁺-haem

group at the O₂-binding site of the binuclear centre Fe_{a3}Cu_B in COX (Cleeter *et al.*, 1994) and this provides the basis for an autoregulatory mechanism in which increasing NO under hypoxia reduces oxygen consumption. The relevance of this for oxygen homeostasis has been demonstrated in normoxic barley roots, where overexpression of Pgb promoted NO scavenging, increased the respiration rate, and decreased the internal oxygen level (Gupta *et al.*, 2014). Overexpression of Pgb also affected the normoxic NO signalling pathways in barley (Cochrane *et al.*, 2017). The physiological significance of this effect has been shown in both seeds (Borisjuk *et al.*, 2007) and isolated mitochondria (Benemar *et al.*, 2008), where it was shown that nitrite reduction at complex III reversibly inhibited COX, and thus contributed to the maintenance of a steady state level of oxygen in the mitochondria.

The role of NO in oxygen homeostasis is also important in seed germination. This process is associated with the production of NO and a decrease in ABA via regulation of *CYP707A2* transcription and (+)-abscisic acid 8'-hydroxylase (Liu *et al.*, 2009). Gibbs *et al.* (2014b) reported that both NO and oxygen availability promote degradation of ERF VII transcription factors during the metabolically active state of seed development, leading to down-regulation of AB15 in the endosperm and the promotion of germination.

Role of NO in mitochondrial activity under hypoxia

Oxygen deprivation can have a marked effect on plant mitochondrial structure, and the observed changes correlate to some extent with the ability of the plant to survive periods of hypoxia or anoxia (Vartapetian *et al.*, 2003; Shingaki-Wells *et al.*, 2014). Nitrate has been shown to have a protective effect on mitochondrial ultrastructure under these conditions (Vartapetian *et al.*, 2003), but recent evidence suggests that nitrite confers similar protection, and that the reduction of nitrite to NO under hypoxia is important for the maintenance of some level of mitochondrial activity (Gupta *et al.*, 2017). Thus incubating hypoxic pea root mitochondria with 0.5 mM nitrite resulted in increased NO production, improved mitochondrial integrity, improved energization of the inner mitochondrial membrane, increased ATP synthesis, lower levels of reactive

oxygen species, and decreased lipid peroxidation. Nitrite also increased the activities of complex I and the supercomplex I + III₂ under hypoxia. These observations highlight the far-reaching effects of nitrite on the hypoxic mitochondrion (Gupta *et al.*, 2017).

The effect of nitrite on the activities of complex I and the supercomplex I + III₂ under hypoxia (Gupta *et al.*, 2017) may well be important in promoting the reduction of nitrite to NO through the maintenance of a fully functional electron transport chain. In tobacco plants deficient in complex I, reduced electron flow in the mitochondrial ETC led to lower NO production (Shah *et al.*, 2013); while supercomplex formation is considered to increase the efficiency of electron transport (Cogliati *et al.*, 2016). The fact that nitrite treatment under hypoxia increases the activities of complex I and the supercomplex I + III₂ hints at a regulatory role for either nitrite or NO under these conditions.

Interpretation of all the effects of nitrite on mitochondrial function under hypoxia is complicated by the potential regulatory effects of the hypoxically-generated NO. For example, COX, the major site for the production of hypoxically-generated NO, is inhibited by NO (Cleeter *et al.*, 1994) and the interaction with NO can increase the efficiency of oxidative phosphorylation (Clerc *et al.*, 2007). These factors are seen in the nitrite-stimulated increase in ATP synthesis in hypoxic pea root mitochondria (Gupta *et al.*, 2017), but at the same time it should be noted that NO scavenging by mitochondria, or via cytosolic scavenging systems, is considerable, with tobacco root mitochondria, for example, consuming 87% of the NO applied within two minutes (Kumari *et al.*, 2016).

Hypoxically-produced NO may also alter mitochondrial activity through changes in AOX activity. There are established links between AOX and NO, with AOX preventing excess NO production in tobacco leaves (Cvetkovska and Vanlerberghe, 2012), and NO inducing AOX under hypoxia (Gupta *et al.*, 2012) and phosphate

deficiency (Royo *et al.*, 2015). Recently, Vishwakarma *et al.* (2018) demonstrated that AOX prevents excess production of NO, peroxyntirite and tyrosine nitration under normoxia. However, it was also found that AOX can generate NO under hypoxia, and that the NO was oxidized via the Pgb-NO cycle (Vishwakarma *et al.*, 2018). Inhibiting AOX under hypoxia led to lower ATP, but AOX overexpressing lines produced more ATP. These data suggested that AOX-mediated NO production plays a role in the production of ATP under hypoxia by supporting proton translocation through complex I. Interestingly, in contrast to normoxia, it was shown that excess NO generated under hypoxia did not lead to the formation of peroxyntirite and tyrosine nitration. Thus, the link between AOX and NO differs between normoxia and hypoxia.

The phenomena of nitrite-driven ATP synthesis and mitochondrial protection are important in specialized structures such as nodules (Berger *et al.*, 2018). *Medicago truncatula* nodules have been shown to increase their production of NO when submitted to hypoxic conditions (Horchani *et al.*, 2011). The nodule oxygen concentration in the cytosol of the host plant cells is typically in the range 5-60 nM due to diffusion resistance and the respiration of the bacteroids. Under these conditions AOX does not contribute to respiration due to its higher K_m , but COX with a K_m value of 50 nM (Millar *et al.*, 1995) is expected to be functional. However, whether the amount of oxygen supplied by leghemoglobin (K_m for oxygen binding 2 nM) to the mitochondria is sufficient for energy production remains an open question (Horchani *et al.*, 2011).

As both Lb and Pgb have the capacity to oxidize NO or nitrate, this may allow the Pgb-NO cycle to operate generating a limited amount of ATP to sustain nodule development and function. The study by Horchani *et al.* (2011) provided evidence that in N_2 -fixing nodules of *M. truncatula*, the energy status of the nodules depends largely on NR functioning under normoxic, or hypoxic conditions. Thus, the Pgb-NO cycle can increase energy efficiency in specialised hypoxic organs such as nodules.

Role of NO in oxygen sensing under hypoxia

The precision and specificity of the control of molecular and physiological responses to low oxygen stress (Geigenberger *et al.*, 2000) suggests that plants possess sensitive oxygen-sensing mechanisms to initiate hypoxic responses. Direct and indirect sensors help in the development of these responses. Direct sensors are specific proteins such as transcriptional activators or repressors that sense oxygen. Prominent examples are the transcription factor hypoxia-inducible factor-1-alpha (HIF-1 α) in animals (Brahimi-Horn *et al.*, 2005), and a heme-binding protein kinase, FixL in rhizobial bacteria (Akimoto *et al.*, 2003). In contrast, indirect sensing relies on hypoxically-induced changes in such properties as calcium levels, energy status and redox status to trigger regulatory mechanisms (Bailey-Serres and Chang, 2005).

Recently, the direct oxygen sensing mechanism known as the N-end pathway has been shown to initiate the response of plants to hypoxia (Licausi *et al.*, 2011; Gibbs *et al.*, 2011). This is an evolutionarily conserved pathway for protein degradation whereby the stability of a protein is determined by the identity of its N-terminal residues (Varshavsky, 2011; Gibbs *et al.* 2014a; 2015). Specifically, the presence of N-degrons and N-terminal destabilizing residues determines whether a protein will be degraded by the proteasome (Graciet *et al.*, 2009; Holman *et al.*, 2009). Group VII ethylene response factors (ERFs), which have been shown to be important regulators of the response to low oxygen (Hinz *et al.*, 2010; Licausi *et al.*, 2010, 2011; Gibbs *et al.*, 2011, 2015), are oxygen-dependent substrates of the N-end rule pathway: these proteins are destabilised in the presence of oxygen and NO and stabilised in their absence. More recently, the polycomb repressive complex 2 component VRN2 has also been identified as an O₂/NO regulated target of the N-end rule pathway, suggesting a potential link between low oxygen/NO and the epigenetic control of gene expression (Gibbs *et al.*, 2018).

The N-terminal (Nt) MCGGAIL/L domain of the ERFVII transcription factors is the target for the N-end rule degradation pathway. Under aerobic conditions, methionine amino peptidase cleaves the Nt-Met to reveal an Nt-Cys, which is then oxidized by

Accepted Article

plant cysteine oxidases (PCOs) to produce Nt-Cys sulfinic acid. The oxidized Nt-Cys is arginylated by an arginyl transferase, creating a substrate for an E3 ligase which leads to polyubiquitination and proteasomal degradation of the ERFVII protein (Figure 2). The role of the PCOs in controlling hypoxic gene expression has been confirmed by genetic studies (Weits *et al.*, 2014) and the molecular mechanism of the oxidation and arginylation steps has been characterized *in vitro* (White *et al.*, 2017).

The extent to which NO influences the oxygen sensing system in plants is unclear. In animal systems, it has been shown that the *in vivo* oxidation of Nt-Cys before arginylation requires NO (Hu *et al.*, 2005) and the hydrolysis of S-nitrosothiols can produce sulfenic acids as the first step in the formation of sulfinic acids (Reddie and Carroll, 2008). In agreement with this, there is some evidence that NO, as well as oxygen, may be required for the degradation of ERFVIIs. Gibbs *et al.* (2014b) showed that ERFVIIs are destabilized in the presence of NO, and stabilized in their absence. In particular, the stability of two ERFVIIs, RAP2.3 and HRE2, was increased in Arabidopsis seedlings in the presence of NO scavengers, and also in the nitrate reductase-deficient *nia1nia2* mutant, which has greatly reduced levels of NO. More recently, Vicente *et al.* (2017) demonstrated that down-regulation of nitrate reductase in Arabidopsis led to lower NO levels and increased stability of ERFVIIs, an effect which was implicated in abiotic stress sensing under normoxia. While it remains the case that low oxygen is the primary determinant of ERFVII stability, the sensitivity of ERFVIIs to NO raises the possibility that the change in NO levels under hypoxia could modulate the oxygen sensing role of the ERFVIIs.

Accepted Article

Another consideration, which needs further investigation, is the potential impact of ROS generation under low oxygen (Vergara *et al.*, 2012). In principle, this could oxidise the Cys residues of ERFVIIs and thus work against their stabilization under hypoxia. At the same time, hypoxia-induced NO generation by mitochondria could play a role in removing excess ROS to ensure the stability of ERFVIIs under hypoxia, but the extent to which this is important has yet to be established and could well

differ between nitrate- and ammonium-grown plants (Wany *et al.*, 2019). More generally the sensitivity of mitochondrial activity to oxygen availability might suggest that mitochondrially-derived NO and ROS could contribute to retrograde signalling in the hypoxic state, but this also remains to be established.

Concluding remarks

It is now clear that nitrite and NO play important and multi-faceted roles in the response of plants to hypoxia. These include classical morphological changes such as hyponasty, the protection of mitochondrial structure, ATP generation and ROS scavenging. However, the role of hypoxically produced NO in various other plant anatomical adaptive responses to flooding, such as aerial lateral root formation, stem elongation, suberin and lignin accumulation needs more investigation. Cross talk between NO and growth hormones such as ethylene, auxin, and ABA during these adaptive responses to hypoxia would also merit further investigation. Another area of interest is the potential effect of the nitrogen supply on hypoxic tolerance given that both nitrite and NO are derived from nitrate. For example, lines with differing nitrogen use efficiency could improve the availability of nitrate and hence affect tolerance to hypoxia. Other targets for further analysis of the role of nitrite and NO under hypoxia include germinating seeds, which experience varying degrees of hypoxia during development, and bulky tissues, where the availability of nitrite might be key to the maintenance of metabolism through its protective effect on the mitochondria. Finally, soil microbes can produce high levels of NO during hypoxia which raises the important question of whether plants are able to distinguish soil-derived NO from that produced endogenously.

Acknowledgements: Work on hypoxia in the KJG lab is supported by a Ramalingaswami Fellowship and an Innovative Young Biotechnologist Award from the Department of Biotechnology, Government of India. KJG also acknowledges the receipt of a Marie Curie Intra-European Fellowship for Career. Development. The authors wish to thank the UKIERI-DST fund for supporting the collaboration between KJG and LAJM. We thank Daniel Gibbs, University of Birmingham for critical reading and suggestions on figure 2.

Figure Legends:

Figure 1: Operation of the phytyglobin/nitric oxide (Pgb/NO) cycle under hypoxic conditions. The reduction of nitrite to NO occurs at complex III (bc_1), complex IV (cytochrome *c* oxidase), and the alternative oxidase (AOX). The NO diffuses to the cytosol where it is converted to nitrate (NO_3^-) by the hypoxia-induced class 1 phytyglobin (PgbO_2), which leads to the formation of metphytyglobin (MetPgb), which is reduced by metphytyglobin reductase (MetPgbR). Nitrate is then reduced by nitrate reductase (NR) to nitrite, which is imported into mitochondria by either a putative nitrite transporter (NT) or passive diffusion. NAD(P)H generated in the cytosol is oxidized by the externally facing calcium-dependent mitochondrial dehydrogenases (ND), or, after import into the mitochondria as reducing equivalents, by complex I. Cyt *c*, cytochrome *c*; IMM, inner mitochondrial membrane; IMS, intermembrane space; UQ, ubiquinone.

Figure 2: A. The role of nitric oxide (NO) and oxygen in the control of group VII ethylene response factor (ERF) stability under normoxia. Methionine amino peptidase (MetAP) cleaves the Nt-Met to reveal an Nt-Cys, which is then oxidized by plant cysteine oxidases (PCOs) to produce Nt-Cys sulfinat (shaded yellow symbol). NO potentially facilitates this oxidation. The oxidized Nt-Cys is arginylated by an arginyl-tRNA protein transferase (ATE1/2), creating a substrate for the N-end rule E3 ligase (PRT6) which leads to polyubiquitination and proteasomal degradation of ERFVII. It has been shown that NO is also required for degradation of ERFVII, and that stress induced reductions in endogenous NO levels can lead to enhanced stability of ERFVII even under normoxia (Vicente *et al.*, 2017), thus identifying NO as a signal controlling the accumulation of these proteins in response to stress.

B. Under hypoxia, high levels of NO are produced via increased activity of nitrate reductase and mitochondrial nitrite reduction. However, because ERFVII degradation also requires oxygen, these increased levels of NO are not able to degrade ERFVII. The NO generated under hypoxia can help in plant survival via aerenchyma formation, protection of mitochondria, hyponasty, regulation of reactive oxygen species (ROS), the Pgb-NO cycle, and the limited production of ATP. Some of these

adaptive responses are mediated by NO, and some are mediated by stabilization of ERFVIs or both.

References

Akimoto S, Tanaka A, Nakamura K, Shiro Y, Nakamura H. 2003. O₂-specific regulation of the ferrous heme-based sensor kinase FixL from *Sinorhizobium meliloti* and its aberrant inactivation in the ferric form. *Biochemical and Biophysical Research Communications* **304**: 136–142.

Armstrong W. 1979. Aeration in higher plants. *Advances in Botanical Research* **7**: 225–232.

Astier J, Gross I, Durner J. 2017. Nitric oxide production in plants: an update. *Journal of Experimental Botany* **69**: 3401–3411.

Bailey-Serres J, Chang R. 2005. Sensing and signalling in response to oxygen deprivation in plants and other organisms. *Annals of Botany* **96**: 507–518.

Bailey-Serres J, Lee SC, Brinton E. 2012. Waterproofing crops: effective flooding survival strategies. *Plant Physiology* **160**: 1698–1709.

Benamar A, Rolletschek H, Borisjuk L, Avelange-Macherel M-H, Curien G, Mostefai HA, Andriantsitohaina R, Macherel D. 2008. Nitrite-nitric oxide control of mitochondrial respiration at the frontier of anoxia. *Biochimica et Biophysica Acta* **1777**: 1268–1275.

Berger A, Brouquisse R, Pathak PK, Hichri I, Inderjit, Bhatia S, Boscari A, Igamberdiev AU, Gupta KJ. 2018. Pathways of nitric oxide metabolism and operation of phytohemoglobins in legume nodules: missing links and future directions *Plant, Cell & Environment* **41**: 2057–2068.

Bologa KL, Fernie AR, Leisse A, Loureiro ME, Geigenberger P. 2003. A bypass of sucrose synthase leads to low internal oxygen and impaired metabolic performance in growing potato tubers. *Plant Physiology* **132**: 2058–2072.

Borisjuk L, Macherel D, Benamar A, Wobus U, Rolletschek H. 2007. Low oxygen sensing and balancing in plant seeds: a role for nitric oxide. *New Phytologist* **176**: 813–823.

Brahimi-Horn C, Mazure N, Pouysségur J. 2005. Signalling via the hypoxia-inducible factor-1alpha requires multiple posttranslational modifications. *Cell Signalling* **17**: 1–9.

Brown GC, Cooper CE. 1994. Nanomolar concentrations of nitric oxide reversibly inhibit synaptosomal respiration by competing with oxygen at cytochrome oxidase. *FEBS Letters* **19**: 295–298.

Cassanova N, O'Brien KM, Stahl BT, McClure T, Poyton RO. 2005. Yeast flavohemoglobin, a nitric oxide oxidoreductase, is located in both the cytosol and the mitochondrial matrix: effects of respiration, anoxia, and the mitochondrial genome on its intracellular level and distribution. *Journal of Biological Chemistry* **280**: 7645–7653.

Chen R, Sun S, Wang C, Li Y, Liang Y, An F, Li C, Dong H, Yang X, Zhang J, Zuo J. 2009. The *Arabidopsis* *PARAQUAT RESISTANT2* gene encodes an S-nitrosoglutathione reductase that is a key regulator of cell death. *Cell Research* **19**: 1377–1387.

Cleeter MW, Cooper JM, Darley-Usmar VM, Moncada S, Schapira AH. 1994. Reversible inhibition of cytochrome *c* oxidase, the terminal enzyme of the mitochondrial respiratory chain, by nitric oxide. Implications for neurodegenerative diseases. *FEBS Letters* **345**: 50–54.

Clerc P, Rigoulet M, Leverve X, Fontaine E. 2007. Nitric oxide increases oxidative phosphorylation efficiency. *Journal of Bioenergetics and Biomembrane* **39**: 158–166.

Cochrane DW, Shah JK, Hebelstrup KH, Igamberdiev AU. 2017. Expression of phytohemoglobin affects nitric oxide metabolism and energy state of barley plants exposed to anoxia. *Plant Science* **265**: 124–130.

Cogliati S, Enriquez JA, Scorrano L. 2016. Mitochondrial cristae: where beauty meets functionality. *Trends in Biochemical Sciences* **41**: 261–273.

Cox MCH, Peeters AJM, Voeselek LACJ. 2006. The stimulating effects of ethylene and auxin on petiole elongation and on hyponastic curvature are independent processes in submerged *Rumex palustris*. *Plant, Cell & Environment* **29**: 282–290.

Cvetkovska M, Vanlerberghe GC. 2012. Alternative oxidase modulates leaf mitochondrial concentrations of superoxide and nitric oxide. *New Phytologist* **195**: 32–39.

de Oliveira HC, Wulff A, Saviani EE, Salgado I. 2008. Nitric oxide degradation by potato tuber mitochondria: evidence for the involvement of external NAD(P)H dehydrogenases. *Biochimica et Biophysica Acta* **1777**: 470–476.

Delledonne M, Xia Y, Dixon RA, Lamb C. 1998. Nitric oxide functions as a signal in plant disease resistance. *Nature* **6**: 585–588.

Drew MC, He C, Morgan PW. 2000. Programmed cell death and aerenchyma formation in roots. *Trends in Plant Science* **3**: 123–127.

Fukao T, Barrera-Figueroa BE, Juntawong P, Peña-Castro JM. 2019. Submergence and waterlogging stress in plants: A review highlighting research opportunities and understudied aspects. *Frontiers in Plant Science* **10**: 340.

Frungillo L, de Oliveira JF, Saviani EE, Oliveira HC, Martínez MC, Salgado I. 2013. Modulation of mitochondrial activity by S-nitrosoglutathione reductase in *Arabidopsis thaliana* transgenic cell lines. *Biochimica et Biophysica Acta* **1827**: 239–247.

Frungillo L, Skelly MJ, Loake GJ, Spoel SH, Salgado L. 2014. S-nitrosothiols regulate nitric oxide production and storage in plants through the nitrogen assimilation pathway. *Nature Communications* **5**: 5401.

Gardner PR. 2005. Nitric oxide dioxygenase function and mechanism of flavohemoglobin, hemoglobin, myoglobin and their associated reductases. *Journal of Inorganic Biochemistry* **99**: 247–266.

Geigenberger P, Fernie AR, Gibon Y, Christ M, Stitt M. 2000. Metabolic activity decreases as an adaptive response to low internal oxygen in growing potato tubers. *Biological Chemistry* **381**: 723–740.

Gibbs DJ, Bacardit J, Bachmair A, Holdsworth MJ. 2014a. The eukaryotic N-end rule pathway: conserved mechanisms and diverse functions. *Trends in Cell Biology* **24**: 603–611.

Gibbs DJ, Conde JV, Berckhan, S, Prasad G, Mendiando GM, Holdsworth MJ. 2015. Group VII ethylene response factors coordinate oxygen and nitric oxide signal transduction and stress responses in plants. *Plant Physiology* **169**: 23–31.

Gibbs DJ, Md Isa N, Movahedi M, Lozano-Juste J, Mendiando GM, Berckhan S, Marín-de la Rosa N, Vicente Conde J, Sousa Correia C, Pearce SP et al. 2011. Homeostatic response to hypoxia is regulated by the N-end rule pathway in plants. *Nature* **479**: 415–418.

Gibbs DJ, Md Isa N, Movahedi M, Lozano-Juste J, Mendiando GM, Berckhan S, Marín-de la Rosa N, Vicente Conde J, Sousa Correia C, Pearce SP et al. 2014b. Nitric oxide sensing in plants is mediated by proteolytic control of group VII ERF transcription factors. *Molecular Cell* **53**: 369–370.

Gibbs DJ, Tedds HM, Labandera AM, Bailey M, White MD, Hartman S, Sprigg C, Mogg SL, Osborne R, Dambire C, Boeckx T, Paling Z, Voesenek LACJ, Flashman E, Holdsworth MJ. 2018 Oxygen-dependent proteolysis regulates the stability of angiosperm polycomb repressive complex 2 subunit VERNALIZATION 2. *Nature Communications* **9**: 5438.

Graciet E, Walter F, Maoiléidigh DÓ, Pollmann S, Meyerowitz EM, Varshavsky A, Wellmer F. 2009. The N-end rule pathway controls multiple functions during Arabidopsis shoot and leaf development. *Proceedings of the National Academy of Science USA* **106**: 13618–13623.

Gupta KJ, Fernie AR, Kaiser WM, van Dongen JT. 2011. On the origins of nitric oxide. *Trends in Plant Science* **16**: 160–168.

Gupta KJ, Hebelstrup KH, Kruger NJ, Ratcliffe RG. 2014. Nitric oxide is required for homeostasis of oxygen and reactive oxygen species in barley roots under aerobic conditions. *Molecular Plant* **7**: 747–750.

Gupta KJ, Igamberdiev AU. 2011. The anoxic plant mitochondrion as a nitrite: NO reductase. *Mitochondrion* **11**: 537–543.

Gupta KJ, Lee CP, Ratcliffe RG. 2017. Nitrite protects mitochondrial structure and function under hypoxia. *Plant and Cell Physiology* **58**: 175–183.

Gupta KJ, Shah JK, Brotman Y, Jahnke K, Willmitzer L, Kaiser WM, Bauwe H, Igamberdiev AU. 2012. Inhibition of aconitase by nitric oxide leads to induction of the alternative oxidase and to a shift of metabolism towards biosynthesis of amino acids. *Journal of Experimental Botany* **63**: 1773–1784.

Gupta KJ, Stoimenova M, Kaiser WM. 2005. In higher plants, only root mitochondria, but not leaf mitochondria reduce nitrite to NO, *in vitro* and *in situ*. *Journal of Experimental Botany* **56**: 2601–2609.

Gupta KJ, Brotman Y, Segu S, Zeier T, Zeier J, Persijn ST, Cristescu SM, Harren FJM, Bauwe H, Fernie AR. 2013. The form of nitrogen nutrition affects resistance against *Pseudomonas syringae* pv. *phaseolicola* in tobacco. *Journal of Experimental Botany* **64**: 553–568.

Hebelstrup KH, Østergaard-Jensen E, Hill RD. 2008. Bioimaging techniques for subcellular localization of plant hemoglobins and measurement of hemoglobin-dependent nitric oxide scavenging in planta. *Methods in Enzymology* **437**: 595–604.

Hebelstrup KH, van Zanten M, Mandon J, Voeselek LACJ, Harren FJM, Cristescu SM, Møller IM, Mur LAJ. 2012. Haemoglobin modulates NO emission and hyponasty under hypoxia-related stress in *Arabidopsis thaliana*. *Journal of Experimental Botany* **63**: 5581–5591.

Hinz M, Wilson IW, Yang J, Buerstenbinder K, Llewellyn D, Dennis ES, Sauter M, and Dolferus R. 2010. Arabidopsis RAP2.2: An ethylene response transcription factor that is important for hypoxia survival. *Plant Physiology* **153**: 757–772.

Holman TJ, Jones PD, Russell L, Medhurst A, Ubeda Tomás S, Talloji P, Marquez J, Schmutts H, Tung SA, Taylor I et al. 2009. The N-end rule pathway promotes seed germination and establishment through removal of ABA sensitivity in *Arabidopsis*. *Proceedings of the National Academy of Science USA* **106**: 4549–4554.

Horchani F, Prévot M, Boscari A, Evangelisti E, Meilhoc E, Bruand C, Raymond P, Boncompagni E, Aschi-Smiti S, Puppo A et al. 2011. Both plant and bacterial nitrate reductases contribute to nitric oxide production in *Medicago truncatula* nitrogen-fixing nodules. *Plant Physiology* **155**: 1023–1036.

Hu RG, Sheng J, Qi X, Xu Z, Takahashi TT, Varshavsky A. 2005. The N-end rule pathway as a nitric oxide sensor controlling the levels of multiple regulators. *Nature* **437**: 981–986.

Igamberdiev AU, Bykova NV, Hill RD. 2006. Nitric oxide scavenging by barley hemoglobin is facilitated by a monodehydroascorbate reductase-mediated ascorbate reduction of methemoglobin. *Planta* **223**: 1033–1040.

Igamberdiev AU, Hill RD. 2004. Nitrate, NO and haemoglobin in plant adaptation to hypoxia: an alternative to classic fermentation pathways. *Journal of Experimental Botany* **55**: 2473–2482.

Kumari A, Wany A, Mishra S, Gupta AK, Gupta KJ. 2016. Nitric oxide measurement from purified enzymes and estimation of scavenging activity by gas phase chemiluminescence. *Methods in Molecular Biology* **1424**: 31–38.

Leterrier M, Chaki M, Valderrama R, Palma JM, Barroso JB, Corpas FJ. 2011. Function of S-nitrosoglutathione reductase (GSNOR) in plant development and under biotic/abiotic stress. *Plant Signaling & Behavior* **6**: 789-793.

Libourel IGL, Van Bodegom PM, Fricker MD, Ratcliffe RG. 2006. Nitrite reduces cytoplasmic acidosis under anoxia. *Plant Physiology* **142**: 1710–1717.

Licausi F, Kosmacz M, Weits DA, Giuntoli B, Giorgi FM, Voesenek LACJ, Perata P, van Dongen, JT. 2011. Oxygen sensing in plants is mediated by an N-end rule pathway for protein destabilization. *Nature* **479**: 419–422.

Licausi F, van Dongen JT, Giuntoli B, Novi G, Santaniello A, Geigenberger P, Perata P. 2010. HRE1 and HRE2, two hypoxia-inducible ethylene response factors, affect anaerobic responses in *Arabidopsis thaliana*. *Plant Journal* **62**: 302–315.

Liu L, Zeng M, Hausladen A, Heitman J, Stamler JS. 2000. Protection from nitrosative stress by yeast flavohemoglobin. *Proceedings of the National Academy of Sciences USA* **97**: 4672–4676.

Liu Y, Shi L, Ye N, Liu R, Jia W, Zhang J. 2009. Nitric oxide-induced rapid decrease of abscisic acid concentration is required in breaking seed dormancy in *Arabidopsis*. *New Phytologist* **183**: 1030–1042.

Liu Z, Shi X, Li S, Hu G, Zhang L, Song X. 2018. Tapetal-delayed programmed cell death (PCD) and oxidative stress-induced male sterility of *Aegilops uniaristata* cytoplasm in wheat. *International Journal of Molecular Sciences* **19**: 1708.

Millar AH, Day DA. 1996. Nitric oxide inhibits the cytochrome oxidase but not the alternative oxidase of plant mitochondria. *FEBS Letters* **398**: 155–158.

Millar AH, Day DA, Bergersen FJ. 1995. Microaerobic respiration and oxidative-phosphorylation by soybean nodule mitochondria: implications for nitrogen fixation. *Plant, Cell and Environment* **18**: 715–726.

Mira MM, Hill RD, Stasolla C. 2016a. Regulation of programmed cell death by phytooglobins. *Journal of Experimental Botany* **67**: 5901–5908.

Mira MM, Hill RD, Stasolla C. 2016b. Phytooglobins improve hypoxic root growth by alleviating apical meristem cell death. *Plant Physiology* **172**: 2044–2056.

Mugnai S, Azzarello E, Baluška F, Mancuso S. 2012. Local root apex hypoxia induces NO-mediated hypoxic acclimation of the entire root. *Plant Cell Physiology* **53**: 912–920.

Narsai R, Secco D, Schultz MD, Ecker JR, Lister R, Whelan J. 2017. Dynamic and rapid changes in the transcriptome and epigenome during germination and in developing rice (*Oryza sativa*) coleoptiles under anoxia and re-oxygenation. *Plant Journal* **89**: 805–824.

Palmieri C, Lindermayr C, Bauwe H, Steinhauser C, Durner J. 2010. Regulation of plant glycine decarboxylase by S-nitrosation and glutathionylation. *Plant Physiology*. **152**: 1514–1528.

Pierik R, Djakovic-Petrovic T, Keuskamp DH, de Wit M, Voeselek LACJ. 2009. Auxin and ethylene regulate elongation responses to neighbor proximity signals independent of gibberellin and DELLA proteins in Arabidopsis. *Plant Physiology* **149**: 1701–1712.

Planchet E, Gupta KJ, Sonoda M, Kaiser WM. 2005. Nitric oxide emission from tobacco leaves and cell suspensions: rate-limiting factors and evidence for the involvement of mitochondrial electron transport. *Plant Journal* **41**: 732–743.

Polko JK, Voeselek LACJ, Peeters AJM, Pierik R. 2011. Petiole hyponasty: an ethylene-driven, adaptive response to changes in the environment. *AoB Plants* plr031.

Pucciariello C, Perata P. 2017. New insights into reactive oxygen species and nitric oxide signalling under low oxygen in plants. *Plant, Cell and Environment* **40**: 473–482.

Reddie KG, Carroll KS. 2008. Expanding the functional diversity of proteins through cysteine oxidation. *Current Opinion in Chemical Biology* **12**: 746–754.

Ricard B, Couee I, Raymond P, Saglio PH, Saint-Ges V, Pradet A. 1994. Plant metabolism under hypoxia and anoxia. *Plant Physiology and Biochemistry* **32**: 1–10.

Royo B, Moran JF, Ratcliffe RG, Gupta KJ. 2015. Nitric oxide induces the alternative oxidase pathway in *Arabidopsis* seedlings deprived of inorganic phosphate. *Journal of Experimental Botany* **66**: 6273–6280.

Sasiharan R, Hartman S, Liu Z, Martopawiro S, Sajeev N, van Veen H, Yeung E, Voeselek LACJ. 2018. Signal dynamics and interactions during flooding stress. *Plant Physiology* **176**: 1106–1117.

Shah JK, Cochrane DW, De Paepe R, Igamberdiev AU. 2013. Respiratory complex I deficiency results in low nitric oxide levels, induction of hemoglobin and upregulation of fermentation pathways. *Plant Physiology and Biochemistry* **63**: 185–190.

Shingaki-Wells R, Millar AH, Whelan J, Narsai R. 2014. What happens to plant mitochondria under low oxygen? An omics review of the responses to low oxygen and reoxygenation. *Plant, Cell & Environment* **37**: 2260–2277.

Tossi V, Amenta M, Lamattina L, Cassia R. 2011. Nitric oxide enhances plant ultraviolet-B protection up-regulating gene expression of the phenylpropanoid biosynthetic pathway. *Plant, Cell & Environment* **34**: 909–921.

Varshavsky A. 2011. The N-end rule pathway and regulation by proteolysis. *Protein Science* **20**: 1298–1345.

Vartapetian BB, Andreeva IN, Generozova IP, Polyakova LI, Masloval, Dolgikh YI, Stepanova AY. 2003. Functional electron microscopy in studies of plant response and adaptation to anaerobic stress. *Annals of Botany* **91**: 155–172.

Vergara R, Parada F, Rubio S, Pérez FJ. 2012. Hypoxia induces H₂O₂ production and activates antioxidant defence system in grapevine buds through mediation of H₂O₂ and ethylene. *Journal of Experimental Botany* **63**: 4123–4131.

Vicente J, Mendiondo GM, Movahedi M, Peirats-Llobet M, Juan YT, Shen YY, Dambire C, Smart K, Rodriguez PL, Charng YY, Gray JE, Holdsworth MJ. 2017. The Cys-Arg/N-End rule pathway is a general sensor of abiotic stress in flowering plants. *Current Biology* **27**: 3183–3190.

Vishwakarma A, Kumari A, Mur LAJ, Gupta KJ. 2018. A discrete role for alternative oxidase under hypoxia to increase nitric oxide and drive energy production. *Free Radical Biology and Medicine* **122**: 40-51.

Voesenek LACJ, Banga AM, Thier RH, Mudde CM, Harren FJM, Barendse GWM, Blom CWPM. 1993. Submergence-induced ethylene synthesis, entrapment, and growth in two plant species with contrasting flooding resistances. *Plant Physiology* **103**: 783–791.

Vreeburg RAM, Benschop JJ, Peeters AJM, Colmer TD, Ammerlaan AHM, Staal M, Elzenga TM, Staals RHJ, Darley CP, McQueen-Mason SJ et al. 2005. Ethylene regulates fast apoplastic acidification and expansin A transcription during submergence-induced petiole elongation in *Rumex palustris*. *Plant Journal* **43**: 597–610.

Wang Y, Loake GJ, Chu C. 2013. Cross-talk of nitric oxide and reactive oxygen species in plant programmed cell death. *Frontiers in Plant Science* **4**: 314

Wany A, Gupta KJ. 2018. Reactive oxygen species, nitric oxide production and antioxidant gene expression during development of aerenchyma formation in wheat. *Plant Signaling & Behaviour* **13**: e1428515.

Wany A, Kumari A, Gupta KJ. 2017. Nitric oxide is essential for the development of aerenchyma in wheat roots under hypoxic stress. *Plant, Cell & Environment* **40**: 3002–3017.

Wany A, Gupta AK, Kumari A, Mishra S, Singh N, Pandey S, Vanvari R, Igamberdiev AU, Fernie AR, Gupta KJ. 2019. Nitrate nutrition influences multiple factors in order to increase energy efficiency under hypoxia in arabidopsis. *Annals of Botany* **123**: 691–705.

Weits DA, Giuntoli B, Kosmacz M, Parlanti S, Hubberten HM, Riegler H, Hoefgen R, Perata P, van Dongen JT, Licausi F. 2014. Plant cysteine oxidases control the oxygen-dependent branch of the N-end-rule pathway. *Nature Communications* **6**: 3425.

White MD, Klecker M, Hopkinson RJ, Weits DA, Mueller C, Naumann C, O'Neill R, Wickens J, Yang J, Brooks-Bartlett JC, Garman EF, Grossmann TN, Dissmeyer N, Flashman E. 2017. Plant cysteine oxidases are dioxygenases that directly enable arginyl transferase-catalysed arginylation of N-end rule targets. *Nature Communications* **8**: 14690.

Accepted Article

Yamauchi T, Watanabe K, Fukazawa A, Mori H, Abe F, Kawaguchi K, Oyanagi A, Nakazono M. 2014. Ethylene and reactive oxygen species are involved in root aerenchyma formation and adaptation of wheat seedlings to oxygen-deficient conditions. *Journal of Experimental Botany* **65**: 261–273.

Yun BW, Feechan A, Yin M, Saidi NB, Le Bihan T, Yu M, Moore JW, Kang JG, Kwon E, Spoel SH et al. 2011. S-nitrosation of NADPH oxidase regulates cell death in plant immunity. *Nature* **478**: 264–268.

Zhan N, Wang C, Chen L, Yang H, Feng J, Gong X, Ren B, Wu R, Mu J, Li Y, Liu Z, Zhou Y, Peng J, Wang K, Huang X, Xiao S, Zuo J. 2018. S-Nitrosation targets GSNO reductase for selective autophagy during hypoxia responses in plants. *Molecular Cell* **71**: 142–154.

Fig 1

