

Aberystwyth University

Cupid, a cell permeable peptide derived from amoeba, capable of delivering GFP into a diverse range of species

Fenton, Daniel; Phillips, Dylan; Maddison, Anne; H. George, Christopher; Ryves, Jonathan; D. Jones, Huw

Published in:
Scientific Reports

DOI:
[10.1038/s41598-020-70532-x](https://doi.org/10.1038/s41598-020-70532-x)

Publication date:
2020

Citation for published version (APA):

Fenton, D., Phillips, D., Maddison, A., H. George, C., Ryves, J., & D. Jones, H. (2020). Cupid, a cell permeable peptide derived from amoeba, capable of delivering GFP into a diverse range of species. *Scientific Reports*, 10(1), Article 13725. <https://doi.org/10.1038/s41598-020-70532-x>

Document License CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


tel: +44 1970 62 2400
email: is@aber.ac.uk

Supplementary Figures

Manuscript title:


Cupid, a cell permeable peptide derived from amoeba, capable of delivering GFP into a diverse range of species.

Supplemental Figure 1.




Supplemental Fig 1. Z-Slice green fluorescent confocal imaging of confluent mouse cardiomyocytes at 1 hour after 5 μ M Cupid-GFP addition. Consecutive Z-Slices from top of cells are shown from left to right to the base of the slide. Scale bar 20 μ m

Supplemental Figure 2.



Supplemental Fig 2. Onion epidermis imaged 1 hour after addition of either water (**A, B**) or 5 μ M native fluorescent Cupid-GFP (**C, D**). Images **A** & **C** show merged phase contrast and green fluorescence channels. **B** & **D** show green fluorescence alone. Scale bar 50 μ m.


Supplemental Figure 3.



Supplemental Fig 3.

UV-Vis absorption spectra of denatured Cupid-GFP (50 μ M) after incubation in a range of cell-culture media for over 24h at room temperature (RT) or 37°C. Native (folded and fluorescent) Cupid-GFP incubated in water at RT or 37°C for 24h also included. Y-axis: absorbance units, X-axis: nanometers.

Supplemental Fig 4.



Supplemental Fig 4.

Full-length SDS-PAGE gels showing that purified Cupid-GFP and H-GFP migrated as single bands at the expected mass of of 32.3 kD and 29.2 kD respectively.

Supplemental Video. After treatment of a living *Phymocycles blakesleanus* mycelia network within a 5 mm deep agar block with 5 μ M Cupid-GFP for 90 minutes, a Z-plane time-lapse video (8 sec/frame) was taken over the course of 1 minute 42 sec. **Video 1** with phase contrast; **Video 2** with green fluorescent filters; and **Video 3** combined overlay.