

Aberystwyth University

Co-expression of lupanine hydroxylase and pyrroloquinoline quinone 2 leads to assembled and active recombinant lupanine hydroxylase in the 3 Escherichia coli periplasm

Stampolidis, Pavlos; Kaderbhai, Naheed; Bryant, David; Winters, Ana; Gallagher, Joe; Kaderbhai, Mustak Ali

Published in:
Archives of Microbiology

DOI:
[10.13140/RG.2.1.4937.7449](https://doi.org/10.13140/RG.2.1.4937.7449)

Publication date:
2013

Citation for published version (APA):
Stampolidis, P., Kaderbhai, N., Bryant, D., Winters, A., Gallagher, J., & Kaderbhai, M. A. (2013). Co-expression of lupanine hydroxylase and pyrroloquinoline quinone 2 leads to assembled and active recombinant lupanine hydroxylase in the 3 Escherichia coli periplasm. *Archives of Microbiology*.
<https://doi.org/10.13140/RG.2.1.4937.7449>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

**Co-expression of lupanine hydroxylase and pyrroloquinoline
quinone
leads to assembled and active recombinant lupanine
hydroxylase in the
Escherichia coli periplasm**


6

Journal:	<i>Archives of Microbiology</i>
Manuscript ID:	ArchMicr-2013-0102.R1
Type of Paper Published:	Short Communication
Date Submitted by the Author:	06-Jun-2013
Complete List of Authors:	Kaderbhai, Naheed; Aberystwyth University, IBERS Stampolidis, Pavlos; Max Planck Institute of Biochemistry, 1Department of Molecular Biology Bryant, David; Aberystwyth university, IBERS Gallagher, Joseph; Aberystwyth university, IBERS Winters, Ana; Aberystwyth university, IBERS
Keywords:	pyrroloquinoline quinone, Escherichia coli periplasmic space, lupanine hydroxylase

Fig. 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

a



b






Fig. 2

a



b



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **SHORT COMMUNICATION**

2 **Co-expression of lupanine hydroxylase and pyrroloquinoline quinone**
3 **leads to assembled and active recombinant lupanine hydroxylase in the**
4 ***Escherichia coli* periplasm**

5
6 Pavlos Stampolidis¹ • Naheed N. Kaderbhai² • David Bryant² • Ana Winters², Joseph
7 Gallagher² • Mustak A. Kaderbhai*

8
9 ¹Department of Molecular Biology, Max Planck Institute of Biochemistry, 82152,
10 Martinsried, Germany

11
12 ²Institute of Biological, Environmental and Rural Sciences, Plas Gogerddan Campus,
13 William Davies Building, Aberystwyth University, Aberystwyth, Ceredigion, SY23 3EE,
14 United Kingdom

15
16 *Dr Mustak A. Kaderbhai is deceased

17
18 **Correspondence:** Naheed N. Kaderbhai; Tel: +44 1970 622294; Fax: +44 1970 823242;
19 e-mail: nnk@aber.ac.uk

20
21 **Keywords:** pyrroloquinoline quinone, lupanine hydroxylase, *Escherichia coli*,
22 periplasmic space, quinohaemoprotein, protein export

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

24 ABSTRACT

25 Lupanine hydroxylase (LH) is a quinohaemoprotein responsible for the conversion of the
26 alkaloid, lupanine to 17-hydroxylupanine. Previous attempts to express the enzyme in
27 *Escherichia coli* required *in vitro* addition of the co-factor pyrroloquinoline quinone
28 (PQQ) and posed some impediments on subsequent structural studies for further
29 characterization of the enzyme. An *E. coli* clone with LH and cytochrome *c* maturation
30 operon was transformed with a third plasmid containing the PQQ operon from *Klebsiella*
31 *pneumoniae*, *luh* gene and resulted in the production of periplasmically-targeted,
32 correctly folded, PQQ and haem inserted active enzyme.

33 Interestingly, LH was less active than the *in vitro* incorporated PQQ-LH, presumably due
34 to the incorporation of PQQ precursors in the periplasm. This is a first report of an active
35 LH enzyme with *in vivo* incorporation of PQQ in *E. coli* and provides the necessary tool
36 for further enzyme structural characterization.

37

38

39

40

41

42

43

44

45

46 Introduction

1
2
3 47 Pyrroloquinoline quinone [4,5-dihydro-4,5-dioxo-1*H*-pyrrolo-[2,3-*f*] quinoline-
4
5 48 2,7,9-tricarboxylic acid (PQQ) is an important non-covalently bound redox cofactor in
6
7
8 49 enzymes such as dehydrogenases, oxygenases and decarboxylases (Anthony 2001) along
9
10 50 with NAD(P) and flavins (Duine and Jongejan 1989a). It is an aromatic, tricyclic, *ortho*-
11
12 51 quinone and is a crucial link between compound oxidation and the respiratory chain
13
14 52 (Duine and Jongejan 1989b). Enzymes containing PQQ as a cofactor are divided into two
15
16 53 classes; Class I quinoproteins contain PQQ and Ca²⁺ as cofactors and class II
17
18 54 quinohaemoproteins, in addition to PQQ and Ca²⁺ also carry haem at their C-terminus
19
20 55 (Stoorvogel et al. 1996). In Class I quinoproteins, catalytic conversion of the substrate
21
22 56 results in PQQ accepting two electrons (PQQH₂) which are then transferred to another
23
24 57 redox protein (Davidson 2004), whereas Class II quinohaemoproteins transfer electrons
25
26 58 from PQQ and then relay onto the haem moiety within the molecule before being donated
27
28 59 to an external acceptor (Oubrie et al. 2002).

30
31
32
33
34 60 In eukaryotes, the presence of PQQ has so far been confirmed in mammalian
35
36 61 tissues and milk, however, no enzymatic reaction which directly necessitates PQQ has
37
38 62 been reported yet (Steinberg et al. 2003).

39
40
41 63 Studies in micro-organisms have revealed that a number of PQQ maturation
42
43 64 chaperones are required for its synthesis and the genes coding for these factors are all
44
45 65 organized into clusters that exhibit a high level of sequence homology (Meulenberg et al.
46
47 66 1990). Examples are *Methylobacterium extorquens* possessing seven PQQ maturation
48
49 67 chaperones, *Klebsiella pneumoniae* has six and *Acinetobacter calcoaceticus* has only four
50
51 68 chaperones (Puehringer et al. 2008).
52
53
54
55
56
57
58
59
60

1
2
3
4 69 Interestingly, there is an absence of PQQ in some prokaryotes as a number of
5
6 70 bacterial species are reported to express quino-enzymes lacking PQQ. An example in
7
8 71 *Escherichia coli* is expression of glucose dehydrogenase minus PQQ (Matsushita et al.
9
10 72 1997) and the reason as to why *E. coli* would produce an enzyme whilst unable to
11
12 73 produce its cofactor remains a mystery. One theory is that *E. coli* is in possession of an
13
14 74 alternative pathway of PQQ production with a yet unidentified inducer (Biville et al.
15
16 75 1991). A second theory is that *E. coli* scavenges PQQ from its environment and the
17
18 76 finding that mobile *E. coli* strains move chemo-tactically towards PQQ reinforces this
19
20 77 argument (DeJonge et al. 1996). Although the complete biosynthetic pathway of PQQ
21
22 78 remains elusive, tyrosine and glutamate (Fig. 1a) are the two main precursors
23
24 79 (Magnusson et al. 2004).

25
26
27
28
29 80 LH is a class II, type I quinohaemoprotein which contains equimolar amounts of
30
31 81 PQQ, Ca²⁺ and haem as co-factors (Stampolidis et al. 2009). It is a 72 kDa monomeric
32
33 82 enzyme responsible for the initial conversion of the alkaloid lupanine to 17-
34
35 83 hydroxylupanine (Toczko et al. 1963). Heterologous expression of the *luh* gene in *E. coli*
36
37 84 also necessitated co-expression of the cytochrome *c* maturation machinery (Thony-Meyer
38
39 85 2003), responsible for haem insertion into the apoform of the enzyme (Stampolidis et al
40
41 86 2009). Based on these factors, we investigated the co-expression and *in vivo*
42
43 87 incorporation of a third plasmid coding for PQQ into the above *E.coli* clone co-
44
45 88 expressing LH and cytochrome *c* maturation machinery.
46
47
48
49
50

51 89

52 53 54 90 **Materials and methods**

55 56 91 **Bacterial strains, growth conditions and plasmids**

57
58
59
60

1
2
3 92 Starter cultures of *E. coli* TB1 [F^- *ara* $\Delta(lac-proAB)$ ($\phi 80dlac$ $\Delta(lacZ)M15$) *rpsL(Str^r)* *thi*
4
5 93 *hsdR*] (New England BioLabs) clones harbouring plasmids pEC86 (cytochrome *c*
6
7 94 maturation operon), pEV-LH32 (untagged LH) and pINK-LH-His₄ (tetra-His tagged LH)
8
9 95 were cultured in Luria Bertani broth (1% (w/v) Tryptone, 0.5% (w/v) yeast extract and
10
11 96 1% (w/v) NaCl) containing 75 $\mu\text{g mL}^{-1}$ ampicillin and/or 50 $\mu\text{g mL}^{-1}$ chloramphenicol
12
13 97 were grown to saturation for 16h at 30⁰C and applied as 2% (v/v) inoculum for batch
14
15 98 cultivation in the MOPS medium (Karim et al. 1993) with the appropriate antibiotics and
16
17 99 orbital agitation at 125 rev min⁻¹ for 18h at 21⁰C. The clone pK-187-PQQ, pEC86/
18
19 100 pK187-PQQ and pEC86/ pINK-LH-His₄/ pK187-PQQ for PQQ synthesis, was induced in
20
21 101 MOPS in the presence of 1% (w/v) Na-gluconate, 50 μM IPTG and kanamycin at 50 μg
22
23 102 mL⁻¹ and grown as above.
24
25
26
27
28
29

30 103 Plasmid pBCP165 was provided by J. C. Arents (Velterop et al. 1995) and
31
32 104 plasmids pEV-LH32, pINK-LH-His₄ and pEC86 were from Dr M. A. Kaderbhai
33
34 105 laboratory.
35
36

37 106

39 107 **Periplasmic extraction from clones**

40
41 108 Periplasmic extract was prepared by the osmotic shock method as described previously
42
43 109 (Kaderbhai et al. 2012) and was recovered by centrifugation at 10,000g for 5min and
44
45 110 stored frozen at -80⁰C.
46
47

48 111

51 112 **His-tagged LH purification using Qiagen Ni-NTA**

52
53 113 His-tagged LH was routinely purified from periplasmic extracts of clone pEC86/pINK-
54
55 114 LH-His₄ via passage through Qiagen Ni-NTA column and eluted with 100mM imidazole
56
57
58
59
60

1
2
3 115 (pH 8.0) and 100mM EDTA. Untagged LH was purified from the pEC86/pEV-LH32
4
5
6 116 clone on DE-52-DEAE-cellulose columns as described previously (Stampolidis et al.
7
8 117 2009).

9
10 118

119 **PQQ purification using DEAE Sepharose Fast Flow**

120 A 10mL volume of periplasmic extract from 2L culture of pKK-187-PQQ (5ml) was
121 mixed with DEAE Sepharose slurry (10ml) and was gently shaken overnight at 4⁰C. The
122 suspension was packed in a column, washed twice with 0.5M KPO₄ (pH 7.4) and eluted
123 in 2mL total volume of 2M NaCl in 25mM Tris-HCl (pH 8). The salt was removed from
124 the eluent by PQQ binding to a 2mL bed volume of C18 reverse phase column followed
125 by two washes with 5mM HCl (pH 2.0) and final elution with 20% (v/v) methanol. PQQ
126 was detected by diode-array detection and ESI-MS as described in Comont et al. (2012).

127

128 **Identification of PQQ**

129 Presence of PQQ in the isolates was detected spectrophotometrically at 257nm
130 absorbance.

131

132 **Measurement of *in vivo* and *in vitro* LH enzyme activity**

133 LH was routinely activated on addition of 4mM CaCl₂ and incubation at room
134 temperature for 1h. Activation of LH apoform *in vitro* necessitated addition of 200μM
135 PQQ. LH was assayed at 25⁰C using horse heart cytochrome *c* as the electron acceptor by
136 increase in absorbance at 550nm due to its reduction as described by Stampolidis et al.

1
2
3 137 (2009). Assays were performed in duplicates, the reaction rate being linear during this
4
5 138 time.
6
7

8 139

10 140 **Results**

13 141 **Construction and expression of PQQ in *E. coli***

14 142 Plasmid pBCP165 comprising a 6.7 Kbp PQQ operon under the transcriptional control of
15
16
17
18 143 a *K. pneumoniae pqq* promoter was cleaved with *NheI* (blunted) and *HindIII* restriction
19
20 144 endonucleases (Fig. 1b). The resulting fragment allowed closer proximity of the operon
21
22
23 145 to the exogenous *lac* promoter present in the host plasmid. Colony PCR with primers,
24
25 146 For-PQQ-Screen: 5'-GCCATCCTGCGGCAGC-3' and Rev-PQQ-Screen: 5'-
26
27 147 CCCCCGGCCATTAATCCC-3' using part of the PQQ operon (For-PQQ-Screen) and
28
29
30 148 part of the plasmid vector (Rev-PQQ-Screen) along with gene mapping with *EcoRI* and
31
32 149 *HindIII* confirmed the presence of the entire PQQ operon in a positive clone designated
33
34 150 as pK187-PQQ.
35
36

37 151 A red eluted band was obtained from the periplasmic extract of pKK187-PQQ
38
39 152 after passage through a DEAE Sepharose Fast Flow column and gave a spectral scan with
40
41 153 a major peak at 257nm, characteristic of PQQ. Comparison of this spectrum with a
42
43 154 control spectrum of PQQ from *Methylophilus methylotrophus* showed no significant
44
45
46 155 difference indicating that the PQQ produced in pKK187-PQQ was of good quality and
47
48 156 purity (Fig. 2b).
49
50

51 157 The PQQ from pKK187-PQQ showed a major peak that eluted at 17 minutes
52
53 158 following separation by C₁₈ reverse phase chromatography and had a molecular mass of
54
55
56 159 331.1 *m/z* [M+H]⁺ using ESI-MS (supplementary data 1).
57
58
59
60

1
2
3 160
4
56 161 **Co-expression of PQQ and LH in *E.coli***
7

8 162 A fully functional LH with *in vivo* PQQ incorporation was obtained with the construction
9
10 163 of a three plasmid system. Isolated plasmid pK187-PQQ (Kam^r), LH encoding pINK-LH-
11
12 164 His₄ (Amp^r) and cytochrome *c* maturation machinery encoding pEC86 (Cm^r) were co-
13
14
15 165 transformed into *E. coli* TB1 cells to generate clone pEC86/ pINK-LH-His₄/ pK187-
16
17 166 PQQ. The TB1 strain, a JM83 derivative, carries the *hsdR* mutation which facilitates
18
19
20 167 transformation efficiency and is reported to provide plasmid stability in conjunction to
21
22 168 protein expression and purification (Yanisch-Perron et al. 1985; Belo et al.
23
24 169 1996). Successful transformants were expressed in the MOPS phosphate-limited medium
25
26
27 170 containing 1% (w/v) Na-gluconate and 50μM IPTG in the presence of the three
28
29 171 antibiotics (see Materials and Methods) for 18h at 21⁰C. Periplasmic extracts from
30
31 172 osmotically shocked cells were prepared as described in the Materials and Methods
32
33
34 173 section.
35

36 174
3738
39 175 **Comparative activity of *in vivo* PQQ incorporated LH apoenzyme and *in vitro***
40
41 176 **activated LH apoenzyme**
42

43
44 177 Comparison of tetra-His-tagged LH activity with the pEC86/pEV-LH32 untagged LH
45
46 178 showed no significant difference in enzymatic activity (Table 1). Comparison of the
47
48 179 purified LH activity from clone pEC86/ pINK-LH-His₄/ pK187-PQQ with LH from clone
49
50 180 pEC86/ pINK-LH-His₄ showed *in vivo* activated form exhibited 10% of the activity of *in*
51
52 181 *vitro* activated form. Periplasmic extracts from clones pEC86, pK187-PQQ and pEC86/
53
54
55
56
57
58
59
60

1
2
3 182 pK187-PQQ were also assayed for LH activity as controls and gave no significant
4
5 183 enzyme activity.
6
7

8 184 Interestingly, although *in vivo* form of LH exhibits only 10% activity of its *in vitro*
9
10 185 counterpart, comparison of the PQQ levels of production between the two respective host
11
12 186 clones appeared to be similar.
13
14

15 187

18 188 **Discussion**

19
20 189 As a result of the inability of *E. coli* to endogenously produce PQQ, initial efforts to
21
22 190 express quinoproteins, heterologously, in a simple and inexpensive way were impeded. In
23
24 191 an effort to tackle this issue, cloning of the entire operon for PQQ from *Klebsiella*
25
26 192 *pneumonia* to *E. coli* was carried out (Meulenberg et al. 1990). Expression of the 6.7 kbp
27
28 193 operon consisting of genes *pqq A, B, C, D, E* and *F* resulted in *in vivo* incorporation of
29
30 194 PQQ into glucose dehydrogenase and subsequent optimization by Velterop and co-
31
32 195 workers (Velterop et al. 1995) with incorporation of plasmid pBCP165 in *E. coli* strain
33
34 196 *JA221* resulted in greater yields in the region of 180nM.
35
36
37
38

39 197 Periplasmic targeting of LH in an earlier study (Stampolidis et al. 2009) resulted
40
41 198 in homogeneous production of the apoform of the enzyme and was readily transformed
42
43 199 into holoform upon subsequent *in vitro* addition of PQQ. However, this approach posed
44
45 200 some impediments to further work in characterization of the enzyme. Preliminary
46
47 201 attempts to examine the effect of replacing Ca^{2+} in the active site of the enzyme with
48
49 202 other ions proved inconclusive due to micro quantities of other ions in PQQ preparation.
50
51 203 Moreover, X-ray diffraction data from crystals of *in vitro* correctly folded LH apoform
52
53 204 failed to provide meaningful diffraction patterns.
54
55
56
57
58
59
60

1
2
3 205 In this study, co-expression of the engineered LH and cytochrome *c* with PQQ
4
5
6 206 resulted in the production of active LH with successful *in vivo* incorporation of PQQ. The
7
8 207 10% activity of the *in vivo* incorporated PQQ-LH might be due to the incorporation of
9
10 208 PQQ precursors into the active site of the enzyme. This occurrence of PQQ precursors is
11
12 209 not uncommon and has been observed in alcohol dehydrogenase from *Pseudomonas*
13
14 210 *testosteroni* with similar effects on the enzyme activity and no change in the quaternary
15
16 211 structure of the protein (Jongejan et al. 1989). Another probable cause for this
17
18 212 phenomenon could be extracellular targeting of PQQ which limits cofactor availability in
19
20 213 the periplasm for apoform incorporation.
21
22
23

24 214 Future attempts to produce LH crystals suitable for X-ray diffraction will
25
26 215 facilitate studies on the structure of quinohaemoproteins and increase our understanding
27
28 216 on the catalytic breakdown of the alkaloid, lupanine. The outcome of our approach is a
29
30 217 first report of *in vivo* recombinant LH apoenzyme synthesized in *E. coli* and provides the
31
32 218 necessary tools for further LH enzymology studies.
33
34
35

36 219

37 220 **Acknowledgement:**

38
39 221 This work was supported by the European Regional Development Fund through funding
40
41 222 provided by the Welsh Government (BEACON: 80561), by ERDF (A4B HE 06 COL
42
43 223 1024) and BBSRC (SPG 03134CFC).
44
45
46
47

48 224

49 225 **Conflict of interest:**

50
51 226 The authors declare no conflict of interest.
52
53
54

55 227

56 228 **References**

57
58
59
60

- 1
2
3 229 Anthony C (2001) Pyrroloquinoline quinone (PQQ) and quinoprotein enzymes.
4 230 Antioxidants and Redox Signalling 3:757-774
5 231 Belo I, Santos JA, Cabral JM, Mota M (1996) Optimization study of *Escherichia coli*
6 232 TB1 cell disruption for cytochrome b5 recovery in a small-scale bead mill.
7 233 Biotechnol Prog 12:201-204
8 234 Biville F, Turlin E, Gasser F (1991) Mutants of *Escherichia coli* producing
9 235 pyrroloquinoline quinone. J Gen Microbiol 137:1775-1782
10 236 Comont DC, Winters AL and Gwynn-Jones D (2012) Acclimation and interaction
11 237 between drought and elevated UV-B in *A. thaliana*: Differences in response over
12 238 treatment, recovery and reproduction. Ecology and evolution 2(11):2695-2709
13 239 Davidson V (2004) Electron transfer in quinoproteins. Arch Biochem Biophys 428:32-40
14 240 DeJonge R, DeMattos MJT, Stock JB, Neijssel OM (1996) Pyrroloquinoline quinone, a
15 241 chemotactic attractant for *Escherichia coli*. Journal of Bacteriology 178:1224-
16 242 1226
17 243 Duine JA, Jongejan JA (1989a) Pyrroloquinoline quinone: a novel cofactor. Vitam Horm
18 244 45:223-262
19 245 Duine JA, Jongejan JA (1989b) Quinoproteins, enzymes with pyrrolo-quinoline quinone
20 246 as cofactor. Annu Rev Biochem 58:403-426
21 247 Kaderbhai MA, Hopper DJ, Akhtar KM, Abbas, SK and Kaderbhai NN (2003) A
22 248 Cytochrome c from a Lupanine-Transforming *Pseudomonas putida* Strain Is
23 249 Expressed in *Escherichia coli* during Aerobic Cultivation and Efficiently
24 250 Exported and Assembled in the Periplasm. Applied and Environmental
25 251 Microbiology 69(8):4727-4731
26 252 Jongejan JA, Groen BW, Duine JA (1989) Structural properties of PQQ involved in the
27 253 activity of quinohaemoprotein alcohol dehydrogenase from *Pseudomonas*
28 254 *testosteroni*. Kluwer Academic Publishers, Dordrecht, The Netherlands
29 255 Karim A, Kaderbhai N, Evans A, Harding V, Kaderbhai MA (1993) Efficient bacterial
30 256 export of a eukaryotic cytoplasmic cytochrome. Biotechnology (N Y) 11:612-618
31 257 Magnusson OT, Toyama H, Saeki M, Schwarzenbacher R, Klinman JP (2004) The
32 258 structure of a biosynthetic intermediate of pyrroloquinoline quinone (PQQ) and
33 259 elucidation of the final step of PQQ biosynthesis. Journal of the American
34 260 Chemical Society 126:5342-5343
35 261 Matsushita K, Arents JC, Bader R, Yamada M, Adachi O, Postma PW (1997)
36 262 *Escherichia coli* is unable to produce pyrroloquinoline quinone (PQQ).
37 263 Microbiology 143 (Pt 10):3149-3156
38 264 Meulenberg JJM, Sellink E, Loenen WAM, Riegman NH, Vankleef M, Postma PW
39 265 (1990) Cloning of *Klebsiella-Pneumoniae* Pqq Genes and Pqq Biosynthesis in
40 266 *Escherichia-coli*. Fems Microbiology Letters 71:337-344
41 267 Oubrie A, Rozeboom HJ, Kalk KH, Huizinga EG, Dijkstra BW (2002) Crystal structure
42 268 of quinohemoprotein alcohol dehydrogenase from *Comamonas testosteroni* -
43 269 Structural basis for substrate oxidation and electron transfer. Journal of Biological
44 270 Chemistry 277:3727-3732
45 271 Puehringer S, Metlitzky M, Schwarzenbacher R (2008) The pyrroloquinoline quinone
46 272 biosynthesis pathway revisited: A structural approach BMC Biochemistry 9

- 1
2
3 273 Stampolidis P, Kaderbhai NN, Kaderbhai MA (2009) Periplasmically-exported lupanine
4 274 hydroxylase undergoes transition from soluble to functional inclusion bodies in
5 275 *Escherichia coli*. Archives of Biochemistry and Biophysics 484:8-15
6
7 276 Stampolidis P, Kaderbhai NN, Kaderbhai MA (2012) Presence and role of a second
8 277 disulphide bond in recombinant lupanine hydroxylase using site-directed
9 278 mutagenesis with 143Cys-->Ser and 124,143Cys-->Ser mutations in *Escherichia*
10 279 *coli*. FEMS Microbiol Lett 334:35-43
11
12 280 Steinberg F, Stites TE, Anderson P, Storms D, Chan I, Eghbali S, Rucher R. (2003)
13 281 Pyrroloquinoline quinone improves growth and reproductive performance in mice
14 282 fed chemically defined diets. Experimental Biology and Medicine 228:160-166
15
16 283 Stoorvogel J, Kraayveld DE, vanSluis CA, Jongejan JA, deVries S, Duine JA (1996)
17 284 Characterization of the gene encoding quinohaemoprotein ethanol dehydrogenase
18 285 of *Comamonas testosteroni*. European Journal of Biochemistry 235:690-698
19
20 286 Thony-Meyer L (2003) A heme chaperone for cytochrome c biosynthesis. Biochemistry
21 287 42:13099-13105
22
23 288 Toczko M, Brzeski W, Kakolewska-Baniuk A (1963) Microbial degradation of lupanine.
24 289 V. Identification of 17-hydroxylupanine. B Pol Acad Sci XI:161-164
25
26 290 Velterop JS, Sellink E, Meulenbergh JJM, David S, Bulder I, Postma PW (1995) Synthesis
27 291 of Pyrroloquinoline Quinone *in-vivo* and *in-vitro* and Detection of an Intermediate
28 292 in the Biosynthetic-Pathway. Journal of Bacteriology 177:5088-5098
29
30 293 Yanisch-Perron C, Vieira J, Messing J (1985) Improved M13 phage cloning vectors and
31 294 host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. Gene
32 295 33:103-119
33
34 296
35 297

298 **Figure Legends**

299 **Fig. 1: Generation of PQQ in *E. coli*.**

300 **a) Precursors of PQQ biosynthesis.** Glutamate and tyrosine have been identified as the
301 two precursors of PQQ. PQQ maturation requires at least four different chaperones

302 **b) The PQQ operon in pBCP165.** ORFX is schematic diagram of an open-reading frame
303 coding for an unidentified protein (not essential for PQQ expression), Ppqq; PQQ
304 promoter and A-F; PQQ maturation genes.

305

306 **Fig. 2: PQQ spectra.**

1
2
3 307 Spectral comparison of PQQ purified preparation from **a)** *Methylophilus methylotrophus*
4
5 308 (control) and **b)** *Escherichia coli* pK187-PQQ.
6
7
8 309
9
10 310
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3
4
5 311 **Table 1: Lupanine hydroxylase (LH) enzyme activity in *Escherichia coli* clones in the presence and/or absence of PQQ and**
6
7 312 ***Cytc* operons.**
8


Clone name and description	LH expression	PQQ production	LH activity (units/mg periplasmic protein)
pEC86/pEV-LH32-untagged LH and <i>Cytc</i> operon	+	-	66
pEC86/pINK-LH-His ₄ tagged- <i>Cytc</i> operon	+	-	54
pEC86/pINK-LH-His ₄ tagged /pK187-PQQ and <i>Cytc</i> operons	+	+	6.8
pEC86- <i>Cytc</i> operon	-	-	0.15
pK187-PQQ operon	-	+	0
pEC86/pK187-PQQ and <i>Cytc</i> operons	-	+	0.30

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30 313

31
32 314 Signs represent presence (+) or absence (-) of LH expression and PQQ production.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


RT: 14.94 - 18.06



NL:
1.93E3
Total Scan
PDA
Naheed1


For Peer Review

Naheed1 #4979-5183 RT: 16.59-17.27 AV: 205 NL: 1.05E4 microAU



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RT: 14.76 - 18.56




NL:
1.23E4
TIC F: ITMS + c
ESI Full ms2
331.00@35.00 [
90.00-400.00]
MS Naheed1


For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Naheed1 #722-747 RT: 16.75-17.18 AV: 9 NL: 6.33E3
4 F: ITMS + c ESI Full ms2 331.00@35.00 [90.00-400.00]




RT: 14.76 - 18.56



NL:
1.76E4
TIC F: ITMS - c
ESI Full ms2
329.00@35.00 [
90.00-400.00]
MS Naheed1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Naheed1 #727-746 RT: 16.83-17.16 AV: 7 NL: 1.02E4
4 F: ITMS - c ESI Full ms2 329.00@35.00 [90.00-400.00]



15% SDS-PAGE OF pEC86, pKK-187, pEC86/pEVLH32 (1), pEC86/pINK-LH-His₄ (2)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

