

Aberystwyth University

Existence results for a nonlinear transmission problem

Dalla Riva, Matteo; Mishuris, Gennady

Published in:

Journal of Mathematical Analysis and Applications

DOI:

[10.1016/j.jmaa.2015.05.019](https://doi.org/10.1016/j.jmaa.2015.05.019)

Publication date:

2015

Citation for published version (APA):

Dalla Riva, M., & Mishuris, G. (2015). Existence results for a nonlinear transmission problem. *Journal of Mathematical Analysis and Applications*, 430(2), 718-741. <https://doi.org/10.1016/j.jmaa.2015.05.019>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Existence results for a nonlinear transmission problem

M. Dalla Riva^{a,*}, G. Mishuris^b

^a*Center for Research and Development in Mathematics and Applications (CIDMA) -
University of Aveiro - Portugal*

^b*Department of Mathematics - Aberystwyth University - UK*

Abstract

Let Ω^o and Ω^i be open bounded regular subsets of \mathbb{R}^n such that the closure of Ω^i is contained in Ω^o . Let f^o be a regular function on $\partial\Omega^o$ and let F and G be continuous functions from $\partial\Omega^i \times \mathbb{R}$ to \mathbb{R} . By exploiting an argument based on potential theory and on the Leray-Schauder principle we show that under suitable and completely explicit conditions on F and G there exists at least one pair of continuous functions (u^o, u^i) such that

$$\begin{cases} \Delta u^o = 0 & \text{in } \Omega^o \setminus \text{cl}\Omega^i, \\ \Delta u^i = 0 & \text{in } \Omega^i, \\ u^o(x) = f^o(x) & \text{for all } x \in \partial\Omega^o, \\ u^o(x) = F(x, u^i(x)) & \text{for all } x \in \partial\Omega^i, \\ \nu_{\Omega^i} \cdot \nabla u^o(x) - \nu_{\Omega^i} \cdot \nabla u^i(x) = G(x, u^i(x)) & \text{for all } x \in \partial\Omega^i, \end{cases}$$

where the last equality is attained in certain weak sense. A simple example shows that such a pair of functions (u^o, u^i) is in general neither unique nor locally unique. If instead the fourth condition of the problem is obtained by a small nonlinear perturbation of a homogeneous linear condition, then we prove the existence of at least one classical solution which is in addition locally unique.

Keywords: nonlinear transmission problem, systems of nonlinear integral equations, fixed-point theorem, potential theory

*Corresponding author. Telephone: +351-234-372545. Complete postal address: Departamento de Matemática, Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

Email addresses: matteo.dallariva@gmail.com (M. Dalla Riva), ggm@aber.ac.uk (G. Mishuris)

URL: <https://sites.google.com/site/matteodallariva/> (M. Dalla Riva)

Preprint submitted to Journal of Mathematical Analysis and Applications November 11, 2015

1. Introduction

We investigate the existence of solutions for a boundary value problem with a nonlinear transmission condition. In order to define such a boundary value problem we introduce some notation. We fix once for all

a natural number $n \in \mathbb{N}$, $n \geq 2$, and a real number $\alpha \in]0, 1[$,

where \mathbb{N} denotes the set of natural numbers including 0. Then we fix two sets Ω^o and Ω^i in the n -dimensional Euclidean space \mathbb{R}^n . The letter ‘ o ’ stands for ‘outer domain’ and the letter ‘ i ’ stands for ‘inner domain’. We assume that Ω^o and Ω^i satisfy the following condition:

Ω^o and Ω^i are open bounded subsets of \mathbb{R}^n of class $C^{1,\alpha}$, $\text{cl}\Omega^i \subseteq \Omega^o$, and the boundaries $\partial\Omega^o$ and $\partial\Omega^i$ are connected.

For the definition of functions and sets of the usual Schauder class $C^{0,\alpha}$ and $C^{1,\alpha}$, we refer for example to Gilbarg and Trudinger [20, §6.2]. Here and in the sequel $\text{cl}\Omega$ denotes the closure of Ω for all $\Omega \subseteq \mathbb{R}^n$. Then we fix a function $f^o \in C^{1,\alpha}(\partial\Omega^o)$ and two continuous functions F and G from $\partial\Omega^i \times \mathbb{R}$ to \mathbb{R} and we consider the following nonlinear transmission boundary value problem for a pair of functions (u^o, u^i) in $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$,

$$\left\{ \begin{array}{ll} \Delta u^o = 0 & \text{in } \Omega^o \setminus \text{cl}\Omega^i, \\ \Delta u^i = 0 & \text{in } \Omega^i, \\ u^o(x) = f^o(x) & \text{for all } x \in \partial\Omega^o, \\ u^o(x) = F(x, u^i(x)) & \text{for all } x \in \partial\Omega^i, \\ \nu_{\Omega^i} \cdot \nabla u^o(x) - \nu_{\Omega^i} \cdot \nabla u^i(x) = G(x, u^i(x)) & \text{for all } x \in \partial\Omega^i, \end{array} \right. \quad (1)$$

where ν_{Ω^i} denotes the outer unit normal to the boundary $\partial\Omega^i$. Our aim is to determine suitably general and completely explicit conditions on F and G which ensure the existence of solutions of problem (1).

The analysis of problems such as (1) is motivated by the role played in continuum mechanics. In particular, nonlinear transmission conditions of this kind arise in the study of composite structures glued together by thin adhesive layers which are thermally or mechanically very different from the components’ constituents. In modern material technology such composites

are widely used (see, *e.g.*, the second named author, Miszuris, and Öchsner [34, 35] and Rosselli and Carbutt [40]), but the numerical treatment of the mathematical model by finite elements methods is still difficult, requires the introduction of highly inhomogeneous meshes, and often leads to poor accuracy and numerical instability (see, *e.g.*, Babuška and Suri [1]). A convenient way to overcome this problem is to replace the thin layers by zero thickness interfaces between the composite's components. Then one has to define on such interfaces suitable transmission conditions which incorporates the thermal and mechanical properties of the original layers. Such a procedure can be rigorously justified by an asymptotic method and leads to the introduction of boundary value problems with nonlinear transmission conditions such as those in (1) (see for example Miszuris and Öchsner [36] and the references therein).

We observe that the existence of solutions of nonlinear boundary value problems has been largely investigated by means of variational techniques (see, *e.g.*, the monographs of Nečas [39] and of Roubíček [41] and the references therein). In fact, under some restrictive assumptions on the functions F and G , the existence of solutions of our problem (1) could be deduced by exploiting some known results. In particular, if it happens that problem (1) can be reformulated into an equation of the form $-\operatorname{div}A(x, U)\nabla U = 0$, where A is a suitable Carathéodory function and the unknown function U belongs to the Sobolev space $H^1(\Omega^o)$ and satisfies a Dirichlet condition on $\partial\Omega^o$, then the existence and uniqueness of a solution can be directly deduced by the results of Hlaváček, Křížek and Malý in [21]. This is for example the case when $G = 0$ and the function $F(x, t)$ of $(x, t) \in \partial\Omega^i \times \mathbb{R}$ is constant with respect to x , is differentiable with respect to t , and the partial differential $\partial_t F(x_0, \cdot)$ is Lipschitz continuous and satisfies the inequality $1/c < \partial_t F(x_0, t) < c$ for a constant $c > 0$ and for all $t \in \mathbb{R}$ (here x_0 is a fixed point of $\partial\Omega^i$).

In this paper instead, we exploit a method based on potential theory to rewrite problem (1) into a suitable nonlinear system of integral equations which can be analysed by a fixed-point theorem. Potential theoretic techniques have been largely exploited in literature to study existence and uniqueness problems for linear or semilinear partial differential equations with non linear boundary conditions. In particular, as far back as in 1921 Carleman [5] has considered the existence of harmonic functions u in a domain Ω which satisfy a non-linear Robin condition $\nu_\Omega(x) \cdot \nabla u(x) = H(x, u(x))$ on the boundary $\partial\Omega$. Since then, such a problem has received the attention of many authors such as Leray [30] (see also Jacob [22]), Nakamori and Suyama [38],

Kilngelhöfer [23, 24], Cushing [8], Efendiev, Schmitz, and Wendland [11], and Kohr, Lanza de Cristoforis, and Wendland [25]. In the case of domains with a small hole we also mention the nonlinear Robin problem for the Laplace operator investigated in Lanza de Cristoforis [27] and the nonlinear traction problem in elasticity addressed in [10]. Moreover, an approach based on coupling of boundary integral and finite element methods has been developed in order to study exterior nonlinear boundary value problems with transmission conditions, we mention for example the papers of Berger [3], Berger, Warnecke, and Wendland [4], Costabel and Stephan [7], and Gatica and Hsiao [18, 19]. In particular, Barrenechea and Gatica considered in [2] the case when the jump of the normal derivative across the interface boundary depends nonlinearly on the Dirichlet data. Boundary integral methods have been applied also by Mityushev and Rogosin for the analysis of transmission problems in the two dimensional plane (cf. [37, Chap. 5]). Finally, we mention the nonlinear transmission problem in a domain with a small inclusion investigated by Lanza de Cristoforis in [28] and the periodic analog studied by Lanza de Cristoforis and Musolino in [29].

2. Description of the main results

We now describe the main results of the present paper. We will exploit the following notation: if H is a function from $\partial\Omega^i \times \mathbb{R}$, then we denote by \mathcal{F}_H the nonlinear non-autonomous composition operator which takes a function f from $\partial\Omega^i$ to \mathbb{R} to the function $\mathcal{F}_H f$ defined by

$$\mathcal{F}_H f(x) \equiv H(x, f(x)) \quad \forall x \in \partial\Omega^i .$$

Since the functions F and G which define the nonlinear condition in (1) are assumed to be continuous from $\partial\Omega^i \times \mathbb{R}$ to \mathbb{R} , one easily verifies that \mathcal{F}_F and \mathcal{F}_G are continuous from $C^0(\partial\Omega^i)$ to itself. Then we consider the following condition:

$$\begin{aligned} &\text{the composition operator } (I_{\Omega^i} + \mathcal{F}_F) \text{ has a continuous} \\ &\text{inverse } (I_{\Omega^i} + \mathcal{F}_F)^{(-1)} \text{ from } C^0(\partial\Omega^i) \text{ to itself.} \end{aligned} \tag{2}$$

Here I_{Ω^i} denotes the identity operator from $C^0(\partial\Omega^i)$ to itself. We observe that for the validity of condition (2) it is not required that the function which takes t to $F(x, t)$ is monotone for all fixed $x \in \partial\Omega^i$. In addition, we introduce

a condition on the growth of F and G : we assume that

$$\begin{aligned} &\text{there exist } c_1, c_2 \in]0, +\infty[, \delta_1 \in]1, +\infty[, \text{ and } \delta_2 \in [0, 1[\text{ such that} \\ &|F(x, t)| \geq c_1 |t|^{\delta_1} - (1/c_1) \quad \text{and} \\ &|G(x, t)| \leq c_2 (1 + |F(x, t)|)^{\delta_2} \quad \forall (x, t) \in \partial\Omega^i \times \mathbb{R}. \end{aligned} \tag{3}$$

The first condition in (3) is a super-linear growth condition for F , while the second one is a sub-linear growth condition for G with respect to F (which is a strictly weaker condition than the standard sub-linear growth condition $|G(x, t)| \leq c_2 (1 + |t|)^{\delta_2}$).

By exploiting an argument based on the invariance of the Leray-Schauder topological degree we show in our main Theorem 4.11 that conditions (2) and (3) imply the existence of at least one pair of continuous functions $(\tilde{u}^o, \tilde{u}^i) \in C^0(\text{cl}\Omega^o \setminus \Omega^i) \times C^0(\text{cl}\Omega^o)$ which satisfies the first four equations of (1) in the classical sense and fulfils the fifth condition in a certain weak sense which will be clarified (see Definition 4.8 and problem (25) below). However, the conditions in (2) and (3) do not imply neither the uniqueness nor the local uniqueness of the pair $(\tilde{u}^o, \tilde{u}^i)$.

This last fact can be evidenced in a simple example. Take $\Omega^o = R\mathbb{B}_n$, $\Omega^i = r\mathbb{B}_n$, with $r, R \in \mathbb{R}$, $r < R$, and $\mathbb{B}_n \equiv \{x \in \mathbb{R}^n : |x| < 1\}$. Then assume that f^o is constant and identically equal to a real number $t^o \in \mathbb{R}$ and that $F(x, t) \equiv f(t)$ and $G(x, t) \equiv g(t)$ for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$, where f and g are continuous functions from \mathbb{R} to \mathbb{R} . We set

$$\Gamma_n(t) \equiv \begin{cases} \frac{1}{2\pi} \log t & \text{if } n = 2, \\ \frac{t^{2-n}}{s_n(2-n)} & \text{if } n \geq 3, \end{cases} \quad \forall t \in]0, +\infty[,$$

where s_n denotes the $(n-1)$ -dimensional measure of $\partial\mathbb{B}_n$ (thus $\Gamma_n(|x|) = S_n(x)$ with S_n the standard fundamental solution of Δ , see also definition (9) below). Then the pair of functions (u^o, u^i) defined by

$$\begin{aligned} u^o(x) &= t^o - \frac{\Gamma_n(R) - \Gamma_n(|x|)}{\Gamma'_n(r)} g(t^i) \quad \forall x \in \text{cl}\Omega^o \setminus \Omega^i, \\ u^i(x) &= t^i \quad \forall x \in \text{cl}\Omega^i \end{aligned} \tag{4}$$

is a solution of problem (1) for all $t^i \in \mathbb{R}$ which are solutions of the equation

$$t^o - \frac{\Gamma_n(R) - \Gamma_n(r)}{\Gamma'_n(r)} g(t^i) = f(t^i). \tag{5}$$




Figure 1: the intersections of the blue graph with the red line correspond to solutions of equation (5)

Now take

$$f(t) \equiv t^3 - 2t^2 + t + 1 \quad \forall t \in \mathbb{R}$$

and assume that g is constant. One immediately verifies that the corresponding functions F and G satisfy the conditions in (2) and (3). In addition, if t^o , R , r , and g are chosen in such a way that the left hand side of (5) is equal to 1, then equation (5) has two solutions: $t^i = 0$ and $t^i = 1$ (see Fig. 1). Accordingly, the corresponding problem (1) has at least two different solutions provided by (4). If instead $f(t) \equiv t^3 - 2t^2 + t + 1$ for $t < 0$ and $t > 1$ and $f(t) \equiv 1$ for $t \in [0, 1]$, then every t^i in $[0, 1]$ is a solution of (5) and the corresponding solutions of problem (1) are not locally unique in any reasonable topology.

We observe that our main existence Theorem 4.11 shows the existence of pair of functions $(\tilde{u}^o, \tilde{u}^i) \in C^0(\text{cl}\Omega^o \setminus \Omega^i) \times C^0(\text{cl}\Omega^i)$ which are solutions of problem (1) in a certain ‘weak’ sense but it would be preferable to have classical solutions in $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ (or at least in $H^1(\Omega^o \setminus \text{cl}\Omega^i) \times H^1(\Omega^i)$). Thus, it is natural to ask what further conditions should one impose on F and G in order to obtain such a regularity. In Theorem 4.12 we show that, if

$$(I_{\Omega^i} + \mathcal{F}_F)^{(-1)} \text{ and } \mathcal{F}_G \text{ map } C^{0,\alpha}(\partial\Omega^i) \text{ to itself,} \quad (6)$$

then problem (1) has at least one weak solution in $C^{0,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{0,\alpha}(\text{cl}\Omega^o)$. However, in order to obtain solutions in $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ by exploiting our argument it does not suffice to increase the regularity of F and G and it seems that a different approach should be implemented.

To illustrate this fact, we consider in the last Section 5 the case when the fourth condition of problem (1) is a small nonlinear perturbation of a homogenous linear condition. Namely, we assume that $F(x, t) = \lambda t + \epsilon \Phi(x, t)$ for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$, where λ is a positive real constant, ϵ is a small real parameter, and Φ is a continuous function from $\partial\Omega^i \times \mathbb{R}$ to \mathbb{R} . Then we consider the nonlinear transmission problem

$$\begin{cases} \Delta u^o = 0 & \text{in } \Omega^o \setminus \text{cl}\Omega^i, \\ \Delta u^i = 0 & \text{in } \Omega^i, \\ u^o(x) = f^o(x) & \text{for all } x \in \partial\Omega^o, \\ u^o(x) = \lambda u^i(x) + \epsilon \Phi(x, u^i(x)) & \text{for all } x \in \partial\Omega^i, \\ \nu_{\Omega^i} \cdot \nabla u^o(x) - \nu_{\Omega^i} \cdot \nabla u^i(x) = G(x, u^i(x)) & \text{for all } x \in \partial\Omega^i, \end{cases} \quad (7)$$

for a pair of functions $(u^o, u^i) \in C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$. In Theorem 5.10 below we show that, under suitable assumptions on Φ , G , and Ω^i (see condition (30)), there exists $\epsilon_* > 0$ such that problem (7) has a solution $(u_\epsilon^o, u_\epsilon^i) \in C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ for all $\epsilon \in] - \epsilon_*, \epsilon_*[$. Such a solution $(u_\epsilon^o, u_\epsilon^i)$ is locally unique in $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ for all fixed $\epsilon \in] - \epsilon_*, \epsilon_*[$ and, in addition, the map which takes ϵ to $(u_\epsilon^o, u_\epsilon^i)$ is continuously Fréchet differentiable from $] - \epsilon_*, \epsilon_*[$ to $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$. However, Theorem 5.10 does not provide any estimate for the value of ϵ_* . Therefore, the existence conditions provided by Theorem 5.10 are not completely explicit, as instead are those of Theorem 4.11.

We also observe that the assumptions of Theorems 4.11 and 5.10 may be simultaneously verified, but the solutions provided by Theorem 4.11 may not coincide with those provided by Theorem 5.10. Consider for example the case introduced here above where $\Omega^o = R\mathbb{B}_n$ and $\Omega^i = r\mathbb{B}_n$, with $r < R$. Assume that $f^o(x) = t^o$, $\Phi(x, t) = \phi(t)$, and $G(x, t) = g(t)$ for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$, where $t^o \in \mathbb{R}$ and ϕ and g are continuous functions from \mathbb{R} to itself. Then we look for solutions of problem (7) in the form (4) with $t^i \in \mathbb{R}$ solution of the equation

$$t^o - \frac{\Gamma_n(R) - \Gamma_n(r)}{\Gamma_n(r)} g(t^i) = \lambda t + \epsilon \phi(t^i). \quad (8)$$




Figure 2: the blue lines are graphs of $\lambda t + \epsilon\phi(t)$ for $\epsilon = 3/2$, $\epsilon = 1$, and $\epsilon = 1/2$. The intersections of the blue graphs with the red line correspond to solutions of (8)

Now we take $\lambda \equiv 1/2$,

$$\phi(t) \equiv t^3 - 2t^2 + \frac{1}{2}t + 1 \quad \forall t \in \mathbb{R},$$

and g constant. One can choose t^o , r , R , and g in such a way that the left hand side of (8) is equal to 1. Then it is easily verified that equation (8) has two solutions for $\epsilon = 1$: $t^i = 0$ and $t^i = 1$. Instead, for $\epsilon > 1$ we only have the solutions provided by Theorem 4.11 due to the behaviour at infinity of ϕ and we loose the solution provided by Theorem 5.10 due to the smallness of ϵ (see Fig. 2). Similar examples can be exhibited to show that the local uniqueness of the solution guaranteed by Theorem 5.10 for ϵ small can be lost when we increase ϵ .

Finally, we observe that potential theoretic methods have been developed for the analysis of linear transmission problems in Lipschitz domains. We mention, for example, the works of Escauriaza, Fabes, and Verchota [12], Escauriaza and Seo [13], and Escauriza and Mitrea [14]. However, the argument used in the present paper for the proof of the main Theorem 4.11 cannot be immediately extended to the case of a Lipschitz contact boundary $\partial\Omega^i$. The reason is that the compactness of the double layer operator W_{Ω^i} plays a crucial role in the proof of Proposition 4.6, where we apply the Leray-Schauder principle to prove that the fixed point equation (16) has solutions

(see Section 3 for the definition of W_{Ω^i}). As is well known, W_{Ω^i} is compact in $L^p(\partial\Omega^i)$, $p \in]1, +\infty[$, if Ω^i is at least of class C^1 , but may be not compact if Ω^i is just a Lipschitz domain (cf., *e.g.*, Fabes, Jodeit, and Lewis [15] and Fabes, Jodeit, and Rivière [16]). Since here we assume Ω^i to be of class $C^{1,\alpha}$, we also have W_{Ω^i} compact from $C^0(\partial\Omega^i)$ to itself, from $C^{0,\alpha}(\partial\Omega^i)$ to itself, and from $C^{1,\alpha}(\partial\Omega^i)$ to itself (cf. Section 3 below).

The paper is organised as follows. Section 3 is a section of preliminaries where we introduce some classical notion of potential theory. In Section 4 we prove our main Theorem 4.11 where we show the existence of continuous solutions of problem (25). Finally, in Section 5 we consider problem (7) and we show the existence of locally unique $C^{1,\alpha}$ solutions for ϵ small.

3. Classical notions of potential theory

We denote by S_n the function from $\mathbb{R}^n \setminus \{0\}$ to \mathbb{R} defined by

$$S_n(x) \equiv \begin{cases} \frac{1}{2\pi} \log |x| & \text{if } n = 2, \\ \frac{1}{s_n(2-n)} |x|^{2-n} & \text{if } n \geq 3, \end{cases} \quad \forall x \in \mathbb{R}^n \setminus \{0\}. \quad (9)$$

As is well known, S_n is a fundamental solution for the Laplace operator in \mathbb{R}^n .

Let Ω be an open bounded subset of \mathbb{R}^n of class $C^{1,\alpha}$. Let $\phi \in L^2(\partial\Omega)$. Then $v_\Omega[\phi]$ denotes the single layer potential with density ϕ . Namely,

$$v_\Omega[\phi](x) \equiv \int_{\partial\Omega} \phi(y) S_n(x-y) d\sigma_y \quad \forall x \in \mathbb{R}^n,$$

where $d\sigma$ denotes the area element on $\partial\Omega$. As is well known, if $\phi \in L^\infty(\partial\Omega)$, then $v_\Omega[\phi]$ is a continuous function from \mathbb{R}^n to \mathbb{R} . In addition, if $\phi \in C^{0,\alpha}(\partial\Omega)$, then the restrictions $v_\Omega^+[\phi] \equiv v_\Omega[\phi]|_{\text{cl}\Omega}$ and $v_\Omega^-[\phi] \equiv v_\Omega[\phi]|_{\mathbb{R}^n \setminus \Omega}$ belong to $C^{1,\alpha}(\text{cl}\Omega)$ and to $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^n \setminus \Omega)$, respectively. Here $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^n \setminus \Omega)$ denotes the space of functions on $\mathbb{R}^n \setminus \Omega$ whose restrictions to $\text{cl}\mathcal{O}$ belong to $C^{1,\alpha}(\text{cl}\mathcal{O})$ for all open bounded subsets \mathcal{O} of $\mathbb{R}^n \setminus \Omega$.

If $\psi \in L^2(\partial\Omega)$, then $w_\Omega[\psi]$ denotes the double layer potential with density ψ . Namely,

$$w_\Omega[\psi](x) \equiv - \int_{\partial\Omega} \psi(y) \nu_\Omega(y) \cdot \nabla S_n(x-y) d\sigma_y \quad \forall x \in \mathbb{R}^n,$$

where ν_Ω denotes the outer unit normal to $\partial\Omega$ and the symbol ‘ \cdot ’ denotes the scalar product in \mathbb{R}^n . If $\psi \in C^{1,\alpha}(\partial\Omega)$, then the restriction $w_\Omega[\psi]|_\Omega$ extends

to a function $w_\Omega^+[\psi]$ of $C^{1,\alpha}(\text{cl}\Omega)$ and the restriction $w_\Omega[\psi]|_{\mathbb{R}^n \setminus \text{cl}\Omega}$ extends to a function $w_\Omega^-[\psi]$ of $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^n \setminus \Omega)$.

Let

$$W_\Omega[\psi](x) \equiv - \int_{\partial\Omega} \psi(y) \nu_\Omega(y) \cdot \nabla S_n(x-y) d\sigma_y \quad \forall x \in \partial\Omega,$$

for all $\psi \in L^2(\partial\Omega)$, and

$$W_\Omega^*[\phi](x) \equiv \int_{\partial\Omega} \phi(y) \nu_\Omega(x) \cdot \nabla S_n(x-y) d\sigma_y \quad \forall x \in \partial\Omega,$$

for all $\phi \in L^2(\partial\Omega)$. As is well known W_Ω and W_Ω^* are compact operator from $L^2(\partial\Omega)$ to itself and are adjoint one to the other. In the sequel we denote by I_Ω the identity map from $L^2(\partial\Omega)$ to itself. Thus $\pm \frac{1}{2}I_\Omega + W_\Omega$ and $\pm \frac{1}{2}I_\Omega + W_\Omega^*$ are Fredholm operators of index 0 from $L^2(\partial\Omega)$ to itself.

We now introduce the following classical result of Schauder [42, 43]:

Lemma 3.1. *Let $\beta \in]0, 1]$. Then the map which takes ψ to $W_\Omega[\psi]$ is continuous from $C^0(\partial\Omega)$ to $C^{0,\alpha}(\partial\Omega)$ and from $C^{1,\beta}(\partial\Omega)$ to $C^{1,\alpha}(\partial\Omega)$. The map which takes ϕ to $W_\Omega^*[\phi]$ is continuous from $C^{0,\beta}(\partial\Omega)$ to $C^{0,\alpha}(\partial\Omega)$.*

As a consequence, the map which takes ψ to $W_\Omega[\psi]$ is compact from $C^0(\partial\Omega)$ to itself, from $C^{0,\alpha}(\partial\Omega)$ to itself, and from $C^{1,\alpha}(\partial\Omega)$ to itself, and the map which takes ϕ to $W_\Omega^*[\phi]$ is compact from $C^{0,\alpha}(\partial\Omega)$ to itself. Then one immediately deduces the validity of the following.

Lemma 3.2. *The operators $\pm \frac{1}{2}I_\Omega + W_\Omega$ are Fredholm of index 0 from $C^0(\partial\Omega)$ to itself, from $C^{0,\alpha}(\partial\Omega)$ to itself, and from $C^{1,\alpha}(\partial\Omega)$ to itself. The operators $\pm \frac{1}{2}I_\Omega + W_\Omega^*$ are Fredholm of index 0 from $C^{0,\alpha}(\partial\Omega)$ to itself.*

In addition we have the following technical Lemma 3.3.

Lemma 3.3. *Let $\psi \in L^2(\partial\Omega)$. Let $\beta \in [0, \alpha]$. Let $\gamma \in \mathbb{R}$. If $(\frac{1}{2}I_\Omega + \gamma W_\Omega^*)\psi$ belongs to $C^{0,\beta}(\partial\Omega)$, then $\psi \in C^{0,\beta}(\partial\Omega)$.*

Proof. If $(\frac{1}{2}I_\Omega + \gamma W_\Omega^*)\psi \in C^{0,\beta}(\partial\Omega)$, then a standard argument based on iterated kernels ensures that $\psi \in C^0(\partial\Omega)$. It follows that $W_\Omega^*\psi \in C^{0,\beta'}(\partial\Omega)$ for all $\beta' \in [0, \alpha[$ (cf. Miranda [33, Chap. II, §14, IV], see also Schauder [43]). Thus $\psi = 2(\frac{1}{2}I_\Omega + \gamma W_\Omega^*)\psi - 2\gamma W_\Omega^*\psi$ belongs to $C^{0,\beta''}(\partial\Omega)$ with $\beta'' \equiv \min\{\beta', \beta\}$ for all $\beta' \in [0, \alpha[$. Now the lemma is proved for $\beta < \alpha$. If instead

$\beta = \alpha$, then we observe that $W_\Omega^* \psi \in C^{0,\alpha}(\partial\Omega)$ by the membership of ψ in $C^{0,\beta'}(\partial\Omega)$ with $\beta' \in]0, 1]$ (cf. Lemma 3.1). Thus $\psi = 2(\frac{1}{2}I_\Omega + \gamma W_\Omega^*)\psi - 2\gamma W_\Omega^* \psi$ belongs to $C^{0,\alpha}(\partial\Omega)$ and the Lemma is proved. \square

By exploiting the operators W_Ω and W_Ω^* we can now write the jump formulas

$$w_\Omega^\pm[\psi]_{|\partial\Omega} = \pm \frac{1}{2}\psi + W_\Omega[\psi] \quad \text{and} \quad \nu_\Omega \cdot \nabla v_\Omega^\pm[\psi]_{|\partial\Omega} = \mp \frac{1}{2}\phi + W_\Omega^*[\psi] \quad (10)$$

which hold for all continuous function $\psi \in C^0(\partial\Omega)$ (cf., *e.g.*, Folland [17, Chap. 3]). In addition, if $\psi \in C^{1,\alpha}(\partial\Omega)$, then we have

$$\nu_\Omega \cdot \nabla w_\Omega^+[\psi]_{|\partial\Omega} = \nu_\Omega \cdot \nabla w_\Omega^-[\psi]_{|\partial\Omega}. \quad (11)$$

In the following Lemma 3.4 we describe the null-spaces $\text{Ker}(\pm \frac{1}{2}I_\Omega + W_\Omega^*)$ and $\text{Ker}(\pm \frac{1}{2}I_\Omega + W_\Omega)$ of the operators $\pm \frac{1}{2}I_\Omega + W_\Omega^*$ and $\pm \frac{1}{2}I_\Omega + W_\Omega$ in $L^2(\partial\Omega)$. To do so, we exploit the following notation: if \mathcal{X} is a subspace of $L^1(\partial\Omega)$ then we denote by \mathcal{X}_0 the subspace of \mathcal{X} consisting of the functions which have 0 integral mean. For a proof of Lemma 3.4 we refer, *e.g.*, to Folland [17, Chap. 3].

Lemma 3.4. *Let $\Omega_1, \dots, \Omega_N$ be the bounded connected components of Ω and $\Omega_0^-, \Omega_1^-, \dots, \Omega_M^-$ be the connected components of $\mathbb{R}^n \setminus \text{cl}\Omega$. Assume that $\Omega_1^-, \dots, \Omega_M^-$ be bounded and that Ω_0^- be unbounded. Then the following statements hold.*

- (i) *The map from $\text{Ker}(\frac{1}{2}I_\Omega + W_\Omega^*)$ to $\text{Ker}(\frac{1}{2}I_\Omega + W_\Omega)$ which takes μ to $v_\Omega[\mu]_{|\partial\Omega}$ is bijective.*
- (ii) *The map from $\text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega^*)_0$ to $\text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega)$ which takes μ to $v_\Omega[\mu]_{|\partial\Omega}$ is one-to-one. If $n \geq 3$, then the map from $\text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega^*)$ to $\text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega)$ which takes μ to $v_\Omega[\mu]_{|\partial\Omega}$ is bijective.*
- (iii) *$\text{Ker}(\frac{1}{2}I_\Omega + W_\Omega)$ consists of the functions from $\partial\Omega$ to \mathbb{R} which are constant on $\partial\Omega_j^-$ for all $j \in \{1, \dots, M\}$ and which are identically equal to 0 on $\partial\Omega_0^-$.*
- (iv) *$\text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega)$ consists of the functions from $\partial\Omega$ to \mathbb{R} which are constant on $\partial\Omega_j$ for all $j \in \{1, \dots, N\}$.*
- (v) *If $\phi \in \text{Ker}(\frac{1}{2}I_\Omega + W_\Omega^*)$ and $\int_{\partial\Omega} \phi \psi \, d\sigma = 0$ for all $\psi \in \text{Ker}(\frac{1}{2}I_\Omega + W_\Omega)$, then $\phi = 0$.*
- (vi) *If $\phi \in \text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega^*)$ and $\int_{\partial\Omega} \phi \psi \, d\sigma = 0$ for all $\psi \in \text{Ker}(-\frac{1}{2}I_\Omega + W_\Omega)$, then $\phi = 0$.*

Finally, we have the following technical Lemma 3.5.

Lemma 3.5. *Let $\tau \in]-1, 1[$. Then $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is an isomorphism from $L^2(\partial\Omega)$ to itself, from $C^0(\partial\Omega)$ to itself, and from $C^{0,\alpha}(\partial\Omega)$ to itself.*

Proof. To prove that $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is an isomorphism from $C^{0,\alpha}(\partial\Omega)$ to itself one can argue as in Lanza de Cristoforis and Musolino [29, Prop. A.1]. Then, to show that $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is an isomorphism from $L^2(\partial\Omega)$ to itself we observe that τW_Ω^* is compact from $L^2(\partial\Omega)$ to itself. Thus $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is a Fredholm operator of index 0 from $L^2(\partial\Omega)$ to itself and it suffices to show that $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is one-to-one. Now, if $\psi \in L^2(\partial\Omega)$ and $(\frac{1}{2}I_\Omega + \tau W_\Omega^*)\psi = 0$, then $\psi \in C^{0,\alpha}(\partial\Omega)$ by Lemma 3.3. Accordingly $\psi = 0$ by the invertibility of $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ in $C^{0,\alpha}(\partial\Omega)$. Finally, to prove that $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is an isomorphism from $C^0(\partial\Omega)$ to itself we observe that $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is continuous from $C^0(\partial\Omega)$ to itself (because W_Ω^* has a weak singularity). Moreover, if $\eta \in L^2(\partial\Omega)$ and $(\frac{1}{2}I_\Omega + \tau W_\Omega^*)\eta \in C^0(\partial\Omega)$ then Lemma 3.3 ensures that $\eta \in C^0(\partial\Omega)$. Then one deduces that $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is a bijective continuous linear operator from $C^0(\partial\Omega)$ to itself. Accordingly $\frac{1}{2}I_\Omega + \tau W_\Omega^*$ is an isomorphism from $C^0(\partial\Omega)$ to itself by the open mapping theorem. \square

4. Existence results for problem (1)

We prove in this section our main Theorem 4.11.

As a first step we deduce in the following Lemma 4.1 a representation for a pair of harmonic functions in $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ in terms of a suitable combination of layer potential. We will exploit the following notation: if Ω is an open bounded subset of \mathbb{R}^n , $k \in \mathbb{N}$, and $\beta \in [0, 1[$, then we denote by $C_{\text{harm}}^{k,\beta}(\text{cl}\Omega)$ the subspace of $C^{k,\beta}(\text{cl}\Omega)$ defined by

$$C_{\text{harm}}^{k,\beta}(\text{cl}\Omega) \equiv \{ \phi \in C^{k,\beta}(\text{cl}\Omega) : \Delta\phi = 0 \text{ in } \Omega \} . \quad (12)$$

Lemma 4.1. *The map from $C^{1,\alpha}(\partial\Omega^o) \times C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ to $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ which takes (μ^o, μ, η) to the pair $(w^o[\mu^o, \mu, \eta], u^i[\mu^o, \mu, \eta])$ given by*

$$w^o[\mu^o, \mu, \eta] \equiv (w_{\Omega^o}^+[\mu^o] + w_{\Omega^i}^-[\mu] + v_{\Omega^i}^-[\eta])|_{\text{cl}\Omega^o \setminus \Omega^i}, \quad u^i[\mu^o, \mu, \eta] \equiv w_{\Omega^i}^+[\mu]$$

is bijective.

Proof. The map is well defined. Indeed $(u^o[\mu^o, \mu, \eta], u^i[\mu^o, \mu, \eta]) \in C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ and $\Delta u^o[\mu^o, \mu, \eta] = 0$, $\Delta u^i[\mu^o, \mu, \eta] = 0$ for all $(\mu^o, \mu, \eta) \in C^{1,\alpha}(\partial\Omega^o) \times C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ (cf. Section 3). We now show that it is bijective. We take a pair of harmonic functions (ϕ^o, ϕ^i) in $C^{1,\alpha}_{\text{harm}}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}_{\text{harm}}(\text{cl}\Omega^i)$ and we prove that there exists unique $(\mu^o, \mu, \eta) \in C^{1,\alpha}(\partial\Omega^o) \times C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ such that $(u^o[\mu^o, \mu, \eta], u^i[\mu^o, \mu, \eta]) = (\phi^o, \phi^i)$. By the standard properties of the double layer potential there exists a unique $\mu \in C^{1,\alpha}(\partial\Omega^i)$ such that $w_{\Omega^i}^+[\mu] = \phi^i$ (cf. (10) and Lemma 3.4 (iii)). Then we have to show that there exists unique $(\mu^o, \eta) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ such that

$$(w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta])|_{\text{cl}\Omega^o \setminus \Omega^i} = \phi^o - w_{\Omega^i}^-[\mu]|_{\text{cl}\Omega^o \setminus \Omega^i}. \quad (13)$$

Let $\psi^o \equiv \phi^o|_{\partial\Omega^o} - w_{\Omega^i}^-[\mu]|_{\partial\Omega^o}$ and $\psi^i \equiv \nu_{\Omega^i} \cdot \nabla \phi^o|_{\partial\Omega^i} - \nu_{\Omega^i} \cdot \nabla w_{\Omega^i}^-[\mu]|_{\partial\Omega^i}$. Then $\psi^o \in C^{1,\alpha}(\partial\Omega^o)$, $\psi^i \in C^{0,\alpha}(\partial\Omega^o)$, and equation (13) is equivalent to

$$\begin{aligned} \left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)\mu^o + v_{\Omega^i}^-[\eta]|_{\partial\Omega^o} &= \psi^o, \\ \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*\right)\eta + \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]|_{\partial\Omega^i} &= \psi^i \end{aligned} \quad (14)$$

by the uniqueness of the classical solution of the Neumann-Dirichlet mixed boundary value problem (see also (10)). By Lemmas 3.2 and 3.4 the operator which takes (μ^o, η) to $((\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})\mu^o, (\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)\eta)$ is a linear isomorphism from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ to itself. Moreover, by the properties of the integral operators with real analytic kernels and no singularities, the operator which takes (μ^o, η) to $(v_{\Omega^i}^-[\eta]|_{\partial\Omega^o}, \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]|_{\partial\Omega^i})$ is compact from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ to itself. Hence, the operator which takes (μ^o, η) to $((\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})\mu^o + v_{\Omega^i}^-[\eta]|_{\partial\Omega^o}, (\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)\eta + \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]|_{\partial\Omega^i})$ is a compact perturbation of an isomorphism and therefore it is a Fredholm operator of index 0 from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ to itself. Thus, to complete the proof it suffices to show that equation (14) with $(\psi^o, \psi^i) = (0, 0)$ implies $(\mu^o, \eta) = (0, 0)$. If $((\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})\mu^o + v_{\Omega^i}^-[\eta]|_{\partial\Omega^o}, (\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)\eta + \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]|_{\partial\Omega^i}) = (0, 0)$, then by the jump properties (10) and by the uniqueness of the classical solution of the Neumann-Dirichlet mixed problem one deduces that $(w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta])|_{\text{cl}\Omega^o \setminus \Omega^i} = 0$. Hence $w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta] = 0$ in $\text{cl}\Omega^i$ by the uniqueness of the classical solution of the Dirichlet problem in Ω^i and by the continuity of $(w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta])|_{\text{cl}\Omega^o}$ (cf. Section 3). Then by (10) we have

$$\begin{aligned} \eta &= \nu_{\Omega^i} \cdot \nabla v_{\Omega^i}^-[\eta]|_{\partial\Omega^i} - \nu_{\Omega^i} \cdot \nabla v_{\Omega^i}^+[\eta]|_{\partial\Omega^i} \\ &= \nu_{\Omega^i} \cdot \nabla (w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta])|_{\partial\Omega^i} - \nu_{\Omega^i} \cdot \nabla (w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^+[\eta])|_{\partial\Omega^i} = 0. \end{aligned}$$

By (14) it follows that $(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})\mu^o = 0$ and thus $\mu^o = 0$ by Lemma 3.4 (iii). Our proof is now complete. \square

In the following Lemma 4.2 we introduce an auxiliary operator which we denote by J . In the sequel we will denote the inverse of an invertible map f with $f^{(-1)}$, as opposed to the reciprocal of a function g which will be denoted with g^{-1} .

Lemma 4.2. *We define*

$$J[\eta] \equiv \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*\right)\eta - \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)^{(-1)} v_{\Omega^i}[\eta]_{|\partial\Omega^o} \right]_{|\partial\Omega^i}$$

for all $\eta \in L^2(\partial\Omega^i)$. Then the map which takes η to $J[\eta]$ is an isomorphism from $L^2(\partial\Omega^i)$ to itself, from $C^0(\partial\Omega^i)$ to itself, and from $C^{0,\alpha}(\partial\Omega^i)$ to itself.

Proof. By the properties of integral operators with real analytic kernels and no singularity, by the invertibility of $\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}$ in $C^{1,\alpha}(\partial\Omega^o)$ (cf. Lemmas 3.2 and 3.4), and by the continuity of the map $w_{\Omega^o}^+[\cdot]$ from $C^{1,\alpha}(\partial\Omega^o)$ to $C^{1,\alpha}(\text{cl}\Omega^o)$ (cf., e.g., Miranda [32]), one deduces that the operator which takes η to

$$\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)^{(-1)} v_{\Omega^i}[\eta]_{|\partial\Omega^o} \right]_{|\partial\Omega^i} \quad (15)$$

is continuous from $L^2(\partial\Omega^i)$ to $C^{0,\alpha}(\partial\Omega^i)$. Then, by the compactness of $W_{\Omega^i}^*$ in $L^2(\partial\Omega^i)$ it follows that J is a Fredholm operator of index 0 from $L^2(\partial\Omega^i)$ to itself. Thus, to show that J is invertible from $L^2(\partial\Omega^i)$ to itself it suffices to prove that $J[\eta] = 0$ implies $\eta = 0$. If $\eta \in L^2(\partial\Omega^i)$ and $J[\eta] = 0$, then $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)\eta \in C^{0,\alpha}(\partial\Omega^i)$ by the membership of (15) in $C^{0,\alpha}(\partial\Omega^i)$. It follows that $\eta \in C^{0,\alpha}(\partial\Omega^i)$ (cf. Lemma 3.3). Then, by setting $\mu^o \equiv -(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})^{(-1)} v_{\Omega^i}[\eta]_{|\partial\Omega^o}$ and by exploiting equality (10) we verify that $u^o[\mu^o, 0, \eta]_{|\partial\Omega^o} = 0$ and $\nu_{\Omega^i} \cdot \nabla u^o[\mu^o, 0, \eta]_{|\partial\Omega^o} = 0$, where $u^o[\mu^o, 0, \eta]$ is defined as in Lemma 4.1. Accordingly $u^o[\mu^o, 0, \eta] = 0$ by the uniqueness of the solution of the mixed boundary value problem. Since $u^i[\mu^o, 0, \eta] = w_{\Omega^i}^+[0] = 0$, Lemma 4.1 implies that $\eta = 0$.

To prove that J is invertible from $C^0(\partial\Omega^i)$ to itself we observe that J is continuous from $C^0(\partial\Omega^i)$ to itself (because $W_{\Omega^i}^*$ has a weak singularity). Moreover, if $\eta \in L^2(\partial\Omega^i)$ and $J[\eta] \in C^0(\partial\Omega^i)$ then $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)\eta \in C^0(\partial\Omega^i)$ by the membership of (15) in $C^{0,\alpha}(\partial\Omega^i)$. Thus Lemma 3.3 ensures that $\eta \in C^0(\partial\Omega^i)$.

Similarly, to prove that J is invertible from $C^{0,\alpha}(\partial\Omega^i)$ to itself we observe that J is continuous from $C^{0,\alpha}(\partial\Omega^i)$ to itself and that $J[\eta] \in C^{0,\alpha}(\partial\Omega^i)$ implies $\eta \in C^{0,\alpha}(\partial\Omega^i)$ for all $\eta \in L^2(\partial\Omega^i)$ (cf. Lemma 3.3). \square

Then we have the following Lemma 4.3 where we rewrite problem (1) into an equivalent system of boundary integral equations.

Lemma 4.3. *Let condition (2) hold. Let $(\mu^o, \mu, \eta) \in C^{1,\alpha}(\partial\Omega^o) \times C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$. Then $(u^o[\mu^o, \mu, \eta], u^i[\mu^o, \mu, \eta])$ is a solution of (1) if and only if*

$$\begin{aligned} \mu^o &= \left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)^{(-1)}(f^o - w_{\Omega^i}^-[\mu]_{|\partial\Omega^o} - v_{\Omega^i}[\eta]_{|\partial\Omega^o}), \\ \mu &= \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}\right)^{(-1)} \left[(I_{\Omega^i} + \mathcal{F}_F)^{(-1)} (w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} + v_{\Omega^i}[\eta]_{|\partial\Omega^i} + 2W_{\Omega^i}\mu) \right], \\ \eta &= J^{(-1)} \left[\mathcal{F}_G \circ (I_{\Omega^i} + \mathcal{F}_F)^{(-1)} (w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} + v_{\Omega^i}[\eta]_{|\partial\Omega^i} + 2W_{\Omega^i}\mu) \right. \\ &\quad \left. - \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)^{(-1)}(f^o - w_{\Omega^i}^-[\mu]_{|\partial\Omega^o}) \right]_{|\partial\Omega^i} \right]. \end{aligned} \tag{16}$$

Proof. Note that $\nu_{\Omega^i} \cdot \nabla w_{\Omega^i}^-[\mu](x) - \nu_{\Omega^i} \cdot \nabla w_{\Omega^i}^+[\mu](x) = 0$ by the membership of μ in $C^{1,\alpha}(\partial\Omega^i)$ (cf. (11)). Then the validity of the statement is a consequence of Lemma 4.1, of the jump properties of single and double layer potentials (cf. (10)), of the invertibility of $(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})$ in $C^{1,\alpha}(\partial\Omega^o)$, of the invertibility of $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i})$ and J in $L^2(\partial\Omega^i)$ (cf. Lemmas 3.4 and 4.2), and of condition (2). \square

In Proposition 4.6 below we prove the existence of a solution $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta})$ in $C^{1,\alpha}(\partial\Omega^i) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ of the system of equations in (16). To do so we exploit the Leray-Schauder principle which is stated in the following Theorem 4.4 and which follows by the invariance of the Leray-Schauder topological degree (for a proof see, *e.g.*, Gilbarg and Trudinger [20, Theorem 11.3]).

Theorem 4.4 (Leray-Schauder principle). *Let \mathcal{X} be a Banach space. Let T be a continuous (nonlinear) operator from \mathcal{X} to itself which maps bounded sets to sets with a compact closure. If there exists a constant $M \in]0, +\infty[$ such that $\|x\|_{\mathcal{X}} \leq M$ for all $(x, t) \in \mathcal{X} \times [0, 1]$ satisfying $x = tT(x)$, then T has at least one fixed point $x \in \mathcal{X}$ such that $\|x\|_{\mathcal{X}} \leq M$.*

In order to apply this principle, we introduce in the following Lemma an elementary consequence of conditions (2) and (3).

Lemma 4.5. *If (2) and (3) hold true, then there exist $C_1, C_2, C_3, C_4 \in]0, +\infty[$ such that*

$$\|(I_{\Omega^i} + \mathcal{F}_F)^{(-1)}f\|_{C^0(\partial\Omega^i)} \leq C_1(C_2 + \|f\|_{C^0(\partial\Omega^i)})^{1/\delta_1} \quad (17)$$

and

$$\|\mathcal{F}_G \circ (I_{\Omega^i} + \mathcal{F}_F)^{(-1)}f\|_{C^0(\partial\Omega^i)} \leq C_3(C_4 + \|f\|_{C^0(\partial\Omega^i)})^{\delta_2} \quad (18)$$

for all functions $f \in C^0(\partial\Omega^i)$.

Proof. To prove (17) we observe that the first inequality in (3) implies that there exist $c_1^*, c_2^* \in]0, +\infty[$ such that $|t + F(x, t)| \geq c_1^*|t|^{\delta_1} - c_2^*$ for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$. Thus we have $\|(I_{\Omega^i} + \mathcal{F}_F)g\|_{C^0(\partial\Omega^i)} \geq c_1^*\|g\|_{C^0(\partial\Omega^i)}^{\delta_1} - c_2^*$ for all $g \in C^0(\partial\Omega^i)$ and the validity of (17) follows by taking $g = (I_{\Omega^i} + \mathcal{F}_F)^{(-1)}f$. To prove (18) we observe that the second inequality in (3) implies that there exist $c_3^*, c_4^* \in]0, +\infty[$ such that $|G(x, t)| \leq c_3^*(c_4^* + |t + F(x, t)|)^{\delta_2}$ for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$. Then we have $\|\mathcal{F}_G g\|_{C^0(\partial\Omega^i)} \leq c_3^*(c_4^* + \|(I_{\Omega^i} + \mathcal{F}_F)g\|_{C^0(\partial\Omega^i)})^{\delta_2}$ for all $g \in C^0(\partial\Omega^i)$ and the validity of (18) follows by condition (2) and by taking $g = (I_{\Omega^i} + \mathcal{F}_F)^{(-1)}f$. \square

Then we have the following.

Proposition 4.6. *Let conditions (2) and (3) hold. Then the nonlinear system (16) has at least one solution $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta})$ in $C^{1,\alpha}(\partial\Omega^o) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$.*

Proof. We plan to apply the Leray-Schauder Theorem 4.4 with \mathcal{X} equal to $C^{1,\alpha}(\partial\Omega^o) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ and $T \equiv (T^o, T_1, T_2)$ given by

$$\begin{aligned} T^o(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) &\equiv \left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)^{(-1)}(f^o - w_{\Omega^i}^-[\tilde{\mu}]_{|\partial\Omega^o} - v_{\Omega^i}[\tilde{\eta}]_{|\partial\Omega^o}) && \text{on } \partial\Omega^o, \\ T_1(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) &\equiv \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}\right)^{(-1)} \left[(I_{\Omega^i} + \mathcal{F}_F)^{(-1)} \left(w_{\Omega^o}^+[\tilde{\mu}^o]_{|\partial\Omega^i} + v_{\Omega^i}[\tilde{\eta}]_{|\partial\Omega^i} + 2W_{\Omega^i}\tilde{\mu} \right) \right] && \text{on } \partial\Omega^i, \\ T_2(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) &\equiv J^{(-1)} \left[\mathcal{F}_G \circ (I_{\Omega^i} + \mathcal{F}_F)^{(-1)} \left(w_{\Omega^o}^+[\tilde{\mu}^o]_{|\partial\Omega^i} + v_{\Omega^i}[\tilde{\eta}]_{|\partial\Omega^i} + 2W_{\Omega^i}\tilde{\mu} \right) \right. \\ &\quad \left. - v_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)^{(-1)}(f^o - w_{\Omega^i}^-[\tilde{\mu}]_{|\partial\Omega^o}) \right]_{|\partial\Omega^i} \right] && \text{on } \partial\Omega^i, \end{aligned} \quad (19)$$

for all $(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta}) \in C^{1,\alpha}(\partial\Omega^\circ) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$. We first verify that T is continuous from $C^{1,\alpha}(\partial\Omega^\circ) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ to itself and maps bounded sets to sets with compact closure. To do so, we consider separately T° , T_1 and T_2 . By Lemmas 3.2 and 3.4 one deduces that $\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ}$ is an isomorphism from $C^{1,\alpha}(\partial\Omega^\circ)$ to itself. In particular, $(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}$ is continuous from $C^{1,\alpha}(\partial\Omega^\circ)$ to itself. Moreover, by the properties of integral operators with real analytic kernel and no singularities $w_{\Omega^i}^-[\cdot]_{|\partial\Omega^\circ}$ and $v_{\Omega^i}[\cdot]_{|\partial\Omega^\circ}$ are compact from $C^0(\partial\Omega^i)$ to $C^{1,\alpha}(\partial\Omega^\circ)$. It follows that T° is continuous from $C^{1,\alpha}(\partial\Omega^\circ) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ to $C^{1,\alpha}(\partial\Omega^\circ)$ and maps bounded sets to sets with compact closure. We now consider T_1 . By Lemmas 3.2 and 3.4 one verifies that $\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}$ is an isomorphism from $C^0(\partial\Omega^i)$ to itself and thus $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i})^{(-1)}$ is continuous from $C^0(\partial\Omega^i)$ to itself. By assumption (2) the map $(I_{\Omega^i} + \mathcal{F}_F)^{(-1)}$ is continuous from $C^0(\partial\Omega^i)$ to itself. Then, by the properties of integral operators with real analytic kernel and no singularities $w_{\Omega^\circ}^+[\cdot]_{|\partial\Omega^i}$ is compact from $C^{1,\alpha}(\partial\Omega^\circ)$ to $C^0(\partial\Omega^i)$. By the mapping properties of the single layer potential (cf., *e.g.*, Kress [26, Thm. 2.22], see also Miranda [33, Chap. II, §14, III]), $v_{\Omega^i}[\cdot]_{|\partial\Omega^i}$ is compact from $C^0(\partial\Omega^i)$ to itself. By Lemma 3.1, W_{Ω^i} is compact from $C^0(\partial\Omega^i)$ to itself. It follows that T_1 is continuous from $C^{1,\alpha}(\partial\Omega^\circ) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ to $C^0(\partial\Omega^i)$ and maps bounded sets to sets with compact closure. Finally we consider T_2 . By Lemma 4.2 the operator $J^{(-1)}$ is continuous from $C^0(\partial\Omega^i)$ to itself. By the continuity of G and by condition (2), the map $\mathcal{F}_G \circ (I_{\Omega^i} + \mathcal{F}_F)^{(-1)}$ is continuous from $C^0(\partial\Omega^i)$ to itself. By the mapping properties of the single layer potential (cf., *e.g.*, Kress [26, Thm. 2.22], see also Miranda [33, Chap. II, §14, III]), $v_{\Omega^i}[\cdot]_{|\partial\Omega^i}$ is compact from $C^0(\partial\Omega^i)$ to itself. By Lemma 3.1, W_{Ω^i} is compact from $C^0(\partial\Omega^i)$ to itself. By the properties of integral operators with real analytic kernel and no singularities and by the continuity of $(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}$ from $C^{1,\alpha}(\partial\Omega^\circ)$ to itself, the map $w_{\Omega^\circ}^+[\cdot]_{|\partial\Omega^i}$ is compact from $C^{1,\alpha}(\partial\Omega^\circ)$ to $C^0(\partial\Omega^i)$ and the map $\nu_{\Omega^i} \cdot \nabla w_{\Omega^\circ}^+[(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}w_{\Omega^i}^-[\cdot]_{|\partial\Omega^\circ}]_{|\partial\Omega^i}$ is compact from $C^0(\partial\Omega^i)$ to itself. Accordingly T_2 is continuous from $C^{1,\alpha}(\partial\Omega^\circ) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ to $C^0(\partial\Omega^i)$ and maps bounded sets to sets with compact closure.

Now let $t \in [0, 1]$ and assume that $(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta}) = tT(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta})$. We show that there exists a constant $M \in]0, +\infty[$ (which does not depend on t) such that

$$\|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} + \|\tilde{\eta}\|_{C^0(\partial\Omega^i)} \leq M. \quad (20)$$

By equality $(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta}) = tT(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta})$ we have that

$$\begin{aligned} \|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} &\leq \|T^\circ(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta})\|_{C^{1,\alpha}(\partial\Omega^\circ)}, \\ \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} &\leq \|T_1(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta})\|_{C^0(\partial\Omega^i)}, \\ \|\tilde{\eta}\|_{C^0(\partial\Omega^i)} &\leq \|T_2(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta})\|_{C^0(\partial\Omega^i)}. \end{aligned} \quad (21)$$

By the first inequality of (21) we deduce that there exists a constant $m_1 \in]0, +\infty[$ which depends only on the norm of the bounded linear operator $(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}$ from $C^{1,\alpha}(\partial\Omega^\circ)$ to itself, on $\|f^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)}$, and on the norm of the linear bounded operators $w_{\Omega^\circ}^-[\cdot]_{|\partial\Omega^\circ}$ and $\nu_{\Omega^i}[\cdot]_{|\partial\Omega^\circ}$ from $C^0(\partial\Omega^i)$ to $C^{1,\alpha}(\partial\Omega^\circ)$, such that

$$\|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} \leq m_1(1 + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} + \|\tilde{\eta}\|_{C^0(\partial\Omega^i)}). \quad (22)$$

By the second inequality of (21) we deduce that there exist real constants $m_2, m_3 \in]0, +\infty[$ which depend on the norm of the linear bounded operator $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i})^{(-1)}$ from $C^0(\partial\Omega^i)$ to itself, on the constants C_1 and C_2 of Lemma 4.5, on the norm of the linear bounded operator $w_{\Omega^\circ}^+[\cdot]_{|\partial\Omega^i}$ from $C^{1,\alpha}(\partial\Omega^\circ)$ to $C^0(\partial\Omega^i)$, and on the norm of the linear bounded operators $\nu_{\Omega^i}[\cdot]_{|\partial\Omega^i}$ and W_{Ω^i} from $C^0(\partial\Omega^i)$ to itself such that

$$\|\tilde{\mu}\|_{C^0(\partial\Omega^i)} \leq m_2(m_3 + \|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} + \|\tilde{\eta}\|_{C^0(\partial\Omega^i)})^{1/\delta_1}. \quad (23)$$

By the third inequality of (21) we deduce that there exist real constants $m_4, m_5 \in]0, +\infty[$ which depend on the norm of the linear bounded operator $J^{(-1)}$ from $C^0(\partial\Omega^i)$ to itself, on the constants C_3 and C_4 of Lemma 4.5, on the norm of the linear bounded operator $w_{\Omega^\circ}^+[\cdot]_{|\partial\Omega^i}$ from $C^{1,\alpha}(\partial\Omega^\circ)$ to $C^0(\partial\Omega^i)$, on the norm of the linear bounded operators $\nu_{\Omega^i}[\cdot]_{|\partial\Omega^i}$ and W_{Ω^i} from $C^0(\partial\Omega^i)$ to itself, on the norm of $\nu_{\Omega^i} \cdot \nabla w_{\Omega^\circ}^+[(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}f^\circ]_{|\partial\Omega^i}$ in $C^0(\partial\Omega^i)$, and on the norm of the bounded linear operator $\nu_{\Omega^i} \cdot \nabla w_{\Omega^\circ}^+[(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}w_{\Omega^i}^-[\cdot]_{|\partial\Omega^\circ}]_{|\partial\Omega^i}$ from $C^0(\partial\Omega^i)$ to itself, such that

$$\begin{aligned} &\|\tilde{\eta}\|_{C^0(\partial\Omega^i)} \\ &\leq m_4 [(m_5 + \|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} + \|\tilde{\eta}\|_{C^0(\partial\Omega^i)})^{\delta_2} + 1 + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)}]. \end{aligned} \quad (24)$$

Then, by inequalities (22), (23), and (24) one deduces that there exists real constants $M_1, M_2, M_3 \in]0, +\infty[$, which depend on m_1, \dots, m_5 , such that

$$\begin{aligned} &\|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} + \|\tilde{\eta}\|_{C^0(\partial\Omega^i)} \\ &\leq M_1 + M_2(M_3 + \|\tilde{\mu}^\circ\|_{C^{1,\alpha}(\partial\Omega^\circ)} + \|\tilde{\mu}\|_{C^0(\partial\Omega^i)} + \|\tilde{\eta}\|_{C^0(\partial\Omega^i)})^{\delta^*} \end{aligned}$$

with $\delta_* \equiv \max\{1/\delta_1, \delta_2\} \in]0, 1[$. Then a straightforward calculation shows that inequality (20) holds with $M \equiv \max\{1, (M_1 + M_2(M_3 + 1)^{\delta_*})^{1/(1-\delta_*)}\}$. Now the validity of the statement follows by Theorem 4.4. \square

With a further regularity request on F and G we can find a solution of (16) in $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$.

Proposition 4.7. *Let conditions (2), (3), and (6) hold. Then the nonlinear system (16) has at least one solution $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta})$ in $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$.*

Proof. Let T be as in (19). By Proposition 4.5 there exists $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta})$ in $C^{1,\alpha}(\partial\Omega^o) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ such that $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) = T(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta})$. Then, by the mapping properties of integral operators with real analytic kernels and no singularities we have that $w_{\Omega^i}^-[\tilde{\mu}]|_{\partial\Omega^o}$ and $v_{\Omega^i}[\tilde{\eta}]|_{\partial\Omega^o}$ belong to $C^{1,\alpha}(\partial\Omega^o)$, that $w_{\Omega^o}^+[\tilde{\mu}^o]|_{\partial\Omega^i}$ belongs to $C^{1,\alpha}(\partial\Omega^i)$, and that $\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\psi]|_{\partial\Omega^i}$ belongs to $C^{0,\alpha}(\partial\Omega^i)$ for all $\psi \in C^0(\partial\Omega^o)$. By a classical result in potential theory (cf., e.g., Miranda [33, Chap. II, §14, III]) we have that $v_{\Omega^i}[\tilde{\eta}]|_{\partial\Omega^i} \in C^{0,\alpha}(\partial\Omega^i)$ and by Lemma 3.1 we have that $W_{\Omega^i}[\mu] \in C^{0,\alpha}(\partial\Omega^i)$. Then, by the invertibility of $\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}$ in $C^{1,\alpha}(\partial\Omega^o)$ and of $\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}$ in $C^{0,\alpha}(\partial\Omega^i)$ (cf. Lemma 3.1 and 3.4), by the invertibility of J in $C^{0,\alpha}(\partial\Omega^i)$ (cf. Lemma 4.2), and by assumption (6) it follows that $T(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$. Thus $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ and our proof is complete. \square

In the following Theorem 4.11 we show that under conditions (2) and (3) there exists a pair of functions $(\tilde{u}^o, \tilde{u}^i) \in C^0(\text{cl}\Omega^o \setminus \Omega^i) \times C^0(\text{cl}\Omega^i)$ which satisfy the first four conditions of problem (1) in the classical sense and which satisfies the fifth condition of (1) in a certain weak sense which we specify below. To do so, we introduce in Definition 4.8 a distribution $[\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w$ which plays the role of a weak counterpart of the difference of the normal derivatives of \tilde{w}^o and \tilde{w}^i and which is defined for all pair of continuous functions $(\tilde{w}^o, \tilde{w}^i) \in C_{\text{harm}}^0(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^0(\text{cl}\Omega^i)$ (see also definition (12)). We observe here that weak counterparts of the normal derivative have being largely considered in literature for functions in suitable Sobolev and Bessel spaces (see for example Costabel [6], Mikhailov [31], and the references therein).

Definition 4.8. *Let $(\tilde{w}^o, \tilde{w}^i)$ be a pair of functions of $C_{\text{harm}}^0(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^0(\text{cl}\Omega^i)$. Then $[\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w$ denotes the distribution on Ω^o*

defined by

$$\begin{aligned} & \langle [\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w, \phi \rangle \\ & \equiv \int_{\partial\Omega^i} (\tilde{w}^o_{|\partial\Omega^i} - \tilde{w}^i_{|\partial\Omega^i}) (\nu_{\Omega^i} \cdot \nabla \phi|_{\partial\Omega^i}) d\sigma + \int_{\Omega^o \setminus \Omega^i} \tilde{w}^o \Delta \phi dx + \int_{\Omega^i} \tilde{w}^i \Delta \phi dx \end{aligned}$$

for all test functions $\phi \in C_c^\infty(\Omega^o)$.

One immediately verifies that the map which takes $(\tilde{w}^o, \tilde{w}^i)$ to $[\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w$ is continuous. Namely we have the following.

Lemma 4.9. *Let $(\tilde{w}^o, \tilde{w}^i) \in C_{\text{harm}}^0(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^0(\text{cl}\Omega^i)$. Let $\{(\tilde{w}_j^o, \tilde{w}_j^i)\}_{j \in \mathbb{N}}$ be a sequence in $C_{\text{harm}}^0(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^0(\text{cl}\Omega^i)$ such that $\lim_{j \rightarrow +\infty} \tilde{w}_j^o = \tilde{w}^o$ in $C^0(\text{cl}\Omega^o \setminus \Omega^i)$ and $\lim_{j \rightarrow +\infty} \tilde{w}_j^i = \tilde{w}^i$ in $C^0(\text{cl}\Omega^i)$. Then*

$$\lim_{j \rightarrow +\infty} \langle [\nu_{\Omega^i} \cdot \nabla \tilde{w}_j^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}_j^i]_w, \phi \rangle = \langle [\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w, \phi \rangle \quad \forall \phi \in C_c^\infty(\Omega^o).$$

Moreover, if (w^o, w^i) belongs to $C_{\text{harm}}^1(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^1(\text{cl}\Omega^i)$, then $[\nu_{\Omega^i} \cdot \nabla w^o - \nu_{\Omega^i} \cdot \nabla w^i]_w$ coincides with $(\nu_{\Omega^i} \cdot \nabla w^o - \nu_{\Omega^i} \cdot \nabla w^i)|_{\partial\Omega^i}$. Namely we have

$$\langle [\nu_{\Omega^i} \cdot \nabla w^o - \nu_{\Omega^i} \cdot \nabla w^i]_w, \phi \rangle = \int_{\partial\Omega^i} (\nu_{\Omega^i} \cdot \nabla w^o(x) - \nu_{\Omega^i} \cdot \nabla w^i(x)) \phi(x) d\sigma_x$$

for all $\phi \in C_c^\infty(\Omega^o)$ and for all pair of functions $(w^o, w^i) \in C_{\text{harm}}^1(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^1(\text{cl}\Omega^i)$. Then we can prove that $[\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w$ is supported on $\partial\Omega^i$.

Lemma 4.10. *For all $(\tilde{w}^o, \tilde{w}^i) \in C_{\text{harm}}^0(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^0(\text{cl}\Omega^i)$ the support of $[\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w$ is contained in $\partial\Omega^i$.*

Proof. By a classical argument one can prove that there exists a sequence $\{(w_j^o, w_j^i)\}_{j \in \mathbb{N}}$ in $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ such that $\lim_{j \rightarrow +\infty} w_j^o = \tilde{w}^o$ in $C^0(\text{cl}\Omega^o \setminus \Omega^i)$ and $\lim_{j \rightarrow +\infty} w_j^i = \tilde{w}^i$ in $C^0(\text{cl}\Omega^i)$. Let $\phi_0 \in C_c^\infty(\Omega^o)$ be such that $\phi_0|_{\partial\Omega^i} = 0$. Then we have

$$\begin{aligned} & \langle [\nu_{\Omega^i} \cdot \nabla w_j^o - \nu_{\Omega^i} \cdot \nabla w_j^i]_w, \phi_0 \rangle \\ & = \int_{\partial\Omega^i} (\nu_{\Omega^i} \cdot \nabla w_j^o(x) - \nu_{\Omega^i} \cdot \nabla w_j^i(x)) \phi_0(x) d\sigma_x = 0 \end{aligned}$$

for all $j \in \mathbb{N}$. Moreover $\lim_{j \rightarrow \infty} \langle [\nu_{\Omega^i} \cdot \nabla w_j^o - \nu_{\Omega^i} \cdot \nabla w_j^i]_w, \phi_0 \rangle = \langle [\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w, \phi_0 \rangle$ by Lemma 4.9, and thus $\langle [\nu_{\Omega^i} \cdot \nabla \tilde{w}^o - \nu_{\Omega^i} \cdot \nabla \tilde{w}^i]_w, \phi_0 \rangle = 0$. \square

We are now ready to prove the main result of this section.

Theorem 4.11. *Assume that F and G satisfy (2) and (3). Then there exists $(\tilde{u}^\circ, \tilde{u}^i) \in C^0(\text{cl}\Omega^\circ \setminus \Omega^i) \times C^0(\text{cl}\Omega^i)$ such that*

$$\left\{ \begin{array}{ll} \Delta \tilde{u}^\circ = 0 & \text{in } \Omega^\circ \setminus \text{cl}\Omega^i, \\ \Delta \tilde{u}^i = 0 & \text{in } \Omega^i, \\ \tilde{u}^\circ(x) = f^\circ(x) & \text{for all } x \in \partial\Omega^\circ, \\ \tilde{u}^\circ(x) = F(x, \tilde{u}^i(x)) & \text{for all } x \in \partial\Omega^i, \\ \langle [\nu_{\Omega^i} \cdot \nabla \tilde{u}^\circ - \nu_{\Omega^i} \cdot \nabla \tilde{u}^i]_w, \phi \rangle & \\ = \int_{\partial\Omega^i} G(x, \tilde{u}^i(x)) \phi(x) d\sigma_x & \text{for all } \phi \in C_c^\infty(\Omega^\circ). \end{array} \right. \quad (25)$$

Proof. Let $(\tilde{\mu}^\circ, \tilde{\mu}, \tilde{\eta}) \in C^{1,\alpha}(\partial\Omega^\circ) \times C^0(\partial\Omega^i) \times C^0(\partial\Omega^i)$ be as in Proposition 4.6 and define

$$\tilde{u}^\circ \equiv (w_{\Omega^\circ}^+[\tilde{\mu}^\circ] + w_{\Omega^i}^-[\tilde{\mu}] + v_{\Omega^i}^-[\tilde{\eta}])|_{\text{cl}\Omega^\circ \setminus \Omega^i}, \quad \tilde{u}^i \equiv w_{\Omega^i}^+[\tilde{\mu}].$$

Then the pair $(\tilde{u}^\circ, \tilde{u}^i)$ belongs to $C^0(\text{cl}\Omega^\circ \setminus \Omega^i) \times C^0(\text{cl}\Omega^i)$ (cf. Folland [17, Chap. 3]) and satisfies the first four conditions of (25) (see also (10)). We now prove that $(\tilde{u}^\circ, \tilde{u}^i)$ satisfies also the fifth condition of (25).

By a standard argument one proves that there exists a sequence $\{v_j^i\}_{j \in \mathbb{N}}$ in $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ such that

$$\lim_{j \rightarrow +\infty} v_j^i = \tilde{u}^i \quad \text{in } C^0(\text{cl}\Omega^i). \quad (26)$$

By Lemmas 3.2 and 3.4 we have that $\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}$ is an isomorphism from $C^0(\partial\Omega^i)$ to itself and from $C^{1,\alpha}(\partial\Omega^i)$ to itself. Then, by (10) one verifies that there exists $\mu_j \in C^{1,\alpha}(\partial\Omega^i)$ such that $v_j^i = w_{\Omega^i}^+[\mu_j]$ for all $j \in \mathbb{N}$. Moreover, by the continuity of $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i})^{(-1)}$ from $C^0(\partial\Omega^i)$ to itself, we have

$$\begin{aligned} \lim_{j \rightarrow +\infty} \mu_j &= \lim_{j \rightarrow +\infty} (\frac{1}{2}I_{\Omega^i} + W_{\Omega^i})^{(-1)} v_j^i|_{\partial\Omega^i} \\ &= (\frac{1}{2}I_{\Omega^i} + W_{\Omega^i})^{(-1)} \tilde{u}^i|_{\partial\Omega^i} = \tilde{\mu} \quad \text{in } C^0(\partial\Omega^i). \end{aligned} \quad (27)$$

Then we set

$$\mu_j^\circ \equiv (\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)} (f^\circ - w_{\Omega^i}^-[\mu_j]|_{\partial\Omega^\circ} - v_{\Omega^i}^-[\tilde{\eta}]|_{\partial\Omega^\circ}) \quad \forall j \in \mathbb{N}.$$

By Lemmas 3.2 and 3.4, we have that $\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ}$ is an isomorphism from $C^{1,\alpha}(\partial\Omega^\circ)$ to itself and from $C^0(\partial\Omega^\circ)$ to itself. In particular, $(\frac{1}{2}I_{\Omega^\circ} + W_{\Omega^\circ})^{(-1)}$

is continuous from $C^0(\partial\Omega^o)$ to itself and maps $C^{1,\alpha}(\partial\Omega^o)$ to itself. Moreover, by the properties of integral operators with real analytic kernel and no singularities $w_{\Omega^i}^-[\cdot]_{|\partial\Omega^o}$ is continuous from $C^0(\partial\Omega^i)$ to $C^{1,\alpha}(\partial\Omega^o)$ and $f^o - v_{\Omega^i}[\tilde{\eta}]_{|\partial\Omega^o}$ belongs to $C^{1,\alpha}(\partial\Omega^o)$. It follows that $\mu_j^o \in C^{1,\alpha}(\partial\Omega^o)$ for all $j \in \mathbb{N}$ and that

$$\begin{aligned} \lim_{j \rightarrow +\infty} \mu_j^o &= \lim_{j \rightarrow +\infty} \left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} (f^o - w_{\Omega^i}^-[\mu_j]_{|\partial\Omega^o} - v_{\Omega^i}[\tilde{\eta}]_{|\partial\Omega^o}) \\ &= \left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} (f^o - w_{\Omega^i}^-[\tilde{\mu}]_{|\partial\Omega^o} - v_{\Omega^i}[\tilde{\eta}]_{|\partial\Omega^o}) = \tilde{\mu}^o \quad \text{in } C^0(\partial\Omega^o). \end{aligned} \tag{28}$$

Now let

$$v_j^o \equiv (w_{\Omega^o}^+[\mu_j^o] + w_{\Omega^i}^-[\mu_j] + v_{\Omega^i}^-[\tilde{\eta}])_{|\text{cl}\Omega^o \setminus \Omega^i} \quad \forall j \in \mathbb{N}.$$

By classical potential theory $v_j^o \in C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i)$ (cf., *e.g.*, Miranda [32]). Moreover, by (10) we have

$$v_j^o|_{\partial\Omega^o} = \left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right) \mu_j^o + w_{\Omega^i}^-[\mu_j]_{|\partial\Omega^o} + v_{\Omega^i}^-[\tilde{\eta}]_{|\partial\Omega^o}$$

and

$$v_j^o|_{\partial\Omega^i} = w_{\Omega^o}^+[\mu_j^o]_{|\partial\Omega^i} + \left(-\frac{1}{2} I_{\Omega^i} + W_{\Omega^i} \right) \mu_j + v_{\Omega^i}^-[\tilde{\eta}]_{|\partial\Omega^i}.$$

Then, by (27) and (28), by the continuity of $\frac{1}{2} I_{\Omega^o} + W_{\Omega^o}$ from $C^0(\partial\Omega^o)$ to itself and of $-\frac{1}{2} I_{\Omega^o} + W_{\Omega^o}$ from $C^0(\partial\Omega^i)$ to itself (cf. Lemma 3.1), and by the properties of integral operators with real analytic kernels and no singularity, we deduce that

$$\lim_{j \rightarrow +\infty} v_j^o|_{\partial\Omega^o} = \tilde{u}|_{\partial\Omega^o} \quad \text{in } C^0(\partial\Omega^o)$$

and

$$\lim_{j \rightarrow +\infty} v_j^o|_{\partial\Omega^i} = \tilde{u}|_{\partial\Omega^i} \quad \text{in } C^0(\partial\Omega^i).$$

It follows that

$$\lim_{j \rightarrow +\infty} v_j^o = \tilde{u} \quad \text{in } C^0(\text{cl}\Omega^o \setminus \Omega^i). \tag{29}$$

In addition, by the jump formulas (10) and (11) and by the validity of equality (16) for $(\mu^o, \mu, \eta) = (\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta})$ (cf. Proposition 4.6) one verifies that the pair (v_j^o, v_j^i) satisfies the equality

$$\nu_{\Omega^i} \cdot \nabla v_j^o(x) - \nu_{\Omega^i} \cdot \nabla v_j^i(x) = G(x, \tilde{u}^i(x)) + \nu_{\Omega^i}(x) \cdot \nabla w_{\Omega^o}^+[\mu_j^o - \tilde{\mu}^o](x) \quad \forall x \in \partial\Omega^i$$

for all $j \in \mathbb{N}$. Hence, by the continuity of the map from $C^0(\partial\Omega^o)$ to $C^0(\partial\Omega^i)$ which takes ϕ to $\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\phi]|_{\partial\Omega^i}$ and by the limit relation in (28) we have that

$$\lim_{j \rightarrow +\infty} (\nu_{\Omega^i} \cdot \nabla v_j^o - \nu_{\Omega^i} \cdot \nabla v_j^i)|_{\partial\Omega^i} = \mathcal{F}_G \tilde{u}^i \quad \text{in } C^0(\partial\Omega^i).$$

Thus, by Lemma 4.9, by the limit relations in (26) and (29), and by the membership of (v_j^o, v_j^i) in $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ for all $j \in \mathbb{N}$, it follows that $(\tilde{u}^o, \tilde{u}^i)$ satisfies the fifth condition in problem (25). The theorem is now proved. \square

If in addition F and G satisfy assumption (6), then the pair $(\tilde{u}^o, \tilde{u}^i)$ belongs to $C^{0,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{0,\alpha}(\text{cl}\Omega^i)$.

Theorem 4.12. *Assume that F and G satisfy (2), (3), and (6). Then there exists $(\tilde{u}^o, \tilde{u}^i) \in C^{0,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{0,\alpha}(\text{cl}\Omega^i)$ which satisfy the conditions in (25).*

Proof. If $(\tilde{\mu}^o, \tilde{\mu}, \tilde{\eta}) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ is as in Proposition 4.12 and

$$\tilde{u}^o \equiv (w_{\Omega^o}^+[\tilde{\mu}^o] + w_{\Omega^i}^-[\tilde{\mu}] + v_{\Omega^i}^-[\tilde{\eta}]|_{\text{cl}\Omega^o \setminus \Omega^i}), \quad \tilde{u}^i \equiv w_{\Omega^i}^+[\tilde{\mu}],$$

then the pair $(\tilde{u}^o, \tilde{u}^i)$ belongs to $C^{0,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{0,\alpha}(\text{cl}\Omega^i)$ (cf. Miranda [32]) and we can prove that it satisfies the conditions of (25) by arguing as in the proof of Theorem 4.11. \square

5. Existence result for problem (7)

We now fix a real number $\lambda > 0$ and a continuous function Φ from $\partial\Omega^i \times \mathbb{R}$ to \mathbb{R} . Then we assume that $F = \lambda \text{id}_{\mathbb{R}} + \epsilon\Phi$, where ϵ is a multiplicative real parameter. Our aim is to study the nonlinear transmission problem (7) for ϵ small. To do so, we find convenient to introduce the following technical assumption:

$$\begin{aligned} & \text{the map from } C^{0,\alpha}(\partial\Omega^i) \text{ to } C^{1,\alpha}(\partial\Omega^i) \text{ which} \\ & \text{takes } \eta \text{ to } v_{\Omega^i}[\eta]|_{\partial\Omega^i} \text{ is an isomorphism.} \end{aligned} \tag{30}$$

We observe that assumption (30) holds for all domains Ω^i in \mathbb{R}^n if $n \geq 3$, and does not hold in \mathbb{R}^2 only in exceptional cases. Indeed we have the following classical result.

Proposition 5.1. *Let Ω be an open bounded connected subset of \mathbb{R}^2 of class $C^{1,\alpha}$. Then there exists a unique function ψ_Ω in $C^{0,\alpha}(\partial\Omega)$ such that*

$$v_\Omega[\psi_\Omega]|_{\partial\Omega} \text{ is constant and } \int_{\partial\Omega} \psi_\Omega d\sigma = 1.$$

Moreover, the following statements hold.

- (i) *If $v_\Omega[\psi_\Omega]|_{\partial\Omega} \neq 0$, then $v_\Omega[\cdot]|_{\partial\Omega}$ is an isomorphism from $C^{0,\alpha}(\partial\Omega)$ to $C^{1,\alpha}(\partial\Omega)$.*
- (ii) *If $v_\Omega[\psi_\Omega]|_{\partial\Omega} = 0$ and $r \in]0, +\infty[\setminus \{1\}$, then $v_{r\Omega}[\cdot]|_{r\partial\Omega}$ is an isomorphism from $C^{0,\alpha}(r\partial\Omega)$ to $C^{1,\alpha}(r\partial\Omega)$.*

In Lemma 5.2 below we introduce an isomorphism between $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ and $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ (cf. definition (12)).

Lemma 5.2. *Let $U \equiv (U^o, U^i)$ be the operator from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ to $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ which takes (μ^o, η^o, η^i) to the pair $(U^o[\mu^o, \eta^o, \eta^i], U^i[\mu^o, \eta^o, \eta^i])$ given by*

$$\begin{aligned} U^o[\mu^o, \eta^o, \eta^i] &\equiv (w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta^o])|_{\text{cl}\Omega^o \setminus \Omega^i}, \\ U^i[\mu^o, \eta^o, \eta^i] &\equiv \lambda^{-1}w_{\Omega^o}^+[\mu^o]|_{\text{cl}\Omega^i} + \lambda^{-1}v_{\Omega^i}^+[\eta^i]. \end{aligned}$$

Then U is a linear isomorphism.

Proof. By the mapping properties of the single and double layer potentials one verifies that the operator U is continuous from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ to $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ (cf. Section 3, see also Miranda [32]).

Therefore, if we prove that U is one-to-one and onto, we can deduce by the open mapping theorem that U is an isomorphism from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ to $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$. So let $(\phi^o, \phi^i) \in C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$. We show that there exists unique triple $(\mu^o, \eta^o, \eta^i) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ such that

$$(U^o[\mu^o, \eta^o, \eta^i], U^i[\mu^o, \eta^o, \eta^i]) = (\phi^o, \phi^i).$$

We first consider $U^o[\mu^o, \eta^o, \eta^i] = \phi^o$ and we verify that there exists unique (μ^o, η^o) such that

$$(w_{\Omega^o}^+[\mu^o] + v_{\Omega^i}^-[\eta^o])|_{\text{cl}\Omega^o \setminus \Omega^i} = \phi^o. \quad (31)$$

By the uniqueness of the classical solution of the Neumann-Dirichlet mixed problem and by the jump properties of the single and double layer potentials (cf. equality (10)), equation (31) is equivalent to

$$\begin{aligned} \left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)\mu^o + v_{\Omega^i}^-[\eta^o]_{|\partial\Omega^o} &= \phi^o_{|\partial\Omega^o}, \\ \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*\right)\eta^o + \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} &= \nu_{\Omega^i} \cdot \nabla \phi^o_{|\partial\Omega^i}. \end{aligned}$$

By classical potential theory, the operator $(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})$ is an isomorphism from $C^{1,\alpha}(\partial\Omega^o)$ to itself and the operator $(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)$ is an isomorphism from $C^{0,\alpha}(\partial\Omega^i)$ to itself (cf. Lemmas 3.2 and 3.4). Then, by the compactness properties of the integral operators with real analytic kernel and with no singularities and by standard properties of Fredholm operators, we deduce that the map which takes (μ, η) to $((\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})\mu + v_{\Omega^i}^-[\eta]_{|\partial\Omega^o}, (\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*)\eta + \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu]_{|\partial\Omega^i})$ is a Fredholm operator of index 0 from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ to itself. Thus, to prove the existence and uniqueness of $(\mu^o, \eta^o) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ which satisfies (31) it suffices to show that $(\mu, \eta) = (0, 0)$ when

$$\left(\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)\mu + v_{\Omega^i}^-[\eta]_{|\partial\Omega^o}, \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*\right)\eta + \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu]_{|\partial\Omega^i} \right) = (0, 0). \quad (32)$$

If (32) holds, then $(w_{\Omega^o}^+[\mu] + v_{\Omega^i}^-[\eta])_{|\text{cl}\Omega^o \setminus \Omega^i} = 0$ by the uniqueness of the classical solution of the Neumann-Dirichlet mixed problem and by (10). Hence $w_{\Omega^o}^+[\mu] + v_{\Omega^i}^-[\eta] = 0$ in $\text{cl}\Omega^i$ by the uniqueness of the classical solution of the Dirichlet problem in Ω^i and by the continuity of $(w_{\Omega^o}^+[\mu] + v_{\Omega^i}^-[\eta])_{|\text{cl}\Omega^o}$ (cf. Section 3). Then, by the jump properties of the single layer potential (cf. equality (10)) we have that

$$\begin{aligned} \eta &= \nu_{\Omega^i} \cdot \nabla v_{\Omega^i}^-[\eta]_{|\partial\Omega^i} - \nu_{\Omega^i} \cdot \nabla v_{\Omega^i}^+[\eta]_{|\partial\Omega^i} \\ &= \nu_{\Omega^i} \cdot \nabla (w_{\Omega^o}^+[\mu] + v_{\Omega^i}^-[\eta])_{|\partial\Omega^i} - \nu_{\Omega^i} \cdot \nabla (w_{\Omega^o}^+[\mu] + v_{\Omega^i}^+[\eta])_{|\partial\Omega^i} = 0 \end{aligned}$$

and thus $\mu = 0$ by the first equality in (32) and by Lemma 3.4 (iii). Now, to complete the proof we observe that $U^i[\mu^o, \eta^o, \eta^i] = \phi^i$ is equivalent to $v_{\Omega^i}^+[\eta^i] = \lambda\phi^i - w_{\Omega^o}^+[\mu^o]_{|\text{cl}\Omega^i}$ and the existence and uniqueness of η^i is guaranteed by the assumption in (30). \square

In the following Lemma 5.3 we introduce an auxiliary operator which we denote by J_λ .

Lemma 5.3. *Let*

$$J_\lambda[\eta] \equiv \left(\frac{1}{2}I_{\Omega^i} + \frac{\lambda-1}{\lambda+1}W_{\Omega^i}^* \right) \eta - \frac{\lambda-1}{\lambda+1}\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} \nu_{\Omega^i}[\eta]_{|\partial\Omega^o} \right]_{|\partial\Omega^i}$$

for all $\eta \in L^2(\partial\Omega^i)$. Then the map which takes η to $J_\lambda[\eta]$ is an isomorphism from $L^2(\partial\Omega^i)$ to itself, from $C^0(\partial\Omega^i)$ to itself, and from $C^{0,\alpha}(\partial\Omega^i)$ to itself.

Proof. By the properties of integral operators with real analytic kernels and no singularity, by the invertibility of $\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}$ in $C^{1,\alpha}(\partial\Omega^o)$ (cf. Lemma 3.2 and Lemma 3.4), and by the continuity of the map $w_{\Omega^o}^+[\cdot]$ from $C^{1,\alpha}(\partial\Omega^o)$ to $C^{1,\alpha}(\text{cl}\Omega^o)$ (cf., e.g., Miranda [32]), one deduces that the operator which takes η to

$$\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} \nu_{\Omega^i}[\eta]_{|\partial\Omega^o} \right]_{|\partial\Omega^i} \quad (33)$$

is continuous from $L^2(\partial\Omega^i)$ to $C^{0,\alpha}(\partial\Omega^i)$. Then, by the invertibility of $\frac{1}{2}I_{\Omega^i} + \frac{\lambda-1}{\lambda+1}W_{\Omega^i}^*$ in $L^2(\partial\Omega^i)$ (cf. Lemma 3.5) it follows that J_λ is a Fredholm operator of index 0 from $L^2(\partial\Omega^i)$ to itself. Thus, to show that J_λ is invertible from $L^2(\partial\Omega^i)$ to itself it suffices to prove that $J_\lambda[\eta] = 0$ implies $\eta = 0$. Now, if $\eta \in L^2(\partial\Omega^i)$ and $J_\lambda[\eta] = 0$, then $(\frac{1}{2}I_{\Omega^i} + \frac{\lambda-1}{\lambda+1}W_{\Omega^i}^*)\eta \in C^{0,\alpha}(\partial\Omega^i)$ by the membership of (33) in $C^{0,\alpha}(\partial\Omega^i)$, and thus $\eta \in C^{0,\alpha}(\partial\Omega^i)$ by Lemma 3.3. Then, by taking $\mu^o \equiv -(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})^{(-1)}\nu_{\Omega^i}[\eta]_{|\partial\Omega^o}$ and by a straightforward calculation based on (10) one verifies that $U^o[\mu^o, \eta, \eta]_{|\partial\Omega^o} = 0$, $U^o[\mu^o, \eta, \eta]_{|\partial\Omega^i} = \lambda U^i[\mu^o, \eta, \eta]_{|\partial\Omega^i}$, and $\nu_{\Omega^i} \cdot \nabla U^o[\mu^o, \eta, \eta]_{|\partial\Omega^i} = \nu_{\Omega^i} \cdot \nabla U^i[\mu^o, \eta, \eta]_{|\partial\Omega^i}$ (where $U^o[\mu^o, \eta, \eta]$ and $U^i[\mu^o, \eta, \eta]$ are defined as in Lemma 5.2). Then, by the uniqueness of the solution of the linear perfect contact problem we have $U^o[\mu^o, \eta, \eta] = 0$ and $U^i[\mu^o, \eta, \eta] = 0$. Accordingly, Lemma 5.2 implies that $\eta = 0$.

To prove that J_λ is invertible from $C^0(\partial\Omega^i)$ to itself, we first observe that J_λ is continuous from $C^0(\partial\Omega^i)$ to itself (see Lemma 3.5). Moreover, if $\eta \in L^2(\partial\Omega^i)$ and $J_\lambda[\eta] \in C^0(\partial\Omega^i)$ then $(\frac{1}{2}I_{\Omega^i} + \frac{\lambda-1}{\lambda+1}W_{\Omega^i}^*)\eta \in C^0(\partial\Omega^i)$ by the membership of (33) in $C^{0,\alpha}(\partial\Omega^i)$, and thus Lemma 3.3 ensures that $\eta \in C^0(\partial\Omega^i)$.

Similarly, to prove that J_λ is invertible from $C^{0,\alpha}(\partial\Omega^i)$ to itself we observe that J_λ is continuous from $C^{0,\alpha}(\partial\Omega^i)$ to itself and that $J_\lambda[\eta] \in C^{0,\alpha}(\partial\Omega^i)$ implies $\eta \in C^{0,\alpha}(\partial\Omega^i)$ for all $\eta \in L^2(\partial\Omega^i)$. \square

We now turn to consider problem (7) for $\epsilon = 0$. By the previous Lemma 5.2 and by the jump properties of the single and double layer potentials (cf. equality (10)) we deduce the following.

Lemma 5.4. *Let (μ^o, η) belong to $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$. Then the pair $(U^o[\mu^o, \eta, \eta], U^i[\mu^o, \eta, \eta])$ is a solution of (7) with $\epsilon = 0$ if and only if*

$$\begin{cases} (\frac{1}{2}I_{\Omega^o} + W_{\Omega^o})\mu^o = f^o - v_{\Omega^i}^-[\eta]_{|\partial\Omega^o}, \\ (\frac{1}{2}I_{\Omega^o} + \frac{\lambda-1}{\lambda+1}W_{\Omega^o}^*)\eta + \frac{\lambda-1}{\lambda+1}\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} \\ = \frac{\lambda}{\lambda+1}\mathcal{F}_G(\lambda^{-1}w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} + \lambda^{-1}v_{\Omega^i}^+[\eta]_{|\partial\Omega^i}). \end{cases} \quad (34)$$

We show the existence of a solution of (34) by an argument based on the invariance of the Leray-Schauder topological degree (cf. Theorem 4.4).

Proposition 5.5. *Assume that \mathcal{F}_G maps $C^{0,\alpha}(\partial\Omega^i)$ to itself and that there exist $C > 0$ and $\delta \in [0, 1[$ such that*

$$|G(x, t)| \leq C(1 + |t|)^\delta \quad \forall (x, t) \in \partial\Omega^i \times \mathbb{R}.$$

Then there exists at least a solution $(\mu_0^o, \eta_0) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ of (34).

Proof. Since $\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}$ is an invertible operator from $C^{1,\alpha}(\partial\Omega^o)$ to itself (cf. Lemmas 3.2 and 3.4) it is enough to show that there exists a solution $\eta_0 \in C^{0,\alpha}(\partial\Omega^i)$ of

$$\begin{aligned} \eta = J_\lambda^{(-1)} & \left[-\frac{\lambda-1}{\lambda+1}\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} f^o \right]_{|\partial\Omega^i} \right. \\ & \left. + \frac{\lambda}{\lambda+1}\mathcal{F}_G \left(\frac{1}{\lambda}w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} (f^o - v_{\Omega^i}^-[\eta]_{|\partial\Omega^o}) \right]_{|\partial\Omega^i} + \frac{1}{\lambda}v_{\Omega^i}^+[\eta]_{|\partial\Omega^i} \right) \right] \end{aligned} \quad (35)$$

(cf. Lemma 5.3). We first show that the equation (35) has a solution in $C^0(\partial\Omega^i)$. By the properties of integral operators with real analytic kernel and no singularities, by the invertibility of $\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}$ in $C^0(\partial\Omega^o)$ (cf. Lemmas 3.2 and 3.4), and by the mapping properties of the single layer potential (cf., e.g., Kress [26, Thm. 2.22], see also Miranda [33, Chap. II, §14, III]) we verify that the map from $C^0(\partial\Omega^i)$ to itself which takes a function η to

$$\frac{1}{\lambda}w_{\Omega^o}^+ \left[\left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} (f^o - v_{\Omega^i}^-[\eta]_{|\partial\Omega^o}) \right]_{|\partial\Omega^i} + \frac{1}{\lambda}v_{\Omega^i}^+[\eta]_{|\partial\Omega^i}$$

is compact (cf. Section 3). In addition, \mathcal{F}_G is continuous from $C^0(\partial\Omega^i)$ to itself (because G is continuous). It follows that the map which takes η to

$$\mathcal{F}_G \left(\frac{1}{\lambda} w_{\Omega^o}^+ \left[\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} (f^o - v_{\Omega^i}^- [\eta]_{|\partial\Omega^o}) \right]_{|\partial\Omega^i} + \frac{1}{\lambda} v_{\Omega^i}^+ [\eta]_{|\partial\Omega^i} \right)$$

is continuous from $C^0(\partial\Omega^i)$ to itself and maps bounded sets to sets with compact closure. Then, Lemma 5.3 implies that the map from $C^0(\partial\Omega^i)$ to itself which takes η to the right hand side of equation (35) is continuous and maps bounded sets to sets with compact closure. Now let $t \in [0, 1]$ and assume that

$$\begin{aligned} \eta = & t J_{\lambda}^{(-1)} \left[-\frac{\lambda-1}{\lambda+1} \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} f^o \right]_{|\partial\Omega^i} \right. \\ & \left. + \frac{\lambda}{\lambda+1} \mathcal{F}_G \left(\frac{1}{\lambda} w_{\Omega^o}^+ \left[\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} (f^o - v_{\Omega^i}^- [\eta]_{|\partial\Omega^o}) \right]_{|\partial\Omega^i} + \frac{1}{\lambda} v_{\Omega^i}^+ [\eta]_{|\partial\Omega^i} \right) \right] \end{aligned}$$

Then, by exploiting inequality $|G(x, t)| \leq C(1 + |t|)^\delta$ one verifies that

$$\|\eta\|_{C^0(\partial\Omega^i)} \leq c_1 + c_2 (c_3 + c_4 \|\eta\|_{C^0(\partial\Omega^i)})^\delta \quad (36)$$

where c_1, \dots, c_4 are real positive numbers which depend on C, t , and λ , on the norm of the bounded operator $J_{\lambda}^{(-1)}$ from $C^0(\partial\Omega^i)$ to itself, on the norm of the bounded operator from $C^0(\partial\Omega^i)$ to itself which takes ϕ to

$$-w_{\Omega^o}^+ \left[\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} v_{\Omega^i}^- [\phi]_{|\partial\Omega^o} \right]_{|\partial\Omega^i} + v_{\Omega^i}^+ [\phi]_{|\partial\Omega^i},$$

and on the $C^0(\partial\Omega^i)$ norms of the functions $w_{\Omega^o}^+ \left[\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} f^o \right]_{|\partial\Omega^i}$ and $\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ \left[\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o} \right)^{(-1)} f^o \right]_{|\partial\Omega^i}$. Then inequality (36) implies that

$$\|\eta\|_{C^0(\partial\Omega^i)} \leq \max \left\{ 1, (c_1 + c_2 (c_3 + c_4)^\delta)^{1/(1-\delta)} \right\}$$

Thus Theorem 4.4 implies that there exists $\eta_0 \in C^0(\partial\Omega^i)$ solution of (35). Then, by classical results of potential theory (cf. Miranda [33, Chap. II, §14, III]), we have $v_{\Omega^i}^+ [\eta_0]_{|\partial\Omega^i} \in C^{0,\alpha}(\partial\Omega^i)$, and, by the properties of integral operators with real analytic kernels and no singularities, by the assumption

that \mathcal{F}_G maps $C^{0,\alpha}(\partial\Omega^i)$ to itself, and by equation (35) we deduce that $\eta_0 \in C^{0,\alpha}(\partial\Omega^i)$. \square

We now pass to consider $\epsilon \neq 0$. We assume that

the composition operator \mathcal{F}_Φ is continuously Fréchet differentiable from $C^{1,\alpha}(\partial\Omega^i)$ to itself and the composition operator \mathcal{F}_G is continuously Fréchet differentiable from $C^{0,\alpha}(\partial\Omega^i)$ to itself. (37)

We observe that condition (37) implies that the partial derivatives $\partial_t\Phi(x, t)$ and $\partial_t G(x, t)$ exist for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$, that the composition operators $\mathcal{F}_{\partial_t\Phi}$ and $\mathcal{F}_{\partial_t G}$ map $C^{1,\alpha}(\partial\Omega^i)$ to itself and $C^{0,\alpha}(\partial\Omega^i)$ to itself, respectively, and that

$$\begin{aligned} d\mathcal{F}_\Phi(v_0).v &= (\mathcal{F}_{\partial_t\Phi} v_0) v \quad \forall v \in C^{1,\alpha}(\partial\Omega^i), \\ d\mathcal{F}_G(w_0).w &= (\mathcal{F}_{\partial_t G} w_0) w \quad \forall w \in C^{0,\alpha}(\partial\Omega^i), \end{aligned}$$

where $d\mathcal{F}_\Phi(v_0)$ denotes the differential of \mathcal{F}_Φ evaluated at a function $v_0 \in C^{1,\alpha}(\partial\Omega^i)$ and $d\mathcal{F}_G(w_0)$ denotes the differential of \mathcal{F}_G evaluated at a function $w_0 \in C^{0,\alpha}(\partial\Omega^i)$ (cf., e.g., Lanza de Cristoforis [27, Prop. 6.3]).

Now we introduce the nonlinear operator $N \equiv (N^o, N_1^i, N_2^i)$ from $\mathbb{R} \times C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)^2$ to $C^{1,\alpha}(\partial\Omega^o) \times C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$ which takes $(\epsilon, \mu^o, \eta^o, \eta^i)$ to

$$\begin{aligned} N^o[\epsilon, \mu^o, \eta^o, \eta^i] &\equiv \left(\frac{1}{2}I_{\Omega^o} + W_{\Omega^o}\right)\mu^o + v_{\Omega^i}^-[\eta^o]_{|\partial\Omega^o} - f^o, \\ N_1^i[\epsilon, \mu^o, \eta^o, \eta^i] &\equiv v_{\Omega^i}[\eta^o - \eta^i]_{|\partial\Omega^i} - \epsilon\mathcal{F}_\Phi(\lambda^{-1}w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} + \lambda^{-1}v_{\Omega^i}^+[\eta^i]_{|\partial\Omega^i}), \\ N_2^i[\epsilon, \mu^o, \eta^o, \eta^i] &\equiv \left(\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*\right)\eta^o - \lambda^{-1}\left(-\frac{1}{2}I_{\Omega^i} + W_{\Omega^i}^*\right)\eta^i \\ &\quad + (\lambda - 1)\lambda^{-1}\nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} - \mathcal{F}_G(\lambda^{-1}w_{\Omega^o}^+[\mu^o]_{|\partial\Omega^i} + \lambda^{-1}v_{\Omega^i}^+[\eta^i]_{|\partial\Omega^i}). \end{aligned}$$

Then, by the mapping and jump properties of single and double layer potentials (cf. Section 3), one verifies the validity of the following Lemmas 5.6 and 5.7.

Lemma 5.6. *If Φ and G satisfy condition (37), then N is continuously Fréchet differentiable map from $\mathbb{R} \times C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)^2$ to $C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$.*

Lemma 5.7. *Let $(\epsilon, \mu^o, \eta^o, \eta^i) \in \mathbb{R} \times C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)^2$. Then we have $N[\epsilon, \mu^o, \eta^o, \eta^i] = 0$ if and only if $(U^o[\mu^o, \eta^o, \eta^i], U^i[\mu^o, \eta^o, \eta^i])$ is a solution of problem (7).*

Moreover, one can prove the following.

Lemma 5.8. *Let Φ and G satisfy the condition in (37). Let $(\tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i) \in C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)^2$. If*

$$\partial_t G(x, \lambda^{-1} w_{\Omega^o}^+[\tilde{\mu}^o](x) + \lambda^{-1} v_{\Omega^i}^+[\tilde{\eta}^i](x)) \geq 0 \quad \forall x \in \partial\Omega^i, \quad (38)$$

then $\partial_{(\mu^o, \eta^o, \eta^i)} N[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i]$ (the partial differential of the map N with respect to (μ^o, η^o, η^i) evaluated at $(0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i)$) is an isomorphism from $C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)^2$ to $C^{1,\alpha}(\partial\Omega^o) \times C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)$.

Proof. We have

$$\begin{aligned} \partial_{(\mu^o, \eta^o, \eta^i)} N^o[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i](\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) &= \left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o}\right) \bar{\mu}^o + v_{\Omega^i}^- [\bar{\eta}^o]_{|\partial\Omega^o}, \\ \partial_{(\mu^o, \eta^o, \eta^i)} N_1^i[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i](\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) &= v_{\Omega^i} [\bar{\eta}^o - \bar{\eta}^i]_{|\partial\Omega^i}, \\ \partial_{(\mu^o, \eta^o, \eta^i)} N_2^i[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i](\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) &= \left(\frac{1}{2} I_{\Omega^i} + W_{\Omega^i}^*\right) \bar{\eta}^o - \lambda^{-1} \left(-\frac{1}{2} I_{\Omega^i} + W_{\Omega^i}^*\right) \bar{\eta}^i \\ &\quad + (\lambda - 1) \lambda^{-1} \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ [\bar{\mu}^o]_{|\partial\Omega^i} - \tilde{\gamma} (\lambda^{-1} w_{\Omega^o}^+ [\bar{\mu}^o]_{|\partial\Omega^i} + \lambda^{-1} v_{\Omega^i}^+ [\bar{\eta}^i]_{|\partial\Omega^i}), \end{aligned}$$

for all $(\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) \in C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)^2$. Here $\tilde{\gamma}$ denotes the function of $C^{0,\alpha}(\partial\Omega^i)$ defined by

$$\tilde{\gamma}(x) \equiv \partial_t G(x, w_{\Omega^o}^+[\tilde{\mu}^o](x) + v_{\Omega^i}^- [\tilde{\eta}^i](x)) \quad \forall x \in \partial\Omega^i.$$

The operator which takes $(\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i)$ to $\partial_{(\mu^o, \eta^o, \eta^i)} N[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i](\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i)$ is Fredholm of index 0 from $C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)^2$ to itself. Indeed the operator which takes $(\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i)$ to

$$\left(\left(\frac{1}{2} I_{\Omega^o} + W_{\Omega^o}\right) \bar{\mu}^o, v_{\Omega^i} [\bar{\eta}^o - \bar{\eta}^i]_{|\partial\Omega^i}, \frac{1}{2} (\bar{\eta}^o + \lambda^{-1} \bar{\eta}^i) \right)$$

is an isomorphism (cf. Lemmas 3.2 and 3.4 and condition (30)) and the operator which takes $(\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i)$ to

$$\begin{aligned} &\left(v_{\Omega^i}^- [\bar{\eta}^o]_{|\partial\Omega^o}, 0, W_{\Omega^i}^* [\bar{\eta}^o] - \lambda^{-1} W_{\Omega^i}^* [\bar{\eta}^i] \right. \\ &\quad \left. + (\lambda - 1) \lambda^{-1} \nu_{\Omega^i} \cdot \nabla w_{\Omega^o}^+ [\bar{\mu}^o]_{|\partial\Omega^i} - \tilde{\gamma} (\lambda^{-1} w_{\Omega^o}^+ [\bar{\mu}^o]_{|\partial\Omega^i} + \lambda^{-1} v_{\Omega^i}^+ [\bar{\eta}^i]_{|\partial\Omega^i}) \right) \end{aligned}$$

is compact (by the properties of integral operators with real analytic kernels and no singularity and by Lemma 3.1). Hence, to prove the statement of the lemma, it suffices to show that

$$\partial_{(\mu^o, \eta^o, \eta^i)} N[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i](\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) = 0$$

implies $(\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) = (0, 0, 0)$. If $\partial_{(\mu^o, \eta^o, \eta^i)} N[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i](\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) = 0$, then by the jump properties of the single and double layer potentials the pair $(U^o[\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i], U^i[\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i])$ is a solution of the problem

$$\begin{cases} \Delta u^o = 0 & \text{in } \Omega^o \setminus \text{cl}\Omega^i, \\ \Delta u^i = 0 & \text{in } \Omega^i, \\ u^o(x) = 0 & \text{for all } x \in \partial\Omega^o, \\ u^o(x) = \lambda u^i(x) & \text{for all } x \in \partial\Omega^i, \\ \nu_{\Omega^i} \cdot \nabla u^o(x) - \nu_{\Omega^i} \cdot \nabla u^i(x) = \tilde{\gamma}(x) u^i(x) & \text{for all } x \in \partial\Omega^i. \end{cases}$$

Then, by inequalities $\lambda > 0$ and $\tilde{\gamma} \geq 0$ (cf. condition (38)) and by a standard energy argument one verifies that $(U^o[\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i], U^i[\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i]) = (0, 0)$. Thus $(\bar{\mu}^o, \bar{\eta}^o, \bar{\eta}^i) = (0, 0, 0)$ by Lemma 5.2 and the proof is completed. \square

Then, by Lemma 5.8 and by the implicit function theorem (see, e.g., Deimling [9, §15]) one verifies the validity of the following proposition.

Proposition 5.9. *Let Φ and G satisfy (37). Let $(\tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i) \in C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)^2$ and $N[0, \tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i] = 0$. Assume that condition (38) holds true. Then there exist $\epsilon^* > 0$, a neighbourhood \mathcal{U} of $(\tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i)$ in $C^{1,\alpha}(\partial\Omega^i) \times C^{0,\alpha}(\partial\Omega^i)^2$, and a continuously Fréchet differentiable map $(\mu^o[\cdot], \eta^o[\cdot], \eta^i[\cdot])$ from $] - \epsilon^*, \epsilon^*[$ to \mathcal{U} such that the set of zeros of N in $] - \epsilon^*, \epsilon^*[\times \mathcal{U}$ coincides with the graph of $(\mu^o[\cdot], \eta^o[\cdot], \eta^i[\cdot])$. In particular, $N[\epsilon, \mu^o[\epsilon], \eta^o[\epsilon], \eta^i[\epsilon]] = 0$ for all $\epsilon \in] - \epsilon^*, \epsilon^*[$ and $(\mu^o[0], \eta^o[0], \eta^i[0]) = (\tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i)$.*

We are now ready to prove the main Theorem 5.10 of this section.

Theorem 5.10. *Let Φ and G satisfy condition (37). Assume that $|G(x, t)| \leq C(1 + |t|)^\delta$ for some $C > 0$, $\delta \in [0, 1[$ and for all $(x, t) \in \partial\Omega^i \times \mathbb{R}$. Then the following statement hold:*

- (i) *there exists at least a solution $(u_0^o, u_0^i) \in C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ of the boundary value problem in (7) with $\epsilon = 0$.*

If in addition we have

$$(\partial_t G)(x, u_0^i(x)) \geq 0 \quad \forall x \in \partial\Omega^i \quad (39)$$

then there exist $\epsilon_* > 0$ and a family of functions $\{(u_\epsilon^o, u_\epsilon^i)\}_{\epsilon \in]-\epsilon_*, \epsilon_*[\setminus \{0\}}$ such that following statements hold:

- (ii) for all $\epsilon \in]-\epsilon_*, \epsilon_*[$ the pair $(u_\epsilon^o, u_\epsilon^i)$ belongs to $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ and is a solution of (7);
- (iii) the map from $] - \epsilon_*, \epsilon_*[$ to $C^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C^{1,\alpha}(\text{cl}\Omega^i)$ which takes ϵ to $(u_\epsilon^o, u_\epsilon^i)$ is continuously Fréchet differentiable;
- (iv) there exists an open subset \mathcal{V} of $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ such that, for all fixed $\epsilon \in] - \epsilon_*, \epsilon_*[$ the pair $(u_\epsilon^o, u_\epsilon^i)$ is the unique solution of (7) belonging to \mathcal{V} .

Proof. (i) By Lemma 5.5 there exists at least a solution $(\mu_0^o, \eta_0) \in C^{1,\alpha}(\partial\Omega^o) \times C^{0,\alpha}(\partial\Omega^i)$ of (34). Then we define $(u_0^o, u_0^i) \equiv (U^o[\mu_0^o, \eta_0, \eta_0], U^i[\mu_0^o, \eta_0, \eta_0])$ and the validity of statement (i) follows by Lemma 5.4 (see also Proposition 5.9).

(ii) Since (μ_0^o, η_0) is a solution of (34), we have $N[0, \mu_0^o, \eta_0, \eta_0] = 0$. Then let $(\tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i) \equiv (\mu_0^o, \eta_0, \eta_0)$. By condition (39) and by the jump properties of single and double layer potential, one verifies that condition (38) is satisfied. Accordingly, the assumption of Proposition 5.9 are fulfilled and we can take $\epsilon_* \equiv \epsilon^*$ and define $(u_\epsilon^o, u_\epsilon^i) \equiv (U^o[\mu^o[\epsilon], \eta^o[\epsilon], \eta^i[\epsilon]], U^i[\mu^o[\epsilon], \eta^o[\epsilon], \eta^i[\epsilon]])$ for all $\epsilon \in] - \epsilon_*, \epsilon_*[$. The validity of (ii) follows by Lemma 5.7.

(iii) It is a consequence of the continuous Fréchet differentiability of $(\mu^o[\cdot], \eta^o[\cdot], \eta^i[\cdot])$, of the definition of (U^o, U^i) in Lemma 5.2, and of the mapping properties of the single and double layer potentials (cf. Miranda [32]).

(iv) Let \mathcal{U} be the open neighbourhood of $(\tilde{\mu}^o, \tilde{\eta}^o, \tilde{\eta}^i)$ introduced in Proposition 5.9. Let $\mathcal{V} \equiv \{(U^o[\mu^o, \eta^o, \eta^i], U^i[\mu^o, \eta^o, \eta^i]) : (\mu^o, \eta^o, \eta^i) \in \mathcal{U}\}$. Since $U = (U^o, U^i)$ is an open operator the set \mathcal{V} is open in $C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^o \setminus \Omega^i) \times C_{\text{harm}}^{1,\alpha}(\text{cl}\Omega^i)$ (cf. Lemma 5.2). Moreover, the pair of functions $(u_\epsilon^o, u_\epsilon^i) = (U^o[\mu^o[\epsilon], \eta^o[\epsilon], \eta^i[\epsilon]], U^i[\mu^o[\epsilon], \eta^o[\epsilon], \eta^i[\epsilon]])$ belongs to \mathcal{V} for all $\epsilon \in] - \epsilon_*, \epsilon_*[$ (cf. Proposition 5.9). Now fix $\epsilon_\# \in] - \epsilon_*, \epsilon_*[$ and assume that $(u_{\epsilon_\#}^o, u_{\epsilon_\#}^i) \in \mathcal{V}$ is a solution of (7) for $\epsilon = \epsilon_\#$. Then there exists $(\mu_{\epsilon_\#}^o, \eta_{\epsilon_\#}^o, \eta_{\epsilon_\#}^i) \in \mathcal{U}$ such that $(u_{\epsilon_\#}^o, u_{\epsilon_\#}^i) = (U^o[\mu_{\epsilon_\#}^o, \eta_{\epsilon_\#}^o, \eta_{\epsilon_\#}^i], U^i[\mu_{\epsilon_\#}^o, \eta_{\epsilon_\#}^o, \eta_{\epsilon_\#}^i])$ (see Lemma 5.2). Moreover, $N[\epsilon_\#, \mu_{\epsilon_\#}^o, \eta_{\epsilon_\#}^o, \eta_{\epsilon_\#}^i] = 0$ by Lemma 5.7 and thus $(\mu_{\epsilon_\#}^o, \eta_{\epsilon_\#}^o, \eta_{\epsilon_\#}^i) = (\mu^o[\epsilon_\#], \eta^o[\epsilon_\#], \eta^i[\epsilon_\#])$ by Proposition 5.9. Accordingly $(u_{\epsilon_\#}^o, u_{\epsilon_\#}^i) = (u_{\epsilon_\#}^o, u_{\epsilon_\#}^i)$ and the proof is complete. \square

Acknowledgment

The research of M. Dalla Riva was supported by Portuguese funds through the CIDMA - Center for Research and Development in Mathematics and Applications, and the Portuguese Foundation for Science and Technology (“FCT–Fundação para a Ciência e a Tecnologia”), within project UID/MAT/04106/2013. The research of M. Dalla Riva was also supported by the Portuguese Foundation for Science and Technology (“FCT–Fundação para a Ciência e a Tecnologia”) with the research grant SFRH/BPD/ 64437/2009. In addition, the work of M. Dalla Riva was supported by “Progetto di Ateneo: Singular perturbation problems for differential operators – CPDA120171/12” of the University of Padova.

G. Mishuris acknowledges the support of the European Community’s Seven Framework Programme under contract number PIAPP-GA-284544-PARM-2.

Finally, M. Dalla Riva wishes to thank the Department of Mathematics of the University of Aberystwyth for the hospitality received during the development of a part of the work.

- [1] I. Babuška, M. Suri, On locking and robustness in the finite element method, *SIAM J. Numer. Anal.* 29 (1992) 1261–1293.
- [2] G.R. Barrenechea, G.N. Gatica, On the coupling of boundary integral and finite element methods with nonlinear transmission conditions, *Appl. Anal.* 62 (1996) 181–210.
- [3] H. Berger, A convergent finite element formulation for transonic flow, *Numer. Math.* 56 (1989) 425–447.
- [4] H. Berger, G. Warnecke, W.L. Wendland, Finite elements for transonic potential flows, *Numer. Methods Partial Differ. Equations* 6 (1990) 17–42.
- [5] T. Carleman, Über eine nichtlineare Randwertaufgabe bei der Gleichung $\Delta u = 0$ (in German), *Math. Zeitschr.* 9 (1921) 35–43.
- [6] M. Costabel, Boundary integral operators in Lipschitz domains: elementary results, *SIAM J. Math. Anal.* 19 (1988) 613–626.

- [7] M. Costabel, E.P. Stephan, Coupling of finite and boundary element methods for an elastoplastic interface problem, *SIAM J. Numer. Anal.* 27 (1990) 1212–1226.
- [8] J.M. Cushing, Some existence theorems for non-linear eigenvalue problems associated with elliptic equations, *Arch. Ration. Mech. Anal.* 42 (1971) 63–76.
- [9] K. Deimling, *Nonlinear functional analysis*, Springer Verlag, Berlin, 1985.
- [10] M. Dalla Riva, M. Lanza de Cristoforis, Hypersingularly perturbed loads for a nonlinear traction boundary value problem. A functional analytic approach, *Eurasian Mathematical Journal* 1 (2010) 31–58.
- [11] M.A. Efendiev, H. Schmitz, W.L. Wendland, On some nonlinear potential problems, *Electron. J. Differ. Equ.* paper 18 (1999) 1–17.
- [12] L. Escauriaza, E.B. Fabes, G. Verchota, On a regularity theorem for weak solutions to transmission problems with internal Lipschitz boundaries, *Proc. Am. Math. Soc.* 115 (1992) 1069–1076.
- [13] L. Escauriaza, J.K. Seo, Regularity properties of solutions to transmissions problems, *Trans. Amer. Math. Soc.* 338 (1993) 405–430.
- [14] L. Escauriaza, M. Mitrea, Transmission problems and spectral theory for singular integral operators on Lipschitz domains, *J. Func. Anal.* 216 (2004) 141–171.
- [15] E.B. Fabes, M. Jodeit, Jr., J.E. Lewis, Double layer potentials for domains with corners and edges, *Indiana Univ. Math. J.* 26 (1977) 95–114.
- [16] E.B. Fabes, M. Jodeit, Jr., N.M. Rivière, Potential techniques for boundary value problems on C^1 -domains, *Acta Math.* 141 (1978) 165–186.
- [17] G.B. Folland, *Introduction to partial differential equations*, second ed., Princeton University Press, Princeton, NJ, 1995.
- [18] G.N. Gatica, G.C. Hsiao, On the coupled BEM and FEM for a nonlinear exterior Dirichlet problem in \mathbb{R}^2 , *Numer. Math.* 61 (1992) 171–214.

- [19] G.N. Gatica, G.C. Hsiao, The uncoupling of boundary integral and finite element methods for nonlinear boundary value problems, *J. Math. Anal. Appl.*, 189 (1995) 442–461.
- [20] D. Gilbarg, N.S. Trudinger, *Elliptic partial differential equations of second order*, reprint of the 1998 ed., Springer Verlag, Berlin, 2001.
- [21] I. Hlaváček, M. Křížek, J. Malý, On Galerkin approximations of a quasilinear nonpotential elliptic problem of a nonmonotone type, *J. Math. Anal. Appl.* 184 (1994) 168–189.
- [22] C. Jacob, Sur la détermination des fonctions harmoniques conjuguées par certaines conditions aux limites. Applications à l'hydrodynamique (in French), *Mathematica, Cluj*, 11, 1935.
- [23] K. Klingelhöfer, Nonlinear harmonic boundary value problems. I, *Arch. Ration. Mech. Anal.* 31 (1968) 364–371.
- [24] K. Klingelhöfer, Modified Hammerstein integral equations and nonlinear harmonic boundary value problems, *J. Math. Anal. Appl.* 28 (1969) 77–87.
- [25] M. Kohr, M. Lanza de Cristoforis, and W.L. Wendland, Nonlinear Neumann–transmission problems for Stokes and Brinkman equations on euclidean Lipschitz domains, *Potential Anal.* 38 (2013) 1123–1171.
- [26] R. Kress, *Linear integral equations*, second ed., Springer Verlag, Berlin, 1989.
- [27] M. Lanza de Cristoforis, Asymptotic behavior of the solutions of a nonlinear Robin problem for the Laplace operator in a domain with a small hole: a functional analytic approach, *Complex Var. Elliptic Equ.* 52 (2007) 945–977.
- [28] M. Lanza de Cristoforis, Asymptotic behaviour of the solutions of a nonlinear transmission problem for the Laplace operator in a domain with a small hole. A functional analytic approach, *Complex Var. Elliptic Equ.* 55 (2010) 269–303.
- [29] M. Lanza de Cristoforis, P. Musolino, A quasi-linear heat transmission problem in a periodic two-phase dilute composite. A functional analytic approach, *Comm. Pure Appl. Anal.* 13 (2014) 2509–2542.

- [30] J. Leray, Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l'hydrodynamique (in French), *J. Math. Pures Appl.*, IX, Sér. 12 (1933) 1–82.
- [31] S.E. Mikhailov, Traces, extensions and co-normal derivatives for elliptic systems on Lipschitz domains, *J. Math. Anal. Appl.* 378 (2011) 324–342.
- [32] C. Miranda, Sulle proprietà di regolarità di certe trasformazioni integrali (in Italian), *Atti Accad. Naz. Lincei Mem. Cl. Sci. Fis. Mat. Natur. Sez. I* 7 (1965) 303–336.
- [33] C. Miranda, *Partial differential equations of elliptic type*, Springer Verlag, Berlin, 1970.
- [34] G. Mishuris, W. Miszuris, A. Öchsner, Evaluation of transmission conditions for thin reactive heat-conducting interphases, *Defect Diffus. Forum* 273–276 (2008) 394–399.
- [35] G. Mishuris, W. Miszuris, A. Öchsner, Transmission conditions for thin reactive heat-conducting interphases: general case, *Defect Diffus. Forum* 283–286 (2009) 521–526.
- [36] W. Miszuris, A. Öchsner, Universal transmission conditions for thin reactive heat-conducting interphases, *Continuum Mechanics and Thermodynamics* 25 (2013) 1–21.
- [37] V.V. Mityushev, S.V. Rogosin, *Constructive methods for linear and nonlinear boundary value problems for analytic functions: theory and applications*, Chapman & Hall/CRC, Boca Raton, FL, 2000.
- [38] K. Nakamori, Y. Suyama, Zum nicht-linearen Randwertproblem der Gleichungen $\Delta u = 0$ and $\Delta u = f(x, y)$ (in Esperanto), *Mem. Fac. Sci. Kyusyu Univ. Ser. A5* (1950) 99–106.
- [39] J. Nečas, *Introduction to the theory of nonlinear elliptic equations*, Teubner-Texte zur Mathematik 52, Teubner, Leipzig, 1983.
- [40] F. Rosselli, P. Carbutt, Structural bonding applications for the transportation industry, *SAMPE J.* 37 (2001) 7–13.

- [41] T. Roubíček, Nonlinear partial differential equations with applications, second ed., Intl. Ser. Numer. Math 153, Birkhäuser, Basel, Boston, Berlin, 2013.
- [42] J. Schauder, Potentialtheoretische Untersuchungen (in German), Math. Z. 33 (1931) 602–640.
- [43] J. Schauder, Bemerkung zu meiner Arbeit “Potentialtheoretische Untersuchungen I (Anhang)” (in German), Math. Z. 35 (1932) 536–538.