

## Aberystwyth University

### *The Diterpenoid 7-Keto-Sempervirol, Derived from Lycium chinense, Displays Anthelmintic Activity against both Schistosoma mansoni and Fasciola hepatica*

Edwards, Jennifer; Brown, Martha; Peak, Emily; Bartholomew, Barbara; Nash, Robert J; Hoffmann, Karl F

*Published in:*

PLoS Neglected Tropical Diseases

*DOI:*

[10.1371/journal.pntd.0003604](https://doi.org/10.1371/journal.pntd.0003604)

*Publication date:*

2015

*Citation for published version (APA):*

Edwards, J., Brown, M., Peak, E., Bartholomew, B., Nash, R. J., & Hoffmann, K. F. (2015). The Diterpenoid 7-Keto-Sempervirol, Derived from *Lycium chinense*, Displays Anthelmintic Activity against both *Schistosoma mansoni* and *Fasciola hepatica*. *PLoS Neglected Tropical Diseases*, 9(3), Article e0003604. <https://doi.org/10.1371/journal.pntd.0003604>

#### Document License

CC BY

#### General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400

email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

## HepG2 cell line / 7-keto sempervirol titration





