

## Aberystwyth University

### *Seed Number and 100-Seed Weight of Pearl Millet (*Pennisetum glaucum* L.) Respond Differently to Low Soil Moisture in Genotypes Contrasting for Drought Tolerance*

Aparna, K.; Hash, C. T.; Yadav, Rattan Singh; Vadez, Vincent

*Published in:*  
Journal of Agronomy and Crop Science

*DOI:*  
[10.1111/jac.12052](https://doi.org/10.1111/jac.12052)

*Publication date:*  
2014

*Citation for published version (APA):*  
Aparna, K., Hash, C. T., Yadav, R. S., & Vadez, V. (2014). Seed Number and 100-Seed Weight of Pearl Millet (*Pennisetum glaucum* L.) Respond Differently to Low Soil Moisture in Genotypes Contrasting for Drought Tolerance. *Journal of Agronomy and Crop Science*, 200(2), 119-131. <https://doi.org/10.1111/jac.12052>

#### **Document License** CC BY

#### **General rights**





Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)



**Supplementary Figure 1** Daily transpiration of two pearl millet test cross hybrids of parental lines PRLT2/89-33 (tolerant), H77/833-2 (sensitive), their NILs ICMH 1046, ICMH 1029 (a) and 863B (tolerant), 841B (sensitive) parental lines of a mapping population (c) grown under well watered (WW) and water stress (WS) conditions i.e. 0.3 FTSW (fraction of transpirable soil water) after flowering. NTR (normalized transpiration ratio) of these parental lines PRLT2/89-33 (tolerant), H77/833-2 (sensitive), their NILs ICMH 1046, ICMH 1029 (b) and 863B (tolerant), 841B (sensitive) (d) at 0.3 FTSW. Each data point represents NTR at each day after flowering.