

Aberystwyth University

Temporal dynamics of the metabolically active rumen bacteria colonising fresh perennial ryegrass

Huws, Sharon; Edwards, Joan Elizabeth; Creevey, Christopher; Rees Stevens, Pauline; Lin, Wanchang; Girdwood, Susan; Pachebat, Justin; Kingston-Smith, Alison

Published in:
FEMS Microbiology Ecology

DOI:
[10.1093/femsec/fiv137](https://doi.org/10.1093/femsec/fiv137)

Publication date:
2016

Citation for published version (APA):
Huws, S., Edwards, J. E., Creevey, C., Rees Stevens, P., Lin, W., Girdwood, S., Pachebat, J., & Kingston-Smith, A. (2016). Temporal dynamics of the metabolically active rumen bacteria colonising fresh perennial ryegrass. *FEMS Microbiology Ecology*, 92(1), [fiv 137]. <https://doi.org/10.1093/femsec/fiv137>

Document License CC BY-NC

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Temporal dynamics of the metabolically active rumen bacteria colonising fresh perennial ryegrass

Sharon A. Huws^{a*}, Joan. E. Edwards^{a,b*}, Christopher J. Creevey, Pauline Rees Stevens^a,
Wanchang Lin^a, Susan E. Girdwood^a, Justin A. Pachebat^a, Alison H. Kingston-Smith^a

^aAberystwyth University, Aberystwyth, SY23 3FG, UK; ^bCurrent address: Wageningen
University, Wageningen, 6703 HB, Netherlands; *these authors contributed equally to the
study.

RUNNING TITLE: Diversity of rumen bacteria attached to perennial ryegrass.

***Correspondence:** Sharon A. Huws, Animal and Microbial Sciences, Institute of Biological,
Environmental and Rural Sciences (IBERS), Aberystwyth University, Penglais Campus,
Aberystwyth, SY23 3FG, UK. E-mail: hnh@aber.ac.uk. Tel: +44 1970 823202 and Fax: +44
1970 823155.

Keywords: Rumen, perennial ryegrass, bacteria, microbiome, metagenome, 16S, RNA.

21 **ABSTRACT**

22 This study investigated successional colonisation of fresh perennial ryegrass (PRG) by the
23 rumen microbiota over time. Fresh PRG was incubated *in sacco* in the rumens of three
24 Holstein x Friesian cows over a period of 8 h, with samples recovered at various times. The
25 diversity of attached bacteria was assessed using 454 pyrosequencing of 16S rRNA (cDNA).
26 Results showed that plant epiphytic communities either decreased to low relative abundances
27 or disappeared following rumen incubation, and that temporal colonisation of the PRG by the
28 rumen bacteria was biphasic with primary (1 & 2 h) and secondary (4-8 h) events evident
29 with the transition period being with 2-4 h. A decrease in sequence reads pertaining to
30 *Succinivibrio* spp. and increases in *Pseudobutyrvibrio*, *Roseburia* and *Ruminococcus* spp.
31 (the latter all order Clostridiales) were evident during secondary colonisation. Irrespective of
32 temporal changes, the continually high abundances of *Butyrivibrio*, *Fibrobacter*, *Olsenella*
33 and *Prevotella* suggest that they play a major role in the degradation of the plant. It is clear
34 that a temporal understanding of the functional roles of these, and the colonisation specific,
35 microbiota within the rumen is now required to unravel the role of these bacteria in the
36 ruminal degradation of fresh perennial ryegrass.

37

38 INTRODUCTION

39 Ruminant animals supply much of our meat and nearly all of our milk requirements,
40 and as such are important to human nutrition. Globally, increased demand coupled with a
41 growing population means that ruminant products will become increasingly scarce
42 (Foresight, 2011). Ruminants convert plant biomass to chemical compounds, which are
43 subsequently metabolised and absorbed by the animal. This process is largely due to the
44 functional capacity of the rumen microbiome (Mackie, 2002; Edwards *et al.*, 2008a; Brulc *et*
45 *al.*, 2009; Kingston Smith *et al.*, 2010; Kav *et al.*, 2012). Furthermore, the fermentative
46 capacity of the rumen microbiota enables microbial breakdown of otherwise undigestible
47 dietary material and thus defines the amount, quality and composition of the meat and milk
48 produced (Edwards *et al.*, 2008a; Brulc *et al.*, 2009; Kim *et al.*, 2009; Kingston Smith *et al.*,
49 2010; Kav *et al.*, 2012; Huws *et al.*, 2014b).

50 The rumen microbiota rapidly colonise ingested feed particles which ultimately
51 results in the microbial degradation of the plant material, causing release of bioavailable
52 nutrients (Cheng *et al.*, 1980; Miron *et al.*, 2001; Russell and Rychlik, 2001; Koike *et al.*,
53 2003; Edwards *et al.*, 2007, 2008b; Huws *et al.*, 2013ab; 2014a). Nonetheless, the process is
54 relatively inefficient in terms of animal production, with as little as 30% of the ingested
55 nitrogen being retained by the animal for milk or meat production and the non-incorporated
56 nitrogen is excreted as urea or ammonia (MacRae and Ulyatt, 1974; Dewhurst *et al.*, 1996;
57 Kingston-Smith *et al.*, 2008, 2010). This presents a major challenge in terms of increasing
58 ruminant productivity and consequently in providing a sustainable supply of meat and milk
59 for the future. As the attachment of rumen microbiota to ingested forage is central for the
60 availability of nutrients to the ruminant (McAllister *et al.*, 1994; Dewhurst *et al.*, 1996;
61 Kingston-Smith *et al.*, 2010), understanding rumen plant-microbe interactions is paramount
62 in order to develop novel methodologies for increasing nutrient use efficiency within

63 ruminants (Leng, 2014). For example, by furthering our fundamental understanding of
64 temporal plant nutrient breakdown and availability it is possible to define key limitations and
65 enhance the chemical characteristics of available forages through targeted plant breeding to
66 circumvent these limitations. Thus, fundamental information on the temporal plant-microbe
67 interactome can inform plant breeding strategies with the ultimate aim of increasing animal
68 nutrient use efficiency whilst decreasing environmental impact.

69 We have previously shown using Denaturing Gradient Gel Electrophoresis (DGGE)
70 that the perennial ryegrass attached microbiome changes in diversity between 2 and 4 h of
71 incubation, but the capacity of this technique is limited in terms of understanding which
72 bacteria change in abundance within these primary and secondary colonisation events (Huws
73 *et al.*, 2013b; Huws *et al.*, 2013a). In this study we used 454 based 16S rRNA (cDNA)
74 sequencing to characterise time-related changes in the diversity of the rumen bacteria
75 attaching to fresh perennial ryegrass (PRG). By basing these experiments on RNA rather
76 than DNA, these data provide an insight specifically into the metabolically active rumen
77 PRG-attached microbiome.

78

79 **MATERIALS AND METHODS**

80 **Growth and preparation of plant material**

81 Perennial ryegrass, (*Lolium perenne* cv. AberDart; PRG) was grown from seed in plastic seed
82 trays (length 38 cm x width 24 cm x depth 5 cm) filled with soil/compost (Levingtons general
83 purpose). The trays were housed in a greenhouse under natural irradiance with additional
84 illumination provided during the winter months (minimum 8 h photoperiod). A temperature
85 of 22/19°C day/night was maintained and plants were watered twice a week. Plants were
86 harvested after 6 weeks by cutting 3 cm above soil level, before cutting with scissors into 1

87 cm sections just prior to incubation in the rumen as described below. Samples of this
88 harvested plant material were also snap frozen in dry ice, and stored at -80°C for bacterial
89 profiling (0 h samples).

90

91 ***In sacco* incubations**

92 Three mature, rumen-cannulated, non-lactating Holstein x Friesian cows were used for this
93 experiment. Experiments were conducted with the authority of Licenses under the United
94 Kingdom Animal Scientific Procedures Act, 1986. For at least 2 weeks prior to the
95 experiments, the cows were fed a diet of straw and grass silage *ad libitum* (~6.5 kg dry matter
96 day⁻¹) and were also permitted field grazing on PRG for at least 4 h/day. For the duration of
97 the experiment animals were fed silage twice daily (07:00 and 16:00). Stitched nylon bags
98 (10 cm × 20 cm) of 100 µm² pore sizes were filled with 15 g (fresh weight) of the processed
99 plant material and sealed at all perimeters by heating (Impulse sealer, American Int, NI
100 Electric, AIE, USA). The nylon bag technique was adopted as described previously (Ørskov
101 *et al.*, 1980; Vanzant *et al.*, 1998). Essentially, bags were connected to a 55-cm, coated
102 flexible plastic cable with lacing cords and this was placed in the rumen and attached to the
103 cap of the fistula. Bags were placed simultaneously in the rumen of each cow shortly after
104 animals were offered the first meal of the morning and removed after 1, 2, 4, 6 and 8 h of
105 incubation. At each time interval, twelve bags (four from each cow) were withdrawn and the
106 residual plant material in six of the bags (two for each cow) was processed by washing with
107 distilled water (500 ml added to plant material within bags and bags gently squeezed
108 thereafter) to remove loosely attached microbes followed by oven drying for two replicate
109 samples from each cow and calculation of plant degradation (% dry matter lost). The

110 remaining six bags (two for each cow) were similarly washed with distilled water before
111 being immediately frozen in dry ice and then stored at -80°C until RNA extraction.

112

113 **RNA extraction**

114 Frozen samples were ground to a fine powder under liquid nitrogen before RNA was
115 extracted using a hot phenol method (Ougham and Davies, 1990). Essentially aquaphenol
116 (10 mL) was added to the ground sample prior to incubation at 65°C for 1 h. Tubes were
117 inverted before chloroform was added (5 mL). Tubes were centrifuged (5,000 x g, 30 mins,
118 20°C) before upper phase was removed then the procedure was repeated by addition of more
119 chloroform (5 mL) and centrifugation as described. Lithium chloride (2M final
120 concentration) was then added, to remove any contaminating DNA, and samples stored
121 overnight at 4°C. Samples were subsequently centrifuged (13,000 x g, 30 mins, 4°C) and
122 supernatant discarded, then the procedure was repeated from addition of lithium chloride to
123 ensure all DNA was removed. Once the supernatant was discarded the pellet was
124 resuspended in ice cold 80% ethanol and centrifuged (13,000 x g, 15 mins, 4°C), this was
125 repeated twice before the pellet was air dried and resuspended in molecular grade water.
126 Quality and quantity of retrieved RNA was checked using the Experion automated
127 electrophoresis system and RNA 'stdsens' chips for standard sensitivity analysis (Bio-rad,
128 Hemel Hempstead, UK).

129

130 **16S rRNA 454 pyrosequencing**

131 RNA (c. 100 ng) was reverse transcribed using the reverse primer R1401
132 (5'GGGTCTTGTACACACCG 3') and Superscript III reverse transcriptase (Invitrogen Ltd,

133 Paisley, UK) in 20 uL reactions, following the manufacturer's guidelines and as previously
134 described by Edwards *et al.* (2007) and Huws *et al.* (2011; 2013a). Control reactions were
135 performed with no reverse transcriptase, and were PCR amplified (as described below) to
136 confirm that the RNA preparations were free of contaminating DNA. Amplicons of the V6-
137 V8 variable region of the bacterial 16S rDNA gene were generated in triplicate per cDNA
138 sample by PCR using the primers 799F2 (5' tagged with Roche B adaptor) and R1401 (5'-
139 tagged with the Roche A adaptor and MID barcode tags specific for each sample as suggested
140 by Roche) as described by Edwards *et al.* (2007), except that 30 cycles of amplification was
141 used. All PCR products were initially verified by electrophoretic fractionation on a 1.0%
142 agarose gel for 1 h, 120 V, 80 mA in 1% TAE (*Tris* base, acetic acid and EDTA) buffer
143 before pooling of triplicate amplifications. The pooled PCR products (30 µl each sample)
144 were subsequently run on a 2.0% agarose gel for 2 h, 120 V, 80 mA in 1% TAE buffer before
145 bands were viewed and cut on a dark reader transilluminator (Clare Chemical Research Inc.,
146 Colorado, USA). Amplicons were retrieved from cut bands using the Isolate II PCR and gel
147 Kit (Bioline, London, UK). Purified amplicons were verified and quantified using the
148 Agilent High Sensitivity Assay Kit (Agilent Technologies, California, USA) prior to
149 pyrosequencing using Titanium chemistry on a Roche GS-FLX 454 sequencer (Roche
150 Diagnostics Ltd, West Sussex, UK) according to the manufacturer's guidelines. These
151 sequences and associated metadata can be accessed through the NCBI bioproject ID
152 PRJNA274256.

153

154 **Data Analysis**

155 All 16S rRNA sequences with a length less than 400bp were discarded and those remaining
156 clustering at 97% identity using CD-HIT-OTU (Li *et al.*, 2012) were analysed to identify
157 Operational Taxonomic Units (OTUs). OTUs with fewer than 5 representatives and those

158 found to be chimeric were removed from subsequent analyses. Abundances of each of the
159 remaining OTUs were calculated using the “clstr_sample_count_matrix.pl” script from the
160 CD-HIT-OTU package. These counts were then used as input to the Bioconductor package
161 DESEQ2 in R (Love *et al.*, 2014) to identify overall changes in the attached microbiota.
162 Taxonomic identification of the OTUs was carried out using the classifier algorithm from the
163 RDP database (Cole *et al.*, 2014). Any taxonomic identification below 90% identity to
164 published sequences was not included in the analysis. Further statistical analyses of changes
165 at the Phylum, Order, Family and Genus level were carried out by ANOVA in Genstat
166 (Payne *et al.*, 2007). Only those genera that were present in greater than 1% of the total
167 microbiome in any time point were included. Dry matter data was also analysed by ANOVA
168 and Genstat (Payne *et al.*, 2007). PCA plots were generated using the ggplot2 library and
169 rarefaction curves were drawn using the vegan package in R. Data was transformed into a
170 heat map using the Heatmap2 package from the Gplots package in R using the summed
171 abundances of each OTU from the genera indicated.

172

173 **RESULTS**

174 **16S rRNA sequencing data**

175 Overall, 1,411,847 sequences were generated, of which 1,016,349 (72%) had a length greater
176 than 400bp (Figure S1). This consisted of an average of 41,646 ($\pm 4,762$ standard deviation)
177 sequences/sample pre-filtering and an average of 31,761 ($\pm 3,523$ standard deviation)
178 sequences/sample post-filtering (Table S1). After filtering, the average sequence length was
179 425 bp. Following removal of low abundance and chimeric OTUs 1,201 OTUs remained,
180 which is consistent with previous reports (Creevey *et al.* 2014). The average number of
181 sequences per sample, assigned to an OUT, which had a taxonomic classification >90%, was

182 9,093 ($\pm 1,274$) (Table S1). The most abundant OTU (from *Butyrivibrio*) had 29,722
183 representatives, representing 10% of all sequences found across all time points. The OTU
184 based rarefaction curve plateaued indicating that a reasonable level of sequencing depth was
185 obtained (Figure S2).

186

187 **Fate of the perennial ryegrass epiphytic microbiota post rumen incubation**

188 16S rRNA pyrosequences showed that the relative abundances of the plant epiphytic
189 communities decreased substantially to very low levels within the first hour of rumen
190 incubation with some decreasing to below detection limits (Tables S2-S5 and Figure 1).

191

192 **Temporal diversity of the PRG attached microbiota post 1 h of rumen incubation**

193 The PCA plot of OTU abundances showed that the microbiota attached to fresh PRG at 1 & 2
194 h differed significantly ($P < 0.05$) from those attached during 4-8 h of incubation (Figure 2).
195 This was in agreement with the results of Denaturing gradient gel electrophoresis (DGGE),
196 performed as described by Edwards *et al.* (2007) prior to sequencing which indicated
197 biphasic colonisation, in which the primary (1 & 2 h) and secondary (4-8 h) phases had
198 approximately 60-75% DGGE profile similarity dependent on cow (data not shown).
199 Shannon diversity boxplots based on OTU abundance showed a higher bacterial diversity ≥ 5
200 following 1 and 2 h of incubation, which then decreased significantly ($P < 0.05$) between 4-6 h
201 of incubation (Figure 3). The Shannon diversity at 8 h was not significantly different from
202 the other time points.

203

204 **Phyla level temporal diversity of the attached microbiota post 1 h of rumen incubation**

205 On a phylum level the most abundant attached bacteria were Firmicutes, Bacteroidetes and
206 Fibrobacteres (on average approx. 75, 17 and 4% of total average normalised reads across
207 time points respectively) (Table 1), whereas a further 8 phyla were relatively minor (<2% of
208 total normalised reads/phyla) in comparison. In terms of temporal changes within the more
209 predominant attached bacterial phyla, Firmicutes changed in abundance significantly over
210 time, with the greatest abundance observed during secondary colonisation (4-8 h) ($P<0.05$)
211 (Table 1). Bacteroidetes and Fibrobacteres read abundances did not change significantly over
212 time ($P>0.05$) (Table 1). Despite their lower abundances the reads pertaining to phyla
213 Actinobacteria, Elusimicrobia, Lentisphaerae and Verrucomicrobia also changed significantly
214 over time ($P<0.05$) (Table 1). More sequences pertaining to the phyla Actinobacteria were
215 present at 2 h of rumen incubation compared to all other time points ($P<0.05$) (Table 1).
216 Elusimicrobia and Verrucomicrobia read abundances decreased significantly post 1 h and 2 h
217 of rumen incubation respectively ($P<0.05$) (Table 1). Reads pertaining to Lentisphaerae were
218 maximal at 1 h of incubation ($P<0.05$) (Table 1). Fusobacteria, Proteobacteria, Spirochaetes,
219 and Tenericutes read abundances did not change significantly over time ($P>0.05$) (Table 1).

220

221 **Order level temporal diversity of the attached microbiome post 1 h of rumen incubation**

222 On an order level the most abundant attached bacteria were Clostridiales, Bacteriodales
223 Selenomonadales, Fibrobacterales, Coriobacteriales and Spirochaetales (on average approx.
224 67, 17, 7, 4, 3 and 2% of total average normalised reads across time points respectively),
225 whereas a further 20 orders were relatively minor (<2% of total normalised reads/phyla) in
226 comparison (Table 2). The order Clostridiales changed significantly in abundance over time,
227 with increased abundances present during secondary colonisation events (4-8 h) ($P<0.05$)
228 (Table 2). Bacteroidales, Fibrobacterales and Spirochaetales read abundances did not change

229 significantly over time ($P>0.05$) (Table 2). Read abundances pertaining to the order
230 Selenomonadales changed significantly over time, with significantly higher abundances
231 present at 2 and 4 h compared with read abundances at 8 h of rumen incubation ($P<0.05$)
232 (Table 2). The order Coriobacteriales varied at each time interval substantially, with no real
233 pattern evident ($P<0.05$) (Table 2). Despite their lower abundances the reads pertaining to
234 orders Aeromonadales, Desulfuromonadales, and Methylophilales also changed significantly
235 over time ($P<0.05$) (Table 2). More sequences pertaining to the order Aeromondales was
236 seen at 1 and 2 h of rumen incubation compared with 6 and 8 h of incubation ($P<0.05$) (Table
237 2). The order Desulfuromonadales varied at each time interval substantially, with no real
238 pattern evident ($P<0.05$) (Table 2). The order Methylophilales decreased significantly after
239 the first 1 h of rumen incubation ($P<0.05$) (Table 2). The remaining bacterial orders were
240 relatively minor and showed no changes in abundance over incubation time ($P>0.05$) (Table
241 2).

242

243 **Family level temporal diversity of the attached microbiome post 1 h of rumen** 244 **incubation**

245 On a family level the most abundant attached bacteria were Lachnospiraceae, Prevotellaceae,
246 Veillonellaceae, Fibrobacteraceae, Ruminococcaceae and Coriobacteriaceae (on average
247 approx. 75, 19, 9, 5, 3 and 2% of total average normalised reads across time points
248 respectively), whereas a further 31 families were relatively minor (<2% of total normalised
249 reads/phyla) in comparison (Table 3). The family Lachnospiraceae changed significantly in
250 abundance over time, with increased abundances present during secondary colonisation
251 events (4-8 h) ($P<0.05$) (Table 3). The family Veillonellaceae changed significantly in
252 abundance over time, with decreased abundances present after 8 h of incubation ($P<0.05$)

253 (Table 3). The families Prevotellaceae and Fibrobacteraceae did not change significantly in
254 abundance over time ($P>0.05$) (Table 3). The family Ruminococcaceae changed significantly
255 in abundance over time, with increased abundances present 8 h after incubation ($P<0.05$)
256 (Table 3). The family Coriobacteriaceae changed significantly in abundance over time, with
257 the highest abundances seen 2 h after incubation ($P<0.05$) (Table 3). Of the other lower
258 abundance families only Methylophilaceae and Succinivibrionaceae changed significantly in
259 abundance over time, with both showing decreased abundance after 1 and 2 h of incubation
260 respectively ($P<0.05$) (Table 3).

261

262 **Genus level temporal diversity of the attached microbiome post 1 h of rumen incubation**

263 On a genus level the most abundant attached bacteria were *Butyrivibrio*, *Pseudobutyrvibrio*,
264 *Selenomonas*, *Prevotella*, *Fibrobacter*, *Olsenella*, and *Ruminococcus* (approx. 44, 17, 12, 10,
265 6, 3 and 2% of total average normalised reads across time points respectively), whereas a
266 further 52 genera were relatively minor ($<2\%$ of total normalised reads/phyla) in comparison
267 (Table 4). The number of sequences pertaining to the genera *Butyrivibrio*, *Prevotella*,
268 *Fibrobacter* and *Olsenella* did not change significantly in abundance over time ($P>0.05$)
269 (Table 4). *Pseudobutyrvibrio* read abundances changed significantly over time, with greater
270 abundances present from 4-8 h of incubation (secondary colonisation phase) ($P<0.05$) (Table
271 4). *Selenomonas* read abundances changed significantly over time, but no decipherable
272 changes in pattern between primary and secondary colonisation could be seen ($P<0.05$)
273 (Table 4). *Ruminococcus* read abundances changed significantly over time, with an increase
274 in abundance evident after 8 h of incubation within the rumen ($P<0.05$) (Table 4). Despite
275 their lower abundances the reads pertaining to genera *Rhodanobacter*, *Roseburia*,
276 *Succinivibrio* and *Murdochiella* also showed temporal variation in abundance ($P<0.05$)

277 (Table 4). *Rhodanobacter* and *Murdochiella* abundance was highest at 2 h post incubation
278 ($P<0.05$) (Table 4), nonetheless even at their highest value they accounted for $>0.1\%$ of the
279 attached diversity. *Roseburia* read abundances were significantly higher in the secondary
280 phase (4-8 h) of rumen incubation ($P<0.05$) (Table 4). Conversely, *Succinivibrio* read
281 abundances were higher during the primary phase (1 & 2 h) of rumen incubation ($P<0.05$)
282 (Table 4). The remaining bacterial genera were relatively minor and showed no changes in
283 abundance over incubation time ($P>0.05$) (Table 4).

284

285 **Temporal niche specialisation of the perennial ryegrass attached microbiota incubated** 286 **within the rumen**

287 Differences were observed in the dynamics of classified OTUs within some of the dominant
288 orders within the transition phase (between 2 and 4 h) of PRG incubation within the rumen.
289 Five OTUs, classified as order Bacteroidales, increased in abundance, whilst 8 decreased in
290 abundance between 2 and 4 h of PRG incubation within the rumen (Table 5). The variability
291 in the proportional representation of the order Bacteroidales also decreased post 2 h of
292 incubation (Figure 4). Conversely, 18 OTUs classified as order Clostridiales increased in
293 abundance, whilst 11 decreased in abundance between 2 and 4 h of incubation within the
294 rumen (Table 5). Again, the variability in the proportional representation of the order
295 Clostridiales also decreased post 2 h of incubation, (Table 5 & Figure 4). Very few changes
296 in OTU representation were apparent for OTUs within any of the other dominant orders
297 (Table 5), nonetheless decreases in proportional variability of reads pertaining to
298 Coriobacterales were seen (Figure 4). Decreases in proportional variability of reads
299 pertaining to the orders Bacteroidales, Clostridiales and Coriobacterales, alongside the
300 decrease in Shannon diversity between primary (1 & 2 h) and secondary colonisation (4-8 h),

301 suggest that the attached microbiota show more niche specialisation during secondary
302 colonisation (Figure 3 & 4). Dry matter (DM) disappearance data showed that a minimal
303 amount (2.8 %) of PRG was degraded within the primary phase (1 & 2 h) of the incubation,
304 unlike the transition (2-4 h) between the two phases where 22.2 % was lost (Figure 5). In the
305 secondary phase between 4 and 8 h of incubation, a further approx. 31.7% of the PRG DM
306 was degraded (Figure 5).

307

308 **DISCUSSION**

309 In this study we characterised the rumen bacteria attached to fresh perennial ryegrass that had
310 been incubated in the rumen over time in order to enhance our understanding of ruminal
311 plant-microbe interactions. Within this study we have demonstrated, using 454 based
312 pyrosequencing of 16S rRNA (cDNA based), that substantial temporal changes occur in the
313 attached microbiota, resulting in primary (1 & 2 h) and secondary (4-8 h) colonisation events
314 by rumen bacteria. The change to a secondary phase was mainly associated with decreases in
315 sequences pertaining to the genera *Succinivibrio* and increases in *Pseudobutyrvibrio*,
316 *Roseburia* and *Ruminococcus*. *Butyrvibrio*, *Fibrobacter*, *Olsenella* and *Prevotella* also
317 dominated the attached microbiome irrespective of incubation time.

318 The depth of sequencing and read length obtained within this study is comparative or
319 higher than those reported in many other published datasets in which 454 technology was
320 used to probe the rumen microbiome. For example Roggenbuck *et al.* (2014) obtained 1,743
321 reads/sample with an average read length of 376 bp, Jami *et al.* (2013) obtained an average of
322 10,938 reads/sample (average read length not specified), Pitta *et al.* (2014) obtained on
323 average 5,199 reads /sample (average read length not specified), Fouts *et al.* (2012) obtained
324 23,493 reads and Jami and Mizrahi (2012) reported an average 9,587 reads/sample with an

325 average read length of 338 bp. In this study we obtained on average 31,761 reads/sample, in
326 the same range as obtained in our previous study (Huws *et al.*, 2014b). Our rarefaction curve
327 based on OTUs, also demonstrated some plateauing. It was suggested in another study (Kim
328 *et al* 2011), that to achieve 99.9% coverage at species level, at least 78,218 bacterial 16S
329 sequences would be needed which equates to approx. 41% of sequences obtained within this
330 study post-filtering. From the reads generated from rumen incubated samples we identified
331 11 phyla, 24 orders, 37 families and 59 genera and an average of 9,093 OTUs, which is
332 similar to that obtained from other previously 454 based rumen microbiome datasets (Fouts *et*
333 *al.*, 2012; Jami and Mizrahi, 2012; Pope *et al.*, 2012; Jami *et al.*, 2013; Huws *et al.*, 2014b).
334 Thus whilst it is possible that our coverage doesn't include all the diversity present, the
335 diversity captured gives a very good indication of the bacterial diversity and temporal
336 changes, post rumen incubation.

337 Our study shows that the plant epiphytic communities rapidly diminished in
338 proportional representation when the rumen microbiota begin to colonise. It should be noted
339 that *Flavobacterium*, *Delftia*, *Cellvibrio* and *Pseudomonas* spp. are still present within the
340 reads obtained post-rumen incubation. This is likely to be because they were the most
341 predominant epiphytes found colonising the PRG pre-incubation.

342 The 16S rRNA sequencing information concurred with our previous DGGE based
343 data showing clear primary (1 & 2 h) and secondary (4-8 h) bacterial colonisation events on
344 fresh perennial ryegrass within the rumen (Huws *et al.*, 2013b; Huws *et al.*, 2014a).
345 Interestingly, a recent publication by Kingston-Smith *et al.* (2013) using FT-IR to investigate
346 the metabolite fingerprint of the interactome (perennial ryegrass coupled with the attached
347 microbiota) did not demonstrate clear differences between 2 and 4 h although a change from
348 6 h onwards was noted. This is probably a consequence of the fact that both the perennial
349 ryegrass and the attached microbiota were analysed together, therefore masking changes

350 occurring in each component separately. A recent DNA based study investigating temporal
351 colonisation of air dried switchgrass showed changes in the microbiome over time, but the
352 greatest changes were observed within the initial 30 mins and after 4 h of rumen incubation
353 (Piao *et al.*, 2014). Nonetheless, previous DGGE analysis of the rumen bacteria attached to
354 fresh PRG, on both a DNA and RNA basis, found no differences within 30 min of incubation
355 (Edwards *et al.*, 2007). Sun *et al.* (2008) found using DGGE that temporal changes in the
356 attached microbiota on Chinese wild rye hay incubated in the rumen occurred between 6 and
357 12 h of incubation. The likely difference between our study and that of others is due to the
358 species of the plant material used, and also our plant material was fresh and not conserved.
359 Furthermore, in this study we also investigated changes on an RNA basis, in order to probe
360 changes in the truly metabolically active bacterial community, whereas other studies used
361 DNA (Piao *et al.*, 2014; Sun *et al.*, 2008). Irrespective of this, however, it is clear from these
362 studies that colonisation events are rapid within the rumen and timings of ecological changes
363 are dependent on the plant characteristics.

364 The data in this study suggest that primary colonising bacteria are likely to utilise
365 soluble nutrients, and that the secondary phase colonisers are adept at degrading plant
366 structural components. This suggestion is based on the fact that only 2.8% plant dry matter
367 disappearance was seen between 1-2 h (primary colonisation phase) of rumen incubation and
368 31.7% dry matter disappearance was seen between 4-8 h (secondary colonisation phase) of
369 rumen incubation. In terms of the temporal changes in the attached microbiota, we observed
370 that *Succinivibrio* (order Aeromonadales) were more abundant during primary colonisation
371 events than secondary colonisation events. The reason for increased abundance of
372 *Succinivibrio* during primary colonisation is unclear as this bacterium is considered to be
373 predominantly amylolytic. Nonetheless the normalised read abundances of *Succinivibrio* are
374 low irrespective of time. Conversely, we observed that *Pseudobutyrvibrio*, *Roseburia* and

375 *Ruminococcus* spp. (all order Clostridiales) were less abundant during primary colonisation
376 events than in secondary colonisation events. Piao *et al.* (2014) also saw increases in
377 *Pseudobutyrvibrio* and *Ruminococcus* spp. during secondary colonisation events, when
378 investigating temporal colonisation of switchgrass incubated within the rumen.
379 *Pseudobutyrvibrio* spp. commonly possess xylanases which randomly cleave the β -1,4
380 backbone of the complex plant cell wall polysaccharide xylan (Krause, 2003). Likewise,
381 *Ruminococcus* spp. are well recognised for their fibrolytic capacity due to the possession of
382 numerous glycosyl hydrolase families (Krause, 2003; Dai *et al.*, 2015). Therefore it is
383 possible that the increase in these bacteria is at least partially responsible for the 31.7% dry
384 matter disappearance seen during the secondary phase of colonisation.

385 Irrespective, of temporal changes it was also noted that *Butyrvibrio*, *Fibrobacter*,
386 *Olsenella* and *Prevotella* spp. read abundances were high irrespective of colonisation phase.
387 Rumen *Butyrvibrio* spp. are known to have proteolytic, biohydrogenating and plant
388 hemicellulolytic activity (Hobson and Stewart, 1997; Krause, 2003). *Fibrobacter* spp. are
389 regarded as being mainly fibrolytic bacteria. Indeed, a recent metatranscriptomic study by
390 Dai *et al.* (2015) suggested that the bulk of ruminal glycosyl hydrolases, including xylanases
391 and endoglucanases, are possessed by *Ruminococcus* and *Fibrobacter* spp. *Olsenella*, on the
392 other hand, is a reasonably newly classified genus (Dewhirst *et al.*, 2001), composed of
393 bacteria that can ferment carbohydrates to lactic acid (Kraatz *et al.*, 2011). A rumen
394 *Olsenella* spp. has also been shown to have β -glucosidase activity, showing its capacity to
395 breakdown glucose (Kraatz *et al.*, 2011). *Prevotella* spp., are mainly known for their starch
396 degrading and proteolytic capacity, but they also have cellulolytic capacity (Gardner *et al.*,
397 1995; Krause, 2003). Due to the fact that most of these attached bacteria have many
398 functions, it is not possible to conclude with absolute certainty what their role is at a given
399 incubation time in terms of plant degradation without gene expression data.

400

401 Nevertheless, variation in proportional representation of Shannon diversity indices
402 and significant OTU changes assigned to order level was observed in this study indicating
403 functional drivers for the succession. It is speculated that the decreased diversity of the
404 secondary colonisers of the orders Clostridiales and Bacteroidales is due to the fact that these
405 bacteria play a more focussed role in plant degradation and nutrient assimilation during the
406 secondary colonisation phase after soluble plant nutrients have been depleted. It has been
407 shown previously that regardless of the concentration, the rate of release of soluble
408 carbohydrate from fresh forage is likely to be limiting to the microbiota (Kingston-Smith *et*
409 *al.*, 2003). This was not apparent in our previous studies (Huws *et al.*, 2013), and is likely to
410 have been due to methodological limitations of the DGGE technique. The study by Piao *et*
411 *al.* (2014) showed increases in Shannon's diversity until 1 h of incubation then a plateau.
412 The reasons for the differences between our findings and those of Piao *et al.* (2014) are
413 unclear but may be due to the different plants analysed, whether they were conserved or not,
414 and also the fact that we analysed the adherent bacterial diversity using RNA as opposed to
415 DNA.

416 In summary, this study demonstrates that fresh perennial ryegrass is rapidly colonised
417 within the rumen with a substantial decrease in active plant epiphytic communities within 1 h
418 of incubation, followed by a biphasic temporal change in the ecology of the adherent
419 bacterial community. These primary (1 & 2 h) and secondary (4-8 h) phases in the attached
420 microbiota were attributable mainly to decreases in *Succinivibrio* spp. and increases in
421 *Pseudobutyrvibrio*, *Roseburia* and *Ruminococcus* spp. during secondary colonisation.
422 Irrespective of temporal changes, the continually high abundances of *Butyrivibrio*,
423 *Fibrobacter*, *Olsenella* and *Prevotella* suggest that they also play a major role in the
424 degradation of the plant. It is clear that a temporal understanding of the functional roles of

425 these microbiota within the rumen is now required to understand the plant-microbe
426 interactome and improve ruminant nutrient use efficiency further. Understanding the plant
427 degradation limitations encountered by the attached microbiota will lead to novel plant
428 breeding targets aimed at increasing the potential degradation of PRG within the rumen and
429 thus increasing animal nutrient use efficiency.

430

431 **SUPPLEMENTARY DATA**

432 Supplementary data is available at FEMSEC online.

433

434 **ACKNOWLEDGEMENTS**

435 This work was supported by the Biotechnology and Biological Sciences Research Council
436 (BBSRC) Institute Strategic Programme Grant, Rumen Systems Biology, (grant number
437 BB/E/W/10964A01) and the Coleg Cymraeg Cenedlaethol.

438

439 **REFERENCES**

440 Brulc JM, Antonopoulos DA, Miller MEB, *et al.* Gene-centric metagenomics of the fiber-
441 adherent bovine rumen microbiome reveals forage specific glycosyl hydrolases. *Proc*
442 *Natl Acad Sci USA* 2009;**106**: 1948-1953.

443 Cheng KJ, Fay JP, Howarth RE, *et al.* Sequence of events in the digestion of fresh legume
444 leaves by rumen bacteria. *Appl Environ Microbiol* 1980;**40**: 613-625.

445 Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun YC, Brown T, Porras-Alfaro A,
446 Kuske CR, & Tiedje JM. Ribosomal Database Project: data and tools for high throughput
447 rRNA analysis. *Nucleic Acids Res* 2014;42:D633-D642.

448 Creevey CJ, Kelly WJ, Henderson G, & Leahy SC. Determining the culturability of the
449 rumen bacterial microbiome. *Microbial biotechnology*, 2014;7(5), 467-479.

450 Dai X, Tian Y, Li J, *et al.* Metatranscriptomic analyses of plant cell wall polysaccharide
451 degradation by microorganisms in the cow rumen. *Appl Environ Microbiol* 2015;**81**:
452 1375-1385.

453 Dewhurst FE, Paster BJ, Tzella N, *et al.* Characterization of the novel human oral isolates and
454 cloned 16s rDNA sequences that fall in the family *Coriobacteriaceae*: description of
455 *Olsenella* gen. nov., reclassification of *Lactobacillus uli* as *Olsenella uli* comb. Nov. and
456 description of *Olsenella profusa* sp. nov. *Int J Syst Evol Microbiol* 2001;51: 1797-1804.

457 Dewhurst RJ, Mitton AM, Offer NW, *et al.* Effect of composition of grass silages on milk
458 production and nitrogen utilisation by dairy cows. *Anim Sci* 1996;**62**: 25- 34.

459 Edwards JE, Huws SA, Kim EJ, *et al.* Characterisation of the dynamics of initial bacterial
460 colonisation of non-conserved (Fresh) forage in the rumen. *FEMS Microbiol Ecol*
461 2007;**62**: 323-335.

462 Edwards JE, Huws SA, Kim EJ, *et al.* Advances in microbial ecosystem concepts and their
463 consequences for ruminant agriculture. *Animal* 2008a;**2**: 653-660.

464 Edwards AH, Kingston-Smith AH, Jiminez HR, *et al.* Dynamics of initial colonisation of
465 non-conserved perennial ryegrass by anaerobic fungi in the bovine rumen. *FEMS*
466 *Microbiol Ecol* 2008b;**66**: 537-545.

467 Foresight. The Future of Food and Farming: Challenges and choices for global sustainability.
468 2011; UK Government report.

469 Fouts DE, Szpakowski S, Purushe J, *et al.* Next generation sequencing to define prokaryotic
470 and fungal diversity in the bovine rumen. *PLOS One* 2012;**7**: e48289.

471 Gardner RG, Wells JE, Russell JB, *et al.* The cellular location of *Prevotella ruminicola* β -
472 1,4-and its occurrence in other strains of ruminal bacteria. *Appl Env Microbiol* 1995;**61**:
473 3288-3292.

474 Huws SA, Kim EJ, Lee MRF, *et al.* As yet uncultured bacteria phylogenetically classified as
475 *Prevotella*, *Lachnospiraceae* incertae sedis, and unclassified *Bacteroidales*, *Clostridiales*
476 and *Ruminococcaceae* may play a predominant role in ruminal biohydrogenation.
477 *Environ Microbiol* 2011;**13**: 1500-1512.

478 Huws SA, Scott MB, Tweed JKS, *et al.* Fatty acid oxidation products ('green odour')
479 released from perennial ryegrass following biotic and abiotic stress, potentially have
480 antimicrobial properties against the rumen microbiota resulting in decreased
481 biohydrogenation. *J Appl Microbiol* 2013a;**15**: 1081-1090.

482 Huws SA, Maorga OL, Theodorou MK, *et al.* Successional colonisation of perennial ryegrass
483 by rumen bacteria. *Letters Appl Microbiol* 2013b;**56**:186-196.

484 Huws SA, Mayorga OL, Theodorou MK, *et al.* Differential colonization of plant parts by the
485 rumen microbiota is likely to be due to different forage chemistries. *J Microb Biochem*
486 *Technol* 2014a;**6**: 080-086.

487 Huws SA, Kim EJ, Cameron SJS, *et al.* Characterization of the rumen lipidome and
488 microbiome of steers fed a diet supplemented with flax and echium oil. *Microb Biotech.*
489 2014b; DOI: 10.1111/1751-7915.12164.

490 Jami E, Mizrahi I. Composition and similarity of bovine rumen microbiota across individual
491 animals. *PLOS One* 2012;**7**: 333306.

492 Jami E, Israel A, Kotser A *et al.* Exploring the bovine rumen bacterial community from birth
493 to adulthood. *ISME J* 2013;**7**: 1069-1079.

494 Kav A-Y, Sasson G, Jami E, *et al.* Insights into the bovine rumen plasmidome. *Proc Natl*
495 *Acad Sci USA* 2012;**109**:5452-5457.

496 Kim EJ, Huws SA, Lee MRF, *et al.* Dietary transformation of lipid in the rumen microbial
497 ecosystem. *Asian-Aus J Anim Sci* 2009;**22**: 1341-1350.

498 Kim M, Morrison M, Yu Z. Status of the phylogenetic diversity census of ruminal
499 microbiomes. *FEMS Microbiol Ecol* 2011;**76**: 49–63.

500 Kingston-Smith AH, Bollard AL, Thomas BJ, *et al.* Nutrient availability during the early
501 stages of colonization of fresh forage by rumen micro-organisms. *New Phytol* 2003;
502 **158**: 119-130.

503 Kingston-Smith AH, Davies TE, Edwards JE, *et al.* From plants to animals; the role of plant
504 cell death in ruminant herbivores. *J Experiment Bot* 2008;**59**: 521-532.

505 Kingston-Smith AH, Edwards JE, Huws SA, *et al.* Plant-based strategies towards minimising
506 livestock's shadow. *Proc Nut Soc* 2010;**4**: 1-8.

507 Kingston-Smith AH, Davies TE, Rees Stevens P, *et al.* Comparative metabolite
508 fingerprinting of the rumen system during colonisation of three forage grass (*Lolium*
509 *perenne* L.) varieties. *PLOS One* 2013;**8**:e82801.

510 Koike S, Pan J, Kobayashi Y, *et al.* Kinetics of *in sacco* fibre-attachment of representative
511 ruminal cellulolytic bacteria monitored by competitive PCR. *J Dairy Sci* 2003;**86**:1429-
512 1435.

513 Krause, DO, Denma SE, Mackie RI, *et al.* Opportunities to improve fiber degradation in the
514 rumen: microbiology, ecology, and genomics. *FEMS Microbiol Rev* 2003;**27**: 663-693.

515 Leng RA. Interactions between microbial consortia in biofilms: a paradigm shift in rumen
516 microbial ecology and enteric methane mitigation. *Anim Prod Sci* 2014;**54**: 519-543.

517 Li W, Fu L, Niu B, *et al.* Ultrafast clustering algorithms for metagenomics sequence analysis.
518 *Briefings Bioinformatics* 2012;**13**: 656-668.

519 Love MI, Huber W, & Anders S. Moderated estimation of fold change and dispersion for
520 RNA-seq data with DESeq2. *Genome Biol* 2014;**15**(12), 550.

521 Mackie RI. Mutualistic fermentative digestion in the gastrointestinal tract: diversity and
522 evolution 1. *Integr Comp Biol* 2002;**42**: 319-326.

523 MacRae JC, Ulyatt MJ. Quantitative digestion of fresh herbage by sheep. 2. Sites of some
524 nitrogenous constituents. *J Agri Sci* 1974;**82**: 309-319.

525 McAllister TA, Bae HD, Jones GA, *et al* Microbial attachment and feed digestion in the
526 rumen. *J Anim Sci* 1994;**72**:3004-3018.

527 Miron J, Ben-Ghedalia D, Morrison M. Invited review: Adhesion mechanisms of rumen
528 cellulolytic bacteria. *J Dairy Sci* 2001;**84**:1294-1309.

529 Ørskov ER, DeB Hovell FD, Mould F. The use of the nylon bag technique for the evaluation
530 of feedstuffs. *Trop Anim Pro* 1980;**5**: 195-213.

531 Ougham HJ, Davies TGE. Leaf development in *Lolium temulentum*-gradients of RNA
532 complement and plastid and nonplastid transcripts. *Phys Plant* 1990;**2**:331-338.

533 their alteration in response to variation in the diet. *Gene* 2014b;**545**: 88-94.

534 Payne, R.W, Murray, D.A., Harding, S.A., Baird, D.B. and Soutar, D.M. (2007) GenStat®
535 for Windows™ 9th Edition, Introduction. Hemel Hempstead, UK: VSN International.

536 Piao H, Lachman M, Malfatti S, *et al.* Temporal dynamics of fibrolytic and methanogenic
537 rumen microorganisms during in situ incubation of switchgrass determined by 16S
538 rRNA gene profiling. *Frontiers Microbiol* 2014;**5**:article 307.

539 Pitta DW, Parmar N, Patel AK, *et al.* (2014) Bacterial diversity dynamics associated with
540 different diets and different primer pairs in the rumen of Kankrej cattle. *PLOS One*
541 2014;**9**:e111710.

542 Roggenbuck M, Sauer C, Poulsen M, *et al.* The giraffe (*Giraffa Camelopardalis*) rumen
543 microbiome. *FEMS Microbiol Ecol* 2014;**90**:237-246.

544 Russell JB, Rychlik, JL. Factors that alter rumen microbial ecology. *Science* 2001;**292**:1119-
545 1122.

546 Sun YZ, Mao SY, Yao W, *et al.* DGGE and 16S rDNA analysis reveals a highly diverse and
547 rapidly colonising bacterial community on different substrates in the rumen of goats.
548 *Animal* 2008;**2**:391-398.

549 Vanzant ES, Cochran RC, Titgemeyer EC. Standardization of *in situ* techniques for ruminant
550 feedstuff evaluation. *J Anim Sci* 1998;**76**: 2717-2729.

551

552

553

FIGURE LEGENDS




Figure 1. Heat map illustrating changes in proportional read abundances of perennial ryegrass epiphytic communities post rumen incubation. Mean data for each time point are shown (n=2 for 0h and n=6 for all other time points).




Figure 2. PCA plot showing the diversity of rumen bacteria attached to perennial ryegrass over time. Data for post rumen incubation of 1, 2, 4, 6 and 8 h (T1, T2, T4, T6 and T8 respectively) within each cow (C1, C2 and C3 respectively) are shown. Mean data for 2 bags incubated within each cow are shown for each time point.




Figure 3. Boxplots of the average Shannon diversity indices at each time point representing each of the three cows sampled in duplicate (n=6). Time points that do not share notations were significantly different ($P<0.05$) according to a t-test. Post rumen incubation time points of 1, 2, 4, 6 and 8 h (T1, T2, T4, T6 and T8 respectively).




Figure 4. Proportional changes in the 6 most abundant bacterial orders attached to perennial ryegrass incubated in the rumen over time (n=6). Post rumen incubation time points of 1, 2, 4, 6 and 8 h (T1, T2, T4, T6 and T8 respectively).




Figure 5. Perennial ryegrass dry matter disappearance (%) following incubation within the rumen over time. Standard error of the mean for each time point are shown.

Table 1. Comparison of the bacterial phyla attached to perennial ryegrass over time within the rumen. Data shown are the average values (n=6) of the normalised reads.

Phylum	Time of incubation (h)					SED	P
	1	2	4	6	8		
<i>Actinobacteria</i>	368.2 ^a	792.6 ^b	146.8 ^a	132.0 ^a	165.6 ^a	162.8	0.016
<i>Bacteroidetes</i>	3356	3954	2341	2528	2726	584.6	NS
<i>Elusimicrobia</i>	11.5 ^b	3.7 ^a	1.3 ^a	0.4 ^a	0.3 ^a	2.6	0.012
<i>Fibrobacteres</i>	675.5	703.2	511.5	744.8	643.4	173.6	NS
<i>Firmicutes</i>	12570 ^{ab}	11840 ^a	14600 ^c	14151 ^{bc}	14136 ^{bc}	819.0	0.040
<i>Fusobacteria</i>	0.9	0.6	0.6	0.3	0.0	0.8	NS
<i>Lentisphaerae</i>	6.2 ^c	5.0 ^{bc}	1.3 ^{ab}	1.8 ^{ab}	0.3 ^a	1.7	0.029
<i>Proteobacteria</i>	515.5	349.9	98.6	160.4	48.7	178.5	NS
<i>Spirochaetes</i>	283.7	122.9	114.7	96.5	95.2	70.9	NS
<i>Tenericutes</i>	0.3	0.6	1.0	1.0	1.6	1.1	NS
<i>Verrucomicrobia</i>	33.0 ^{ab}	49.3 ^b	7.6 ^a	8.9 ^a	7.7 ^a	10.9	0.015

Values with different superscripts on the same row differed significantly ($P < 0.05$), whereas values that were not significantly (NS) different ($P > 0.05$) have no superscripts in the same row.

Table 2. Comparison of the bacterial orders attached to perennial ryegrass over time within the rumen. Data shown are the average values (n=6) of the normalised reads.

Order	Time of incubation (h)					SED	P
	1	2	4	6	8		
<i>Actinomycetales</i>	11.9	27.9	0.6	0.4	0.0	15.0	NS
<i>Aeromonadales</i>	57.4 ^b	64.3 ^b	39.4 ^{ab}	19.3 ^a	18.8 ^a	11.8	0.013
<i>Anaeroplasmatales</i>	0.3	0.6	1.0	1.0	1.6	1.1	NS
<i>Bacilliales</i>	7.8	2.2	0.0	3.5	0.0	3.5	NS
<i>Bacteroidales</i>	3135	3737	2285	2457	2693	546.1	NS
<i>Burkholderiales</i>	18.7	21.6	13.0	14.2	15.3	3.1	NS
<i>Caulobacterales</i>	6.0	7.0	0.0	3.1	0.7	2.9	NS
<i>Clostridiales</i>	10705 ^a	9871 ^a	12625 ^b	12660 ^b	12943 ^b	749.2	0.011
<i>Coriobacteriales</i>	356.3 ^a	764.4 ^b	145.6 ^a	129.5 ^b	165.0 ^a	152.7	0.014
<i>Desulfuromonadales</i>	0.0 ^a	1.9 ^c	0.3 ^{ab}	1.1 ^{bc}	0.7 ^{ab}	0.4	0.011
<i>Enterobacteriales</i>	2.3	5.4	0.3	0.7	0.0	2.7	NS
<i>Fibrobacterales</i>	675.2	702.9	510.9	744.1	643.4	173.8	NS
<i>Flavobacteriales</i>	50.4	24.5	0.6	17.5	0.0	22.3	NS
<i>Lactobacillales</i>	1.2	25.0	2.6	5.3	4.7	13.5	NS
<i>Methylophilales</i>	3.0 ^b	0.9 ^a	0.0 ^a	0.0 ^a	0.0 ^a	0.7	0.012
<i>Neisseriales</i>	0.9	27.9	0.0	2.1	0.0	14.6	NS
<i>Pseudomonadales</i>	51.4	20.4	0.9	16.4	0.0	25.1	NS
<i>Rhizobiales</i>	17.5	3.5	0.6	3.5	0.3	9.2	NS
<i>Rhodobacterales</i>	0.3	1.6	0.0	0.0	0.0	0.5	NS
<i>Rhodocyclales</i>	1.2	0.0	0.0	0.0	0.0	0.7	NS
<i>Selenomonadales</i>	1357 ^{ab}	1455 ^b	1663 ^b	1135 ^{ab}	764 ^a	251.2	0.054
<i>Sphingobacteriales</i>	27.2	6.6	0.0	8.4	0.0	0.2	NS
<i>Spirochaetales</i>	222.5	84.6	81.9	70.7	76.8	76.2	NS
<i>Xanthomonadales</i>	14.3	8.4	0.0	4.9	0.3	5.5	NS

Values with different superscripts on the same row differed significantly ($P < 0.05$), whereas values that were not significantly (NS) different ($P > 0.05$) have no superscripts in the same row.

Table 3. Comparison of the bacterial families attached to perennial ryegrass over time within the rumen. Data shown are the average values (n=6) of the normalised reads.

Family	Time of incubation (h)					SED	P
	1	2	4	6	8		
<i>Alcaligenaceae</i>	10.4	4.5	0.3	1.4	0.3	4.8	NS
<i>Anaeroplasmataceae</i>	0.3	0.6	1.0	1.1	1.6	1.1	NS
<i>Beijerinckiaceae</i>	0.3	0.0	0.0	0.0	0.0	0.2	NS
<i>Bradyrhizobiaceae</i>	0.0	0.0	0.6	0.0	0.0	0.4	NS
<i>Burkholderiaceae</i>	0.89	0.0	0.0	0.0	0.0	0.6	NS
<i>Caulobacteraceae</i>	6.0	7.0	0.0	3.1	0.7	2.9	NS
<i>Comamonadaceae</i>	21.1	8.7	0.2	5.1	0.3	9.8	NS
<i>Coriobacteriaceae</i>	356.3 ^a	764.4 ^b	145.6 ^a	129.5 ^a	165.0 ^a	152.7	0.014
<i>Cryomorphaceae</i>	1.2	0.3	0.0	0.0	0.0	0.7	NS
<i>Cytophagaceae</i>	0.3	0.0	0.0	0.0	0.0	0.2	NS
<i>Enterobacteriaceae</i>	0.8	1.8	0.1	0.2	0.0	0.9	NS
<i>Eubacteriaceae</i>	5.9	4.3	4.5	2.9	2.3	2.9	NS
<i>Fibrobacteraceae</i>	675.2	702.9	510.9	744.1	643.4	173.8	NS
<i>Flavobacteriaceae</i>	49.2	24.2	0.6	17.5	0.0	21.7	NS
<i>Hyphomicrobiaceae</i>	3.8	0.0	0.0	0.7	0.3	2.3	NS
<i>Lachnospiraceae</i>	9506 ^a	8631 ^a	11701 ^b	11569 ^b	11458 ^b	814.6	0.016
<i>Methylobacteriaceae</i>	0.0	0.2	0.0	0.0	0.0	0.1	NS
<i>Methylophilaceae</i>	3.1 ^a	0.9 ^b	0.0 ^b	0.0 ^b	0.0 ^b	0.0	0.012
<i>Moraxellaceae</i>	0.5	1.8	0.0	0.0	0.0	1.1	NS
<i>Neisseriaceae</i>	0.9	27.9	0.0	2.1	0.0	14.6	NS
<i>Oxalobacteraceae</i>	18.6	13.5	0.9	0.7	0.4	7.6	NS
<i>Paenibacillaceae</i>	1.6	0.4	0.0	0.7	0.0	0.7	NS
<i>Porphyromonadaceae</i>	9.7	8.0	18.4	16.5	18.1	9.0	NS
<i>Prevotellaceae</i>	2632	3170	1816	1925	2012	550.2	NS
<i>Pseudomonadaceae</i>	50.4	16.8	0.9	16.4	0.0	25.0	NS
<i>Rhizobiaceae</i>	12.9	3.2	0.0	2.8	0.0	6.3	NS
<i>Rhodobacteraceae</i>	0.3	1.6	0.0	0.0	0.0	0.5	NS
<i>Rhodocyclaceae</i>	1.2	0.0	0.0	0.0	0.0	0.7	NS
<i>Ruminococcaceae</i>	445.2 ^a	437.5 ^a	317.0 ^a	400.8 ^a	660.4 ^b	83.0	0.031
<i>Sinobacteraceae</i>	0.3	0.0	0.0	0.0	0.0	0.2	NS
<i>Sphingobacteriaceae</i>	26.9	6.6	0.0	8.4	0.0	15.2	NS
<i>Spirochaetaceae</i>	11.1	4.2	4.1	3.5	3.8	3.8	NS
<i>Streptococcaceae</i>	1.2	25.0	2.6	5.3	4.7	13.5	NS
<i>Succinivibrionaceae</i>	57.0 ^a	62.1 ^a	39.4 ^{ab}	19.3 ^b	18.8 ^b	11.8	0.016
<i>Sutterellaceae</i>	29.2	39.6	40.3	31.0	8.1	11.5	NS
<i>Veillonellaceae</i>	1357 ^{ab}	1455 ^b	1663 ^b	1135 ^{ab}	764 ^a	251.2	0.054
<i>Xanthomonadaceae</i>	14.00	8.54	0.00	4.9	0.33	5.34	NS

Values with different superscripts on the same row differed significantly ($P < 0.05$), whereas values that were not significantly (NS) different ($P > 0.05$) have no superscripts in the same row.

Table 4. Comparison of the bacterial genera attached to perennial ryegrass over time within the rumen. Data shown are the average values (n=6) of the normalised reads.

Genus	Time of incubation (h)					SED	P
	1	2	4	6	8		
<i>Acidovorax</i>	27.9	12.2	1.2	3.5	0.3	12.8	NS
<i>Acinetobacter</i>	0.5	1.8	0.0	0.0	0.0	1.1	NS
<i>Advenella</i>	0.3	1.3	0.0	0.0	0.0	0.8	NS
<i>Anaeroplasma</i>	0.3	0.6	1.0	1.0	1.6	1.1	NS
<i>Anaerovibrio</i>	25.4	21.0	16.0	14.9	9.9	7.0	NS
<i>Asticcacaulis</i>	0.0	0.3	0.0	0.4	0.0	0.3	NS
<i>Blautia</i>	1.7	2.5	0.3	0.0	2.0	1.2	NS
<i>Bosea</i>	0.0	0.0	0.6	0.0	0.0	0.4	NS
<i>Brevundimonas</i>	5.1	6.0	0.0	2.8	0.3	2.8	NS
<i>Butyrivibrio</i>	4918	4110	5115	4573	3896	474.3	NS
<i>Camelimonas</i>	0.29 ^a	0.0 ^a	0.0 ^a	0.0 ^a	0.0 ^a	10.9	0.015
<i>Caulobacter</i>	0.9	0.6	0.0	0.0	0.3	0.7	NS
<i>Cellvibrio</i>	18.5	3.8	0.0	3.8	0.0	10.0	NS
<i>Chryseobacterium</i>	24.1	6.3	0.0	4.2	0.3	12.6	NS
<i>Clostridium</i>	114.8	118.8	47.7	45.4	37.9	30.7	NS
<i>Comamonas</i>	38.1	16.9	0.0	14.3	0.0	19.6	NS
<i>Coprococcus</i>	0.0	1.2	0.9	0.7	0.7	0.9	NS
<i>Delftia</i>	148.3	46.8	1.6	36.0	1.0	73.5	NS
<i>Devosia</i>	3.8	0.0	0.0	0.7	0.3	2.3	NS
<i>Duganella</i>	2.5	0.9	0.0	0.0	0.0	1.2	NS
<i>Dyadobacter</i>	0.3	0.0	0.0	0.0	0.0	0.2	NS
<i>Epilithonimonas</i>	0.6	1.6	0.0	0.3	0.0	1.0	NS
<i>Erwinia</i>	0.9	4.2	0.0	0.3	0.0	2.2	NS
<i>Eubacterium</i>	5.9	4.3	4.4	2.9	2.3	2.9	NS
<i>Fibrobacter</i>	675.2	702.9	510.9	744.1	643.4	173.8	NS
<i>Flavobacterium</i>	39.8	17.9	0.6	12.6	0.0	17.3	NS
<i>Helococcus</i>	0.29	1.6	0.0	0.0	0.0	0.7	NS
<i>Herbaspirillum</i>	12.4	8.5	0.9	0.7	0.3	5.5	NS
<i>Howardella</i>	0.9	2.8	0.7	0.0	0.3	0.8	NS
<i>Lachnobacterium</i>	16.3	10.0	61.1	43.9	37.8	22.0	NS
<i>Luteibacter</i>	0.6	1.5	0.0	0.7	0.0	1.1	NS
<i>Methylobacterium</i>	0.0	0.3	0.0	0.0	0.0	0.2	NS
<i>Mogibacterium</i>	2.6	3.1	0.6	0.0	0.7	1.9	NS
<i>Mucilaginibacter</i>	0.6	0.0	0.0	0.0	0.0	0.4	NS
<i>Murdochiella</i>	1.6 ^a	8.0 ^b	1.0 ^a	0.0 ^a	0.3 ^a	4.1	0.014
<i>Olsenella</i>	356.3	765.7	146.5	130.2	165.7	152.2	NS
<i>Oxalicibacterium</i>	0.9	0.0	0.0	0.0	0.0	0.6	NS
<i>Paenibacillus</i>	6.3	1.9	0.0	3.5	0.0	3.4	NS
<i>Pandoraea</i>	0.9	0.0	0.0	0.0	0.0	0.6	NS
<i>Pantoea</i>	0.0	0.6	0.0	0.0	0.0	0.4	NS
<i>Pelomonas</i>	4.2	7.6	0.0	1.4	0.7	2.7	NS
<i>Prevotella</i>	1144	1341	840	939	865	271.3	NS
<i>Propionibacterium</i>	3.5	1.9	0.3	0.4	0.0	1.3	NS
<i>Pseudobutyrvibrio</i>	701 ^a	805 ^a	2500 ^b	2262 ^b	2306 ^b	275.6	<0.001
<i>Pseudomonas</i>	31.9	13.0	0.9	12.6	0.0	15.3	NS

<i>Rheinheimera</i>	3.8	0.0	0.0	0.0	0.0	2.4	NS
<i>Rhizobium</i>	12.9	3.2	0.0	2.8	0.0	6.3	NS
<i>Rhodanobacter</i>	0.0 ^a	1.3 ^b	0.0 ^a	0.0 ^a	0.0 ^a	0.4	0.046
<i>Roseburia</i>	2.0 ^a	0.9 ^a	9.5 ^b	11.8 ^b	10.4 ^b	2.7	0.010
<i>Ruminococcus</i>	151.4 ^a	155.9 ^a	140.5 ^a	204.1 ^{ab}	337.4 ^b	59.0	0.050
<i>Saccharofermentans</i>	5.5	5.6	2.6	1.1	7.0	2.0	NS
<i>Selenomonas</i>	1325 ^{ab}	1426 ^b	1641 ^b	1117 ^{ab}	747 ^a	244.7	0.050
<i>Shinella</i>	1.17	0.0	0.0	0.0	0.0	0.7	NS
<i>Sphingobacterium</i>	23.2	4.7	0.0	8.4	0.0	13.4	NS
<i>Streptococcus</i>	1.2	25.0	2.6	5.3	4.7	13.47	NS
<i>Succinomonas</i>	0.4	2.2	0.0	0.0	0.0	0.9	NS
<i>Succinivibrio</i>	57.0 ^b	62.1 ^b	39.4 ^{ab}	19.3 ^a	18.8 ^a	11.8	0.016
<i>Treponema</i>	275.5	169.8	121.0	89.6	94.6	76.7	NS
<i>Variovorax</i>	19.8	13.6	0.0	1.7	0.3	7.2	NS

Values with different superscripts on the same row differed significantly ($P < 0.05$), whereas values that were not significantly (NS) different ($P > 0.05$) have no superscripts in the same row.

Table 5. Comparison of OTU changes within the dominant orders of bacteria attached to perennial ryegrass over time

Order	T1 to T2			T2 to T4			T4 to T6			T6 to T8		
	Number of OTUs increased	Number of OTUs decreased	Average overall change in abundance	Number of OTUs increased	Number of OTUs decreased	Average overall change in abundance	Number of OTUs increased	Number of OTUs decreased	Average overall change in abundance	Number of OTUs increased	Number of OTUs decreased	Average overall change in abundance
<i>Bacteroidales</i>	-	-	-	5	8	-321	-	-	-	-	1	-73
<i>Clostridiales</i>	-	-	-	18	11	2746	-	-	-	1	4	-55
<i>Coriobacteriales</i>	-	-	-	-	6	-601	-	-	-	-	-	-
<i>Fibrobacterales</i>	-	-	-	-	-	-	-	-	-	-	-	-
<i>Seimonadales</i>	-	-	-	-	-	-	-	-	-	-	-	-
<i>Spirochaetales</i>	-	-	-	1	-	15	-	-	-	-	1	-24
Total	0	0		24	25	1824	0	0	0	1	6	-152

T= Time (h)

Downloaded from <http://fems.oxfordjournals.org/> by guest on February 11, 2016