

## Aberystwyth University

### *Draft Genome Sequence of Probiotic Enterococcus faecium Strain L-3*

Karaseva, Alena; Tsapieva, Anna; Pachebat, Justin Alexander; Suvorov, Alexander

*Published in:*

Genome Announcements

*DOI:*

[10.1128/genomeA.01622-15](https://doi.org/10.1128/genomeA.01622-15)

*Publication date:*

2016

*Citation for published version (APA):*

Karaseva, A., Tsapieva, A., Pachebat, J. A., & Suvorov, A. (2016). Draft Genome Sequence of Probiotic Enterococcus faecium Strain L-3. *Genome Announcements*, 4(1), Article e01599-15.  
<https://doi.org/10.1128/genomeA.01622-15>

#### Document License

CC BY

#### General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

# Draft Genome Sequence of Probiotic *Enterococcus faecium* Strain L-3

Alena Karaseva,<sup>a</sup> Anna Tsapieva,<sup>a</sup> Justin Pachebat,<sup>b</sup> Alexander Suvorov<sup>a,c</sup>

FSBSI Institute of Experimental Medicine, St. Petersburg, Russian Federation<sup>a</sup>; Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Aberystwyth, Ceredigion, United Kingdom<sup>b</sup>; St. Petersburg State University, St. Petersburg, Russian Federation<sup>c</sup>

**We report here the draft genome sequence of the bacteriocin producer *Enterococcus faecium* strain L-3, isolated from a probiotic preparation, Laminolact, which is widely used in the Russian Federation. The draft genome sequence is composed of 74 contigs for a total of 2,643,001 bp, with 2,646 coding genes. Five clusters for bacteriocin production were found.**

Received 2 December 2015 Accepted 7 December 2015 Published 28 January 2016

Citation Karaseva A, Tsapieva A, Pachebat J, Suvorov A. 2016. Draft genome sequence of probiotic *Enterococcus faecium* strain L-3. *Genome Announc* 4(1):e01622-15. doi:10.1128/genomeA.01622-15.

Copyright © 2016 Karaseva et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Alena Karaseva, [xathaelru@gmail.com](mailto:xathaelru@gmail.com).

**E**nterococci are Gram-positive lactic acid bacteria that inhabit the gastrointestinal tracts of diverse hosts (1–3). On one hand, the species *Enterococcus faecium* was considered a harmless commensal of the mammalian gastrointestinal (GI) tract and is commonly used as a probiotic (4) and in food products (5); on another hand, some enterococcal strains were associated with infection, especially in immunocompromised patients (6, 7). *E. faecium* strain L-3 was isolated from a probiotic preparation. This strain has a wide range of antimicrobial activities against organisms, including *Listeria monocytogenes*, *Streptococcus pyogenes*, and *Streptococcus agalactiae* (8).

Genomic DNA was extracted from the cultured bacterium by the DNeasy blood and tissue kit (Qiagen, USA). The genome was sequenced *de novo* using 454 pyrosequencing on the GS FLX platform (Roche 454 Life Sciences). Barcoded DNA libraries were prepared using the GS DNA library preparation kit in combination with GS multiplex identifiers kits. The GS emulsion-based clonal amplification (emPCR) kit II was used for the Multiplex Identifier-adapted library. GS LR70 sequencing kit in combination with GS FLX PicoTiterPlate kit (70 × 75 bp) was used to obtain the sequences. All procedures were carried out according to the manufacturer's instructions (Roche). Reads were *in silico* assembled using the Newbler assembler, giving 74 contigs of different sizes (smallest contig, 102 bp; largest contig, 211,071 bp). Genome annotation was performed using the standard operation procedures from the Rapid Annotations using Subsystems Technology (RAST) server (9).

The draft genome sequence consists of 2,643,001 bases, with a mean of G+C content of 38%. A total of 2,646 coding sequences (CDSs) and 60 structural RNAs were predicted. There were 331 RAST subsystems determined in the genome, which represents only 50% of the sequences assigned.

Several CDSs for the production of bacteriocins, namely, enterocin A, enterocin B, two-component enterocin X (X- $\alpha$  and X- $\beta$  subunits), and lactobin-like bacteriocins, were found.

Seventeen contigs (total size of 115,433 bp) were recognized as a part of a plasmid related to the rep17 family (10).

The sequence was analyzed for the presence of genes that are

considered to be virulence markers employing PCR. The genes for gelatinase (*gelE*), serine protease (*sprE*), quorum-sensing regulator (*fsrB*), putative adherence factors (*asa1*, *efaA*, and *esp*), cytolytins (*cylA*, *cylB*, *cylM*, and *cylL*), hemolysin (*hyl*), and genes related to vancomycin resistance (*vanA*) were not found in this strain (11). The sequences of these genes have not been determined in any contigs under study.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [JRGX000000000](https://www.ncbi.nlm.nih.gov/nuccore/JRGX000000000). The version described in this paper is version [JRGX010000000](https://www.ncbi.nlm.nih.gov/nuccore/JRGX010000000).

## ACKNOWLEDGMENT

This work was supported by RFBR (grant 13-04-01861 to A. Suvorov). The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

## FUNDING INFORMATION

Russian Foundation for Basic Research (RFBR) provided funding to Alexander Suvorov under grant number 13-04-01861.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

1. Franz CMAP, Holzapfel WH, Stiles ME. 1999. Enterococci at the crossroads of food safety? *Int J Food Microbiol* 47:1–24. [http://dx.doi.org/10.1016/S0168-1605\(99\)00007-0](http://dx.doi.org/10.1016/S0168-1605(99)00007-0).
2. Giraffa G. 2002. Enterococci from foods. *FEMS Microbiol Rev* 26:163–171. <http://dx.doi.org/10.1111/j.1574-6976.2002.tb00608.x>.
3. Murray BE. 1990. The life and times of the *Enterococcus*. *Clin Microbiol Rev* 3:46–65.
4. Lund B, Edlund C. 2001. Probiotic *Enterococcus faecium* strain is a possible recipient of the *vanA* gene cluster. *Clin Infect Dis* 32:1384–1385. <http://dx.doi.org/10.1086/319994>.
5. Foulquié Moreno MR, Sarantinopoulos P, Tsakalidou E, De Vuyst L. 2006. The role and application of enterococci in food and health. *Int J Food Microbiol* 106:1–24. <http://dx.doi.org/10.1016/j.jiffoodmicro.2005.06.026>.
6. Moellering RC, Jr. 1992. Emergence of *Enterococcus* as a significant pathogen. *Clin Infect Dis* 14:1173–1176. <http://dx.doi.org/10.1093/clinids/14.6.1173>.
7. Mundy LM, Sahn DF, Gilmore M. 2000. Relationships between entero-

- coccal virulence and antimicrobial resistance. *Clin Microbiol Rev* 13: 513–522. <http://dx.doi.org/10.1128/CMR.13.4.513-522.2000>.
8. Yermolenko E, Chernysh A, Kolobov A, Suvorov A. 2011. Influence of synthetic peptide inducers on antibacterial activity of enterococci. *Benef Microbes* 2:9–13. <http://dx.doi.org/10.3920/BM2010.0006>.
  9. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
  10. Jensen LB, Garcia-Migura L, Valenzuela AJS, Løhr M, Hasman H, Aarestrup FM. 2010. A classification system for plasmids from enterococci and other Gram-positive bacteria. *J Microbiol Methods* 80:25–43. <http://dx.doi.org/10.1016/j.mimet.2009.10.012>.
  11. Bondarenko V, Suvorov A, Vershinin A, Yermolenko E, Kolodzhieva V, Grabovskaya K, Klimovitsch B. 2008. Differences in the pattern of the virulence genes among the industrial and clinical strains. *Biopreparations* 4:3–6. (In Russian).