

Aberystwyth University

The paradox of invasion

Freeman, M. S.; Beatty, G. E.; Dick, J. T. A.; Reid, N.; Provan, J.

Published in:

Journal of Zoology

DOI:

[10.1111/jzo.12283](https://doi.org/10.1111/jzo.12283)

Publication date:

2016

Citation for published version (APA):

Freeman, M. S., Beatty, G. E., Dick, J. T. A., Reid, N., & Provan, J. (2016). The paradox of invasion: Reeves' muntjac deer invade the British Isles from a limited number of founding females. *Journal of Zoology*, 298(1), 54-63. <https://doi.org/10.1111/jzo.12283>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

The paradox of invasion: Reeves' muntjac deer invade the British Isles from a limited number of founding females

Marianne S. Freeman^{1,2}, Gemma E. Beatty^{1,2,3}, Jaimie T. A. Dick^{1,2,3},
Neil Reid^{1,2,3} & Jim Provan^{1,2,3*}

¹ School of Biological Sciences, Queen's University Belfast, Belfast BT9 7BL

² *Quercus*, School of Biological Sciences, Queen's University Belfast, Belfast BT9 7BL

³ Institute for Global Food Security, Queen's University Belfast, Belfast BT9 7BL

Running title: Population genetics of invasive muntjac deer

Correspondence: Dr. Jim Provan (address as above)

Tel: +44 28 9097 2280

Fax: +44 28 9097 5877

E-mail: J.Provan@qub.ac.uk

1 **Abstract**

2

3 High levels of genetic diversity and high propagule pressure are favoured by conservation
4 biologists as the basis for successful re-introductions and ensuring the persistence of
5 populations. However, invasion ecologists recognise the “paradox of invasion”, as successful
6 species introductions may often be characterised by limited numbers of individuals and
7 associated genetic bottlenecks. In the present study, we used a combination of high-
8 resolution nuclear and mitochondrial genetic markers to investigate the invasion history of
9 Reeves’ muntjac deer in the British Isles. This invasion has caused severe economic and
10 ecological damage, with secondary spread currently a concern throughout Europe and
11 potentially globally. Microsatellite analysis based on eight loci grouped all 176 introduced
12 individuals studied from across the species’ range in the UK into one genetic cluster, and
13 seven mitochondrial D-loop haplotypes were recovered, two of which were present at very
14 low frequency and were related to more common haplotypes. Our results indicate that the
15 entire invasion can be traced to a single founding event involving a low number of females.
16 These findings highlight the fact that even small releases of species may, if ignored, result in
17 irreversible and costly invasion, regardless of initial genetic diversity or continual genetic
18 influx.

19 **Introduction**

20

21 Conservation biologists have long appreciated the importance of high levels of genetic
22 diversity and high propagule pressure as the basis for the successful introduction and
23 enhanced persistence of populations (Frankham et al. 2002). The parallel case of species
24 invasions, however, is often characterised by limited numbers of individuals and associated
25 genetic bottlenecks, a so-called “paradox of invasion” (Sax & Brown 2000; Dlugosch &
26 Parker 2008a). Although there are several well-known cases of invasive species that exhibit
27 extremely low levels of genetic variation (Hollingsworth & Bailey 2000), many successful
28 invasions have been facilitated by multiple introductions, resulting in high levels of genetic
29 diversity in the invasive range (Lockwood et al. 2005, 2009; Lavergne & Molofsky 2007;
30 Roman & Darling 2007; Simberloff 2009). In particular, we need to know if particularly
31 damaging species are likely to establish and spread from small founder events or if high
32 propagule pressure is required (Sax & Brown 2000; Lockwood et al. 2005, 2009; Simberloff
33 2009). The use of genetic studies to gain insight into various aspects of the invasion process,
34 such as the mode and frequency of introduction, can help predict the potential likelihood and
35 impacts of further invasion (e.g. Provan et al. 2008; Xavier et al. 2009) and also offer
36 information helpful for management and control (Allendorf & Lundquist 2003).

37 The impacts of invasive species are of global concern (Lowe et al. 2000; Simberloff et al.
38 2013). Introduced deer species represent one such problem, with more than a quarter of deer
39 species having been introduced outside of their native range (Dolman & Waeber 2008). Over-
40 abundant deer negatively impact biodiversity (Cote et al. 2004), commercial land use
41 (Putman & Moore 1998), and human health and well-being through potentially fatal deer-
42 vehicle collisions (Bruiderink & Hazebroek 1996). The arrival of non-native deer species can
43 impose ecological pressure on woodland ecosystems, with new species moving into

44 unoccupied niches detrimentally impacting native flora and fauna (White et al. 2008). With
45 deer introductions and secondary spread continuing, such as the arrival and establishment of
46 muntjac throughout Ireland (Dick et al. 2010, 2012), knowledge of invasion history is critical
47 to assess risks of future ecological and economic damage through population expansion.

48 Outside of their native range of south-east China and Taiwan, Reeves's Chinese muntjac
49 (*Muntiacus reevesi*, Ogilby, 1839) have been introduced to France, Japan and the British
50 Isles, though only the latter two countries now have established populations (Lever 2009;
51 Ohdachi et al. 2009). The post-introduction natural range expansion of muntjac in the British
52 Isles has been around 1km per year, similar to other introduced deer (Chapman et al. 1994).
53 However, their full range expansion has been supplemented by secondary inocula via
54 human-mediated dispersal (Smith-Jones 2004). Their invasive success is further facilitated
55 by year-round breeding, rapid reproductive maturity (Chapman et al. 1997), and an ability to
56 inhabit anthropogenically modified habitats (Dansie, 1983). Indeed, the Game and Wildlife
57 Conservancy Trust reported a 1,756% increase in the numbers of muntjac shot between 1961-
58 2009 (Aebische et al, 2011), with the national population estimated very conservatively at
59 52,250 individuals by 2008 (Harris and Yalden, 2008). This rapid dispersal and increase in
60 abundance in such a relatively short time has resulted in considerable ecological and
61 economic damage (Cooke & Farrell 2001; Dolman and Waeber 2008; Mayle 2002).
62 Browsing and grazing pressure by muntjac has had major impacts on woodland ground flora
63 diversity and tree regeneration (Joys et al. 2004) and, as a result, there have been cascade
64 effects on other taxa, including rare butterflies such as the heath fritillary (*Melitaea athalia*)
65 and wood white (*Leptidea sinapis*) (Tabor 1998). The removal of ground cover by muntjac
66 reduces nesting sites for woodland songbirds (Holt et al. 2011) and is also thought to be
67 responsible for the reduction in woodland small mammal populations (Flowerdew & Elwood
68 2001).

69 Historical information on the sourcing, supply and release of muntjac in the British Isles is
70 highly confusing (Chapman et al, 1994). Records indicate that a pair of Reeves' muntjac were
71 presented to the Zoological Society of London in 1838 by John Russell Reeves, possibly
72 sourced from Guangdong (Canton) Province, South China. In 1867, the Zoo purchased a
73 replacement male obtained by Robert Swinhoe, possibly sourced from Formosa (also known
74 as Chinese Taipei, or more commonly now, Taiwan, which has its own subspecies known as
75 Taiwanese Reeves's muntjac *Muntiacus reevesi micrurus*). The female died shortly after
76 giving birth and was replaced in 1873 with a female from Ningpo. The species is known to
77 have bred at the London Zoo during this period. In November 1874, another pair from
78 Formosa (also known as Chinese Taipei, or more commonly now, Taiwan, which has its own
79 subspecies known as Taiwanese Reeves's muntjac *Muntiacus reevesi micrurus*) were
80 presented to the Zoological Society of London and, in the same year, a further female was
81 sent from Ningpo and a male from Hong Kong, with the parents of the latter believed to have
82 originated from Formosa (Sclater, 1875). Out of the 14 births that were recorded at London
83 Zoo between 1874 and 1881, nine were sold to Tring Park, England, and Jardin des Plants,
84 France, amongst other possible locations and between 1890 and 1928 no records of muntjac
85 in London Zoo exist (Chapman et al 1994).

86 Woburn Abbey had an initial import of three pairs of muntjac from a dealer on three
87 separate occasions in 1893 (Chapman et al. 1994). These individuals may have come direct
88 from wild caught animals in China, but it is likely that they were captive-bred individuals, as,
89 by this time, at least six other collections were known to keep and trade muntjac in Europe,
90 including Rambouillet, France (Dansie, 1983) and Berlin Zoo, Germany (Chapman et al.
91 1994). The 11th Duke of Bedford was also well known for travelling around Europe
92 collecting deer for his park in Woburn Abbey (Dansie, 1983). What is known is that he
93 received a male and two females, most likely of the London Zoo descendents, from Jardin

94 des Plants in 1902 (Chapman et al 1994). In total between 1894 and 1906, 13 males and 15
95 females were brought into Woburn Park and records show that eleven individuals were
96 released from there in 1901 (Chapman 1993).

97 Until 2009, it was widely assumed that muntjac had not reached Scotland or Ireland
98 (Lever 2009). Recent deer surveys, however, suggest that muntjac have spread westward into
99 Wales and south-west England as well as northward up the eastern half of England to the
100 Scottish Borders (Ward et al. 2008), and most recently have been discovered in Ireland
101 (Hailstone 2012). The first confirmed sighting of muntjac in the wild in the Republic of
102 Ireland was a culled animal in Co. Wicklow (Carden et al. 2011), swiftly followed by a
103 carcass from a deer-vehicle collision in Co. Down, Northern Ireland (Dick et al. 2010). The
104 question remains, however, as to the size of the actual founding ‘propagule’ that led to this
105 dramatic and continuing invasion of the British Isles. Such information is important to help
106 understand and predict invasion success in general and with regards to the likelihood of
107 further invasions by non-native deer globally.

108 In the only population genetic analysis of muntjac in Britain carried out to date, it was
109 suggested that there were at least eight maternal lineages, and most likely more, based on
110 restriction fragment length polymorphism (RFLP) analysis of the mitochondrial genome
111 (Williams et al. 1993). However, mitochondrial markers only give an indication of female-
112 mediated gene flow, and analysis of high-resolution nuclear microsatellite markers offers a
113 more complete picture of the invasion history of a species (Guillemaud et al. 2010).
114 Consequently, in the present study we carried out a combined analysis using microsatellites
115 and sequencing of the mitochondrial D-loop region to resolve the invasion history of the
116 species in the British Isles, specifically the likely number of founding females.

117 **Materials and methods**

118

119 **Sampling and DNA extraction**

120 Tissue samples were collected from 176 muntjac across the majority of their known
121 distribution in the British Isles (Figure 1; Table 1; Appendix 1). Samples were collected as
122 part of ongoing control programme by deer managers from the British Deer Society (BDS).
123 Tissue samples, mostly tongue or ear clippings (1 cm³), were collected and stored in absolute
124 ethanol. In addition, five DNA samples of the Taiwanese subspecies *Muntiacus reevesi*
125 *micrurus* were obtained directly from Taiwan. DNA was extracted from tissue samples using
126 a high salt extraction technique (Aljanabi & Martinez 1997).

127

128 **Genotyping**

129 All samples were genotyped for eight microsatellite loci originally developed for *M.*
130 *crinifrons* (Wu et al. 2008): Mcr-3, Mcr-4, Mcr-5, Mcr-6, Mcr-7, Mcr-13, Mcr-14 and Mcr-
131 19. The remaining three loci reported by Wu et al. (2008) could not be amplified reliably.
132 PCR was carried out in a total volume of 10 µl containing 100 ng genomic DNA, 5 pmol of
133 6-FAM-labelled M13 primer, 0.5 pmol of M13-tailed forward primer, 5 pmol reverse primer,
134 1x PCR reaction buffer, 200 µM each dNTP, 2.5 mM MgCl₂ and 0.25 U GoTaq Flexi DNA
135 polymerase (Promega, Sunnyvale, CA, USA). PCR was carried out on a MWG Primus
136 thermal cycler (Ebersberg, Germany) using the following conditions: initial denaturation at
137 94 °C for 3 min followed by 40 cycles of denaturation at 94 °C for 30 s, annealing at 55 °C
138 for 30 s, extension at 72 °C for 30 s and a final extension at 72 °C for 5 min. Genotyping was
139 carried out on an AB3730xl capillary genotyping system. (Life Technologies; Carlsbad,
140 California, USA). Allele sizes were scored using the GENEMAPPER software package (v4.1;

141 Applied Biosystems) using LIZ-500 size standards, and were checked by comparison with
142 previously sized control samples. All chromatograms were inspected visually.

143

144 **Mitochondrial D-loop sequencing**

145 The complete mitochondrial D-loop region was amplified using a pair of primers designed
146 from the complete mitochondrial genome of *M. reevesi* (GenBank accession number
147 AF527537): trn-Pro-F 5'-TCAACACCCAAAGCTGAAGTT-3 and trn-Phe-R 5'-
148 TCAGTGCCTTGCTTTATTGC-3. PCR was carried out in a total volume of 20 µl
149 containing 200 ng genomic DNA, 10 pmol of each primer, 1x PCR reaction buffer, 200 µM
150 each dNTP, 2.5 mM MgCl₂ and 0.5 U GoTaq Flexi DNA polymerase (Promega, Sunnyvale,
151 CA). PCR was carried out on a MWG Primus thermal cycler (Ebersberg, Germany) using the
152 following parameters: initial denaturation at 94 °C for 3 min followed by 40 cycles of
153 denaturation at 94 °C for 30 s, annealing at 58 °C for 30 s, extension at 72 °C for 1 min and a
154 final extension at 72 °C for 5 min. Five µl PCR product were resolved on 1.5% agarose gels
155 and visualised by ethidium bromide staining, and the remaining 15 µl were EXO-SAP
156 purified and sequenced in both direction using the BigDye sequencing kit (V3.1; Applied
157 Biosystems) using the primers Munt-DLOOP-IN-F 5'-ATCCTTGTC AACATGCGTATC-3'
158 and Munt-DLOOP-IN-R 5'-TTATGTGTGAGCATGGGCTG-3' and run on an AB 3730XL
159 DNA analyser (Life Technologies; Carlsbad, California, USA).

160

161 **Data analysis**

162 GENEPOP (V3.4; Raymond and Rousset, 1995) was used to test for linkage disequilibrium
163 between nuclear microsatellite loci. MICRO-CHECKER (van Oosterhout et al. 2004) was
164 used to check for the possible occurrence of null alleles. To estimate genetic diversity within
165 populations containing six or more individuals (Table 1), levels of observed (H_o) and

166 expected (H_E) heterozygosity, levels of allelic richness (A_R) and fixation indices (F_{IS}) were
167 calculated using the FSTAT software package (V2.9.3.2; Goudet, 2001). Significance of F_{IS}
168 was determined by 10,000 randomisation steps. Levels of genetic diversity (H) based on
169 mtDNA D-loop haplotype frequencies were calculated using the ARLEQUIN software package
170 (V3.5.1.2; Excoffier and Lischer, 2010).

171 The overall level of genetic differentiation between populations was estimated using Φ_{ST} ,
172 which gives an analogue of F_{ST} (Weir and Cockerham, 1984) calculated within the analysis
173 of molecular variance (AMOVA) framework (Excoffier et al. 1992) using ARLEQUIN. To
174 further identify possible patterns of genetic structuring, the software package BAPS (V5;
175 Corander et al. 2003) was used to identify clusters of genetically similar populations based on
176 the complete microsatellite data set (181 individuals) using a Bayesian approach. Ten
177 replicates were run for all possible values of the maximum number of clusters (K) up to $K =$
178 40, with a burn-in period of 10,000 iterations followed by 100,000 iterations. Multiple
179 independent runs always gave the same outcome. To further identify possible spatial patterns
180 of gene flow, a principal coordinate analysis (PCA) was carried out on the population-level
181 data set (105 individuals; Table 1) in GENALEX (V6.1; Peakall & Smouse 2006). Inter-
182 individual genetic distances were calculated as described in Smouse & Peakall 1999, and the
183 PCA was carried out using the standard covariance approach.

184 To test for the occurrence of a genetic bottleneck, the Wilcoxon test for heterozygote
185 excess was performed under the infinite alleles model (IAM), the stepwise mutation model
186 (SMM) and a two-phase model (TPM) incorporating 90% single-stepwise mutations using
187 the program BOTTLENECK (V1.2; Piry *et al.* 1999). The Wilcoxon test was used as it is
188 recommended for less than 20 microsatellite loci.

189 Results

190

191 No evidence of linkage disequilibrium was detected between any of the eight nuclear
192 microsatellites analysed. Between eight (Mcr-3) and 16 (Mcr-5) alleles were detected per
193 locus, with a total of 98 (mean = 12.25 per locus). Within populations for which a minimum
194 of six individuals were sampled, levels of allelic richness (A_R) averaged over loci ranged
195 from 3.723 (Kineton) to 5.120 (Sandlings), with a mean value of 4.136 (Table 2). Levels of
196 observed (H_O) and expected (H_E) heterozygosity ranged from 0.450 (Welford) to 0.663
197 (Wytham Wood; mean = 0.558), and from 0.675 (Kineton) to 0.855 (Sandlings; mean =
198 0.755) respectively. The heterozygote deficit observed in the majority of the populations gave
199 rise to F_{IS} values which were significantly different from zero in all of the populations
200 studied, ranging from 0.140 (Wytham Wood) to 0.420 (Welford; mean = 0.270), which is
201 consistent with the departure from Hardy-Weinberg equilibrium generally associated with
202 invasive species. MICRO-CHECKER indicated the possibility of null alleles at four of the
203 eight loci (Mcr-5, Mcr-7, Mcr-13 and Mcr-14), which could at least in part explain this
204 heterozygosity deficit.

205 Complete D-loop sequences were obtained for 121 individuals, with an alignment length
206 of 815 bp. A total of 23 substitution mutations gave rise to seven haplotypes (Figure 2). No
207 indels were observed, which can often make the alignment of D-loop sequences difficult.
208 There was a notable east-west cline in the frequency of haplotypes (Figure 1). Haplotype
209 diversity values for populations for which a minimum of six individuals were sampled ranged
210 from zero (Ickworth Park, for which only two complete sequences were obtained) to 0.800
211 (Sennowe Park; mean = 0.430). As the mitochondrial D-loop region exhibits an extremely
212 fast mutation rate, and thus provides a high-resolution marker for female lineages, since the
213 mitochondrial genome is maternally inherited in mammals (Harrison 1989), the results

214 indicate a low number of founding females. Two of the seven haplotypes occurred at very
215 low frequency. All seven haplotypes were split into four very divergent groups, each
216 separated by at least seven mutations. It is possible that Haplotype H6 evolved from
217 Haplotype H2 after the introduction, since the two differ by a single mutation, although it is
218 difficult to be more certain in the absence of an accurate mutation rate for this region.
219 Likewise, Haplotype H7 is only found in a single individual and is one mutation removed
220 from Haplotype H5, suggesting very recent divergence.

221 The overall level of differentiation estimated by nuclear microsatellites was low ($\Phi_{ST} =$
222 $0.050, P < 0.001$), whilst the level based on mitochondrial D-loop sequences was much
223 higher ($\Phi_{ST} = 0.470, P < 0.001$; Table 2). The BAPS analysis indicated that all the
224 individuals analysed from Britain and Ireland were grouped into a single genetic cluster
225 (100% probability), separate from the Taiwanese subspecies *Muntiacus reevesi*. This was
226 reflected in the PCA, which showed no evidence of geographical structuring of individual
227 multilocus genotypes (Figure 3). Finally, the Wilcoxon test for heterozygote excess
228 suggested a bottleneck under the IAM ($P = 0.006$), but not under the SMM or the TPM,
229 although it should be borne in mind that the number of loci used (eight) may be insufficient
230 to detect the latter (see Discussion).

231 Discussion

232

233 The probability of invasion success generally increases with propagule pressure, in terms of a
234 high number of viable founding individuals and repeated introductions (Simberloff 2009).

235 However, we also know that some invasions appear to establish from small founding events,

236 including the well-known case of the green seaweed *Caulerpa taxifolia*, which spread

237 throughout the Mediterranean rapidly following a single aquarium release (Jousson et al.

238 1998), and the suggestion that a single pair of squirrels would have a greater than 50%

239 chance of establishing a new population (Bertolino 2009). However, the invasion history of

240 many alien species is unknown. Resolving the size of founding propagules of major invasions

241 could help predict future invasions. Here, the use of high-resolution microsatellite markers

242 suggested that the invasion of the British Isles by Chinese Reeve's muntjac resulted from a

243 very small founding population. Indeed, our analysis is not only consistent with the known

244 introduction history of muntjac, but implies, through combined nuclear and mtDNA data, that

245 the current population was founded by a single group including a low number of females.

246 From Chapman et al. (1994), it seems probable to suggest those females are descendents of

247 some of the original five females imported to London Zoo amongst others imported

248 elsewhere.

249 The distribution of mtDNA haplotypes across southeast England is consistent with

250 separate escapes and releases from Woburn and other captive collections, since Haplotypes

251 H2 and H3, represented in yellow and blue respectively, tend to be primarily found west of

252 Woburn, which lies just northwest of Site 4 in Figure 1, and indeed are the only two

253 haplotypes found in this area, consistent with an extreme maternal founder effect. Likewise,

254 Haplotypes H1 and H4, shown in red and green, are not found west of Woburn, and are the

255 dominant haplotypes in sites to the east. The only previous genetic study on muntjac in the

256 UK suggested “at least eight maternal lineages in the UK” (Williams et al. 1993). This was
257 based on RFLP analysis of the same D-loop region analysed in the present study, but this
258 approach, unlike the sequencing analysis carried out here, cannot elucidate the genetic
259 relationships between haplotypes. Three of the eight haplotypes detected by Williams et al.
260 (1993) occurred at frequencies of 1.7%, 0.5% and 0.25% and could, as in the present study,
261 represent recent, post-introduction mutations. However, the lack of information on the
262 genealogical relationships between the RFLP haplotypes meant that this aspect could not be
263 addressed in the earlier study, but their identification of four or five haplotypes at relatively
264 high frequency (5% or above) is consistent with our findings, particularly when considering
265 that they analysed over three times as many samples. It is difficult to quantify accurately the
266 number of females involved in the introduction, since many of the original mitochondrial
267 lineages may have become extinct since the original founding event, but both studies indicate
268 a low number of individuals. The additional use of high-resolution microsatellite markers in
269 the present study suggests a single introduction of muntjac into the Britain Isles, since levels
270 of genetic differentiation were very low. As invasion events, single or multiple, are generally
271 characterised by founder effects, the random nature of these episodes means that multiple
272 events tend to involve separate gene pools from the original source population (Provan et al.
273 2005). This is contrary to the assignment of all individuals sampled from Britain and Ireland
274 in the present study to a single genetic cluster in the BAPS analysis, and the lack of any
275 geographical structuring in the PCA. The results of the BOTTLENECK analysis were
276 somewhat inconclusive, with a genetic bottleneck suggested under the IAM but not under the
277 other two models. This could be due to the fact that the two models that assume stepwise
278 mutation, particularly the SMM, are more conservative than the IAM (Cornuet and Luikart
279 1996; Luikart and Cornuet 1998). Alternatively, it may be that the short generation time of
280 muntjac, which can start breeding at around 36 weeks (Chapman et al. 1997), means that

281 sufficient generations have passed since introduction to mask the signature of any genetic
282 bottleneck.

283 Our findings highlight the risk of directly equating propagule pressure with the success of
284 an invasion, since, despite a potentially high number of released individuals across multiple
285 sites, the data here show that the invasion is descended from the same import source which
286 consisted of a limited number of founders. Given that invasive deer species have been found
287 to have earlier weaning and sexual maturity age (Fautley et al. 2012), the success of muntjac
288 invasion undoubtedly appears to be due their high fecundity and rate of increase in the initial
289 stages, as well as secondary introductions at multiple locations over a sustained period of
290 time. However, instead of offering a chance for an increase in genetic variation, due to the
291 restricted original gene pool, the sole advantage of these multiple release sites appears to be
292 demographic, by offering additional mates if a nearby population begins to fail. Invasion
293 success, in spite of population bottlenecks, has previously been reported in species once kept
294 as exotic pets or decorative plants (Le Page et al. 2000; Goodman et al. 2001; Dlugosch &
295 Parker 2008b). It is possible that human-mediated species introductions such as those from
296 ornamental and/or pet species have an increased chance of invasion due to *ex-situ* breeding,
297 despite low genetic diversity and limited primary introductions from the native range.

298 More optimistically, this finding also has implications for captive management of
299 conservation reintroductions. Though low genetic diversity is not considered ideal in species
300 reintroduction programmes (Frankham et al. 2002), this study supports the idea that a
301 successful introduction can result from a low number of individuals with limited genetic
302 variation. Many examples of successful reintroductions from low number of founding
303 individuals exist. Taylor et al. (2005) found that the number of released New Zealand
304 saddlebacks (*Philesturnus carunculatus*) and robins (*Petroica australis*) on different islands
305 did not affect the success of introductions. The alpine ibex (*Capra ibex ibex*), carefully bred

306 in captivity from a very low number of founding individuals, was reintroduced successfully
307 on several occasions (Stüwe & Nievergelt 1991). Most notably, the milù (*Elaphurus*
308 *davidianus*), also known as Père David's deer, was successfully reintroduced to China having
309 been rescued from just two females and a single male after careful captive breeding by the
310 11th Duke of Bedford (Zeng et al. 2007). In all cases, success was based on a combination of
311 selective breeding to increase the effective captive population size and multiple releases.

312 Conversely, our paper offers an insight for legislative policy in invasive species
313 management. Our data are consistent with four or five females leading to a major geographic
314 invasive species problem. This should serve as a warning for future muntjac invasions, such
315 as those unfolding in Ireland currently (Dick et al. 2010; Freeman et al. 2011; Dick et al.
316 2012; Hogg et al. 2014) and a suspected introduction in Belgium (T. Adrianens, pers.
317 comm.). It appears quite plausible that, in both cases, muntjac could colonise a large area,
318 giving rise to large numbers of individuals, from relatively few founding individuals. We
319 raise the point to caution against complacency if invasive species such as muntjac are
320 suspected in a new area and we would advocate heightened biosecurity and a need to react to
321 sightings rapidly (Caffrey et al. 2014). Indeed, under recent EU legislation, inclusion of
322 muntjac on the list of high-risk species would restrict the movement and release of the
323 species (Genovesi et al. 2014). The present study is an example of the value of genetics to
324 invasion ecology, and also helps to illuminate the origin of muntjac deer populations in the
325 British Isles. Vigilance should be exercised even with small releases of species such as
326 muntjac, which may, if ignored, result in an irreversible invasion, regardless of initial genetic
327 diversity or continuous genetic influx.

328 **Acknowledgements**

329

330 Marianne Freeman was supported by Department of Employment and Learning (DEL) and
331 Neil Reid was supported by the Natural Heritage Research Partnership (NHRP) between the
332 Northern Ireland Environment Agency (NIEA) and *Quercus*, Queen's University Belfast
333 (QUB). We are grateful to the members of the BDS for taking the time to send tissue samples
334 and to Kurtis Pei (Professor of Wildlife Ecology & Management, National Pingtung
335 University of Science & Technology, Taiwan) for the *M. reevesi micrurus* samples. Required
336 permits for the export of samples from Taiwan were obtained from the Council of Agriculture
337 and import to the UK from the Department for Environment, Food and Rural Affairs
338 (Authorisation number POAO/2010/151). We are also grateful to Doug Armstrong and an
339 anonymous reviewer, whose comments improved the manuscript.

340 **References**

341

342 Aebischer, N.J., Davey, P.D. & Kingdon, N.G. (2011). National Gamebag Census: Mammal
343 Trends to 2009. Game & Wildlife Conservation Trust, Fordingbridge
344 (<http://www.gwct.org.uk/ngcmammals>)

345 Allendorf, F.W. & Lundquist, L.L. (2003). Introduction: population biology, evolution and
346 control of invasive species. *Conservation Biology* **17**, 24-30.

347 Aljanabi, S.M. & Martinez, I. (1997). Universal and rapid salt-extraction of high quality
348 genomic DNA for PCR-based techniques. *Nucleic Acids Research* **25**, 4692-4693

349 Bertolino, S. (2009) Animal trade and non-indigenous species introduction: the world-wide
350 spread of squirrels. *Diversity and Distributions* **15**, 701–708.

351 Bruiderink, G.W.T.A.G. & Hazebroek, E. (1996). Ungulate traffic collisions in Europe.
352 *Conservation Biology* **10**, 1059-1067.

353 Caffrey, J.M., Baars, J., Barbour, J.H., Boets, P., Boon, P., Davenport, K., Dick, J.T., Early,
354 J., Edsman, L. & Gallagher, C. (2014). Tracking invasive alien species in Europe: the top
355 20 issues. *Management of Biological Invasions* **5**, 1-20.

356 Carden, R.F., Carlin, C.M., Marnell, F., McElholm, D., Hetherington, J. & Gammell, M.P.
357 (2011). Distribution and range expansion of deer in Ireland. *Mammal Review* **41**, 313-325.

358 Chapman, N.G. (1993). Current distribution, status and habitat preferences of Reeves’
359 muntjac. in Mayle BAE ed (1993) *Muntjac Deer; Their Biology, Impact and Management*
360 *in Britain*. Forestry Commission, Farnham and British Deer Society, Trentham.

361 Chapman, N., Harris, S. & Stanford, A. (1994). Reeves’ muntjac, *Muntiacus reevesi*, in
362 Britain – their history, spread, habitat selection, and the role of human intervention in
363 accelerating their dispersal. *Mammal Review* **24**, 113-160.

364 Chapman, N.G., Furlong, M. & Harris, S. (1997). Reproductive strategies and the influence
365 of date of birth on growth and sexual development of an aseasonally-breeding ungulate:
366 Reeves' muntjac (*Muntiacus reevesi*). *Journal of Zoology* **241**, 551-570.

367 Cooke, A. & Farrell, L. (2001). Impact of muntjac deer (*Muntiacus reevesi*) at Monks Wood
368 National Nature Reserve, Cambridgeshire, eastern England. *Forestry* **74**, 241.

369 Corander, J., Waldmann, P. & Sillanpää, M.J. (2003). Bayesian analysis of genetic
370 differentiation between populations. *Genetics* **163**, 367-374.

371 Cornuet, J-M. & Luikart, G. (1996). Description and power analysis of two tests for detecting
372 recent population bottlenecks from allele frequency data. *Genetics* **144**, 2001-2014.

373 Cote, S.D., Rooney, T.P., Tremblay, J.P., Dussault, C. & Waller, D.M. (2004). Ecological
374 impacts of deer overabundance. *Annual Review of Ecology, Evolution and Systematics* **35**,
375 113-147.

376 Dansie, O. (1983) Muntjac. In: *Muntjac and Chinese Water Deer*. pp 3-24. The British Deer
377 Society, Warminster, UK.

378 Dick, J.T.A., Freeman, M., Provan, J. & Reid N. (2010). First record of free-living Reeves'
379 muntjac deer (*Muntiacus reevesi* Ogilby, 1939) in Northern Ireland. *Irish Naturalists'*
380 *Journal* **31**, 151.

381 Dick, J.T.A., McKillen, J., Chapman, N., Collins, L., Provan, J., Freeman, M., Hogg, K.,
382 Reid, N. (2012). *Post-mortem Examination of a Wild Muntjac from Northern Ireland*.
383 Report prepared by the Natural Heritage Research Partnership (NHRP) between *Quercus*,
384 Queen's University Belfast, and the Northern Ireland Environment Agency (NIEA).
385 Northern Ireland Environment Agency Research and Development Series No 12/12,
386 Belfast, UK.

387 Dlugosch, K.M. & Parker, I.M. (2008a) Founding events in species invasions: genetic
388 variation, adaptive evolution, and the role of multiple introductions. *Molecular Ecology*
389 **17**, 431-449.

390 Dlugosch, K.M. & Parker, I.M. (2008b) Invading populations of an ornamental shrub show
391 rapid life history evolution despite genetic bottlenecks. *Ecology Letters* **11**, 701-709.

392 Dolman, P.M. & Waeber, K. (2008). Ecosystem and competition impacts of introduced deer.
393 *Wildlife Research* **35**, 202-214.

394 Excoffier, L., Smouse, P.E. & Quattro, J.M. (1992). Analysis of molecular variance inferred
395 from metric distances among DNA haplotypes - application to human mitochondrial DNA
396 restriction data. *Genetics* **131**, 479-491.

397 Excoffier, L. & Lischer, H.E.L. (2010). ARLEQUIN suite ver 3.5: a new series of programs to
398 perform population genetics analyses under Linux and Windows. *Molecular Ecology*
399 *Resources* **10**, 564-567.

400 Fautley, R., Coulson, T. & Savolainen, V. (2012) A comparative analysis of the factors
401 promoting deer invasion. *Biological Invasions* **14**, 2271-2281.

402 Flowerdew, J. & Ellwood, S. (2001) Impacts of woodland deer on small mammal ecology.
403 *Forestry* **74**, 277.

404 Frankham, R., Briscoe, D.A. & Ballou, J.D. (2002). *Introduction to Conservation Genetics*.
405 Cambridge University Press, London, UK.

406 Freeman, M., Reid, N. & Dick, J.T.A. (2011) *Evaluation of the Veracity of Anecdotal*
407 *Muntjac Records in Northern Ireland*. Report prepared by the Natural Heritage Research
408 Partnership (NHRP) between *Quercus*, Queen's University Belfast, and the Northern
409 Ireland Environment Agency (NIEA). Northern Ireland Environment Agency Research
410 and Development Series No 11/07, Belfast, UK.

411 Genovesi P., Carboneras C., Vilà M. & Walton P. (2014) EU adopts innovative legislation on
412 invasive species: a step towards a global response to biological invasions? *Biological*
413 *Invasions* **17**, 1307–1311.

414 Goodman, S.J., Tamate, H.B., Wilson, R., Nagata, J., Tatsuzawa, S., Swanson, G.M.,
415 Pemberton, J.M. & MuCullough, D.R. (2001). Bottlenecks, drift and differentiation; the
416 population structure and demographic history of sika deer (*Cervus nippon*) in the Japanese
417 archipelago. *Molecular Ecology* **10**, 1357-1370.

418 Goudet, J. (2001). FSTAT, A program to estimate and test gene diversities and fixation
419 indices (Version 2.9.3.) <http://www2.unil.ch/popgen/softwares/fstat.htm> Accessed 16
420 January 2013.

421 Guillemaud, T., Beaumont, M.A., Ciosi, M., Cornuet, J-M., Estoup, A. (2010). Inferring
422 introduction routes of invasive species using approximate Bayesian computation on
423 microsatellite data. *Heredity* **104**, 88-99.

424 Hailstone, M. (2012). The Great British Deer Survey. *Deer* **16**, 12-15.

425 Harris, S., Yalden, D.W. (2008). *Mammals of the British Isles Handbook*. Mammal Society,
426 Southampton, UK..

427 Harrison, R.G. (1989). Animal mitochondrial DNA as a genetic marker in population and
428 evolutionary biology. *Trends in Ecology and Evolution* **4**, 6-11.

429 Hogg, K., Freeman, M., Burdekin, O., Lennon, J., Reid, N., Dick, J.T.A. (2014). *Ards*
430 *Peninsula muntjac deer (Muntiacus reevesi) survey*. Report prepared by the Natural
431 Heritage Research Partnership (NHRP) between *Quercus*, Queen’s University Belfast and
432 the Northern Ireland Environment Agency (NIEA) for the Research and Development
433 Series No. 13/XX.

434 Hollingsworth, M.L. & Bailey, J.P. (2000). Evidence for massive clonal growth in the
435 invasive weed *Fallopia japonica* (Japanese Knotweed). *Botanical Journal of the Linnean*
436 *Society* **133**, 463-472.

437 Holt, C.A., Fuller, R.J. & Dolman, P.M. (2011). Breeding and post-breeding responses of
438 woodland birds to modification of habitat structure by deer. *Biological Conservation* **144**,
439 2151-2162.

440 Jousson, O., Pawlowski, J., Zaninetti, L., Meinesz, A. & Boudouresque, C.F. (1998)
441 Molecular evidence for the aquarium origin of the green alga *Caulerpa taxifolia*
442 introduced to the Mediterranean Sea. *Marine Ecology Progress Series* **172**, 275-280.

443 Joys, A., Fuller, R. & Dolman, P. (2004) Influences of deer browsing, coppice history and
444 standard trees on the growth and development of vegetation structure in coppiced woods
445 in lowland England. *Forest Ecology and Management* **202**, 23-37.

446 Lavergne, S. & Molofsky, J. (2007). Increased genetic variation and evolutionary potential
447 drive the success of an invasive grass. *Proceedings of the National Academy of Sciences*
448 *USA* **104**, 3883-3888.

449 Le Page, S., Livermore, R., Cooper, D. & Taylor, A. (2000). Genetic analysis of a
450 documented population bottleneck: introduced Bennett's wallabies (*Macropus rufogriseus*
451 *rufogriseus*) in New Zealand. *Molecular Ecology* **9**, 753-763.

452 Lever, C. (2009). *The Naturalised Animals of Britain and Ireland*. New Holland Ltd.,
453 London.

454 Lockwood, J.L., Cassey, P. & Blackburn, T. (2005) The role of propagule pressure in
455 explaining species invasions. *Trends in Ecology and Evolution* **20**, 223-228.

456 Lockwood, J.L., Cassey, P. & Blackburn, T.M. (2009) The more you introduce the more you
457 get: the role of colonization pressure and propagule pressure in invasion ecology. *Diversity*
458 *and Distributions* **15**, 904-910.

459 Lowe, S., Browne, M., Boudjela, S. & De Poorter, M. (2000) *100 of the world's worst*
460 *invasive alien species: a selection from the global invasive species database*. Invasive
461 Species Specialist Group Auckland, New Zealand

462 Luikart, G. & Cornuet, J.-M. (1998). Empirical evaluation of a test for identifying recently
463 bottlenecked populations from allele frequency data. *Conservation Biology* **12**, 228-237.

464 Mayle, B.A.E. ed. (1993). *Muntjac Deer; Their Biology, Impact and Management in Britain*.
465 Forestry Commission, Farnham and British Deer Society, Trentham.

466 Ohdachi, S., Ishibashi, Y., Iwasa, M. & Saitoh, T. eds. (2009). *The Wild Mammals of Japan*.
467 Shoukadoh, Kyoto.

468 van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M. & Shipley, P. (2004) MICRO-
469 CHECKER: software for identifying and correcting genotyping errors in microsatellite
470 data. *Molecular Ecology Notes* **4**, 535-538.

471 Peakall, R. & Smouse, P.E. (2006). GENALEX 6 Genetic analysis in Excel. Population
472 genetic software for research and teaching. *Molecular Ecology Notes* **6**, 288-295.

473 Piry, A., Luikart, G. & Cornuet, J.M. (1999). Bottleneck: a computer program for detecting
474 recent reductions in the effective population size using allele frequency data. *Journal of*
475 *Heredity* **90**, 502-503.

476 Provan, J., Murphy, S., Maggs, C.A. (2005). Tracking the invasive history of the green alga
477 *Codium fragile* ssp. *tomentosoides*. *Molecular Ecology* **14**, 189-194.

478 Provan, J., Booth, D., Todd, N.P., Beatty, G.E. & Maggs, C.A. (2008) Tracking biological
479 invasions in space and time: elucidating the invasive history of the green alga *Codium*
480 *fragile* using old DNA. *Diversity and Distributions* **14**, 343-354.

481 Putman, R.J. & Moore, N.P. (1998). Impact of deer in lowland Britain on agriculture, forestry
482 and conservation habitats. *Mammal Review* **28**, 141-163.

483 Raymond, M. & Rousset, F. (1995). GENEPOP (version 1.2): population genetic software for
484 exact tests and ecumenicism. *Journal of Heredity* **86**, 248-249.

485 Roman, J. & Darling, J.A. (2007). Paradox lost: genetic diversity and the success of aquatic
486 invasions. *Trends in Ecology and Evolution* **22**, 454-464.

487 Sax, D.F. & Brown, J.H. (2000) The paradox of invasion. *Global Ecology and Biogeography*
488 **9**, 363-371.

489 Sclater, P.L. (1875). Notes on rare mammals in the Society's collection. *Proceedings of the*
490 *Scientific Meetings of the Zoological Society of London* 421-422

491 Simberloff, D. (2009). The role of propagule pressure in biological invasions. *Annual Review*
492 *of Ecology, Evolution and Systematics* **40**, 81-102.

493 Simberloff, D., Martin, J-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp,
494 F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E. & Vila, M.
495 (2013). Impacts of biological invasions: what's what and the way forward. *Trends in*
496 *Ecology and Evolution* **28**, 58-66.

497 Smith-Jones, C. (2004). *Muntjac: Managing an Alien Species*. Coch-Y-Bonddu Books,
498 Powys.

499 Smouse, P.E. & Peakall, R. (1999). Spatial autocorrelation analysis of individual multiallele
500 and multilocus genetic structure. *Heredity* **82**, 561-573.

501 Stüwe, M. & Nievergelt, B. (1991). Recovery of alpine ibex from near extinction: the result
502 of effective protection, captive breeding, and reintroductions. *Applied Animal Behaviour*
503 *Science* **29**, 379-387.

504 Tabor, R.C.C. (1998). The oxlip in Britain - is its future in doubt? *British Wildlife* **10**, 92-
505 101

- 506 Taylor, S.S., Jamieson, I.G. & Armstrong, D.P. (2005). Successful island reintroductions of
507 New Zealand robins and saddlebacks with small numbers of founders. *Animal*
508 *Conservation* **8**, 415-420.
- 509 Ward, A., Etherington, T. & Ewald, J. (2008). Five years of change. *Deer* **12**, 17-20.
- 510 Weir, B.S. & Cockerham, C.C. (1984). Estimating F-statistics for the analysis of population
511 structure. *Evolution* **38**, 1358-1370.
- 512 White, P., Ford, A., Clout, M., Engeman, R., Roy, S. & Saunders, G. (2008) Alien invasive
513 vertebrates in ecosystems: pattern, process and the social dimension. *Wildlife Research* **3**,
514 171.
- 515 Williams, T., Harris, S., Chapman, N., Wayne, R., Beaumont, M. & Bruford, M. (1993). A
516 molecular analysis of the introduced Reeves' muntjac in southern England: genetic
517 variation in the mitochondrial genome. in Mayle BAE ed (1993) *Muntjac Deer; Their*
518 *Biology, Impact and Management in Britain*. Forestry Commission, Farnham and British
519 Deer Society, Trentham.
- 520 Wu, H., Meng, K. & Zhu, G. (2008). Isolation and characterization of microsatellite markers
521 in black muntjac (*Muntiacus crinifrons*). *Molecular Ecology Resources* **8**, 584-586.
- 522 Xavier, R., Santos, A., Lima, F. & Branco, M. (2009). Invasion or invisibility: using genetic
523 and distributional data to investigate the alien or indigenous status of Atlantic populations
524 of the peracarid isopod *Stenosoma nadejda* (Rezig 1989). *Molecular Ecology* **18**, 3283-
525 329.
- 526 Zeng, Y., Jiang, Z. & Li, C. (2007) Genetic variability in relocated Père David's deer
527 (*Elaphurus davidianus*) populations—Implications to reintroduction program.
528 *Conservation Genetics* **8**, 1051-1059.

Table 1 Details of populations studied where six or more individuals were analyzed with nuclear microsatellites. N – number of individuals analysed; A_R – allelic richness; H_O – observed heterozygosity; H_E – expected heterozygosity; F_{IS} – inbreeding coefficient; H1-H7 – frequency of mitochondrial haplotypes; H – haplotype diversity.

No	Name	Lat	Long	Nuclear microsatellites					Mitochondrial D-loop								
				(N)	(E)	N	A_R	H_O	H_E	F_{IS}	N	H1	H2	H3	H4	H5	H6
1	Kineton, Warwickshire	52.136	-1.470	12	3.723	0.563	0.675	0.173 ^{**}	12	-	11	1	-	-	-	-	0.167
2	Welford, Berkshire	51.478	-1.407	6	3.933	0.450	0.746	0.420 ^{***}	4	-	2	2	-	-	-	-	0.667
3	Wytham Wood, Oxfordshire	51.769	-1.334	10	4.099	0.663	0.764	0.140 [*]	10	-	7	3	-	-	-	-	0.467
4	Hexton, Bedfordshire	51.938	0.359	8	4.056	0.627	0.778	0.205 ^{**}	8	-	-	1	7	-	-	-	0.250
5	Ickworth Park, Suffolk	52.218	0.649	6	4.223	0.583	0.753	0.242 ^{**}	2	-	-	-	2	-	-	-	-
6	Stanta, Suffolk	52.494	0.720	9	3.768	0.500	0.700	0.298 ^{***}	8	5	-	-	-	3	-	-	0.536
7	Shadwell Estate, Norfolk	53.294	0.837	14	3.826	0.537	0.718	0.259 ^{***}	11	9	-	1	1	-	-	-	0.346
8	Sennowe Park, Norfolk	52.773	0.912	14	4.218	0.499	0.774	0.364 ^{***}	6	2	2	2	-	-	-	-	0.800
9	Sandlings, Suffolk	52.097	1.426	12	5.120	0.626	0.855	0.277 ^{***}	8	5	-	-	3	-	-	-	0.536
10	Sotterley, Suffolk	52.404	1.625	14	4.389	0.536	0.785	0.326 ^{***}	10	1	-	1	7	1	-	-	0.533

Table 2 Analysis of molecular variance (AMOVA).



Markers	Source of variation	Sum of squares	Variance	% variation	<i>P</i>
Nuclear microsatellites	Among populations	55.584	0.155	5.01	<i>P</i> < 0.001
	Within populations	588.655	2.943	94.99	
Mitochondrial D-loop	Among populations	15.206	0.190	47.00	<i>P</i> < 0.001
	Within populations	14.769	0.214	53.00	


Figure Legends

Figure 1 Map showing the distribution of mitochondrial D-loop haplotypes in south-eastern England. Inset shows Ireland. Colours refer to haplotypes in Figure 2. Circle size is proportional to the number of samples, with the largest circle representing $N = 12$ and the smallest $N = 1$. Dots indicate sites for which only microsatellite data were obtained. Numbers refer to Table 1 and Appendix 1. “W” indicates the location of Woburn Abbey.


Figure 2 Median-joining network showing relationships between the seven haplotypes detected by sequencing the mtDNA D-loop region. Circle sizes are approximately proportional to haplotype frequency: smallest circle represents a single individual, largest circle represents 22 individuals. Each dash between haplotypes represents a single mutation. The diamond represents an unsampled ancestral haplotype.

Figure 3 Results of the PCA. The first three axes accounted for 23.17%, 20.03% and 17.59% respectively of the total variation (60.78%).






Principal Coordinates (1 vs 2)



Principal Coordinates (1 vs 3)



- ◆ Bedfordshire
- Sennowe Park
- ▲ Shadwell Estate
- × Wytham Wood
- ✧ Ickworth Park
- Sandlings
- + Sotterley
- Stanta
- Kineton
- ◆ Welford

Appendix 1 Samples collected with less than six individuals at a site.

No	Location	Lat (N)	Long (E)	N_{nuc}	N_{mt}
11	Yarkhill, Hertfordshire	52.082	-2.573	3	3
12	Shuckburgh Estate, Warwickshire	52.254	-1.274	2	1
13	Cothill, Oxfordshire	51.670	-1.288	1	1
14	Wittenham Clumps, Oxfordshire	51.631	-1.189	1	1
15	Horsemoor Wood, Oxfordshire	51.456	-1.078	3	3
16	Lower Ianham, Hampshire	51.128	-1.131	1	1
17	Alton, Hampshire	51.157	-1.051	1	1
18	Old Elvendon, Oxfordshire	52.426	-1.068	2	2
19	Broughton Wood, Northamptonshire	52.432	-0.661	3	1
20	Finesmade, Northamptonshire	52.581	-0.557	1	-
21	Duddington, Northamptonshire	52.569	-0.542	1	-
22	Apethorpe, Northamptonshire	52.551	-0.487	2	-
23	Fothorngay, Northamptonshire	52.437	-0.444	1	1
24	Forty Acres Wood, Bedfordshire	52.230	-0.620	1	-
25	Great Hayes Wood, Bedfordshire	52.246	-0.588	4	2
26	Englefield Estate, Berkshire	51.430	-0.569	1	-
27	Chilworth, Surrey	51.214	-0.539	1	-
28	Bardney, Lincolnshire	53.215	-0.308	1	1
29	Woodhall Spa, Lincolnshire	53.158	-0.215	2	2
30	Welton Wood, Lincolnshire	53.202	0.197	2	2
31	Quendon, Essex	51.953	0.201	2	2
32	Ugley, Essex	51.931	0.198	1	-
33	Little Walden, Essex	52.067	0.239	1	1
34	Hamperden End, Essex	51.980	0.263	2	2
35	Lode, Cambridgeshire	52.241	0.239	1	-
36	Thurlow, Suffolk	52.108	0.384	1	-
37	Great Thurlew, Suffolk	52.084	0.447	1	-
38	Thurlow, Suffolk	52.137	0.480	1	1
39	Beachamwell, Norfolk	52.610	0.622	1	1
40	Thetford Forest, Norfolk	52.506	0.651	3	-
41	Rushbrooke, Suffolk	52.208	0.779	2	-
42	East Wretham, Norfolk	52.481	0.830	3	3
43	Haughley Park, Suffolk	52.222	0.920	1	1
44	Wolves' Wood, Suffolk	52.054	0.995	1	-
45	Tom's Wood, Suffolk	52.033	0.997	1	-
46	Holkham, Norfolk	52.950	0.804	4	2
47	Foxley Wood, Norfolk	52.765	1.036	2	1
48	Shelton, Norfolk	52.464	1.274	5	3
49	Sundy Green, Norfolk	52.484	1.293	1	1
50	Kirby Cane, Norfolk	52.493	1.493	2	-
51	Mount Stewart, Co. Down	54.557	-5.625	2	2