PRIFYSGOL ABERYSTWYTH UNIVERSITY

Aberystwyth University

Cultivars to face climate change effects on crops and weeds: a review

Korres, N. E.; Norsworthy, J. K.; Tehranchian, P.; Gitsopoulos, T. K.; Loka, D. A.; Oosterhuis, D. M.; Gealy, D. R.; Moss, S. R.; Burgos, N. R. O.

Published in:
Agronomy for Sustainable Development

DOI:
10.1007/s13593-016-0350-5

Publication date:
2016

Citation for published version (APA):
Korres, N. E., Norsworthy, J. K., Tehranchian, P., Gitsopoulos, T. K., Loka, D. A., Oosterhuis, D. M., Gealy, D. R., Moss, S. R., & Burgos, N. R. O. (2016). Cultivars to face climate change effects on crops and weeds: a review. *Agronomy for Sustainable Development*, *36*(1), Article 12. https://doi.org/10.1007/s13593-016-0350-5

General rights
Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
• You may not further distribute the material or use it for any profit-making activity or commercial gain
• You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy
If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Download date: 08. Sep. 2026

Developing cultivars to face climate change effects on crops and weeds. A review

Nicholas E. Korres[1*], Jason K. Norsworthy[1], Parsa Tehranchian[1], Thomas C. Gitsopoulos[2], Dimitra A. Loka[3], Derrick M. Oosterhuis[1], David R. Gealy[4], Stephen R. Moss[5], Nilda R. Burgos[1], M. Ryan Miller[1], Matheus Palhano[1]

[1]Dept. of Crop, Soil & Environmental Sciences, University of Arkansas, Fayetteville, AR, 72704 USA; [2]Hellenic Agricultural Organization-Demeter, Institute of Plant Breeding and Phytogenetic Resources, P.O. Box 60324, Thermi, Thessaloniki, Greece; [3]Aberystwyth University, UK; [4]USDA-ARS, Dale Bumpers National Rice Research Center, Stuttgart, AR, 72160, USA; [5]Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, UK

*Corresponding author: korres@uark.edu; nkorres@yahoo.co.uk

Abstract

Climate change caused by the release of greenhouse gases in the atmosphere due to various anthropogenic activities will impact many aspects of the human and natural world, but effects on agricultural production could be of particular significance. Estimates of annual damages in agriculture due to temperature increase or extended periods of drought, for example, will be more costly compared to those in other sectors and activities. Yield losses, as a result of climate change, are caused either through the direct effects of climate change components on crops or through indirect effects such as increased inputs in crop production necessary for the control of weeds, for example. To counteract the effects of climate change various adaptation strategies have been suggested. It has been suggested by several authors that the farmers' primary response to climate change would be to seek and crop cultivars that are most adapted to highly variable, extreme climatic conditions and pest changes brought forth by global warming.

Here we review the effects of climate change on crop cultivars alongside with these on weeds. Increases in marketable yield, mainly through increases in biomass, of cereals, particularly those that exhibit C3 photosynthetic pathway, range between 8-70%, these of row, cash and vegetable crops between 20-+144% and these of flowers between 6-35%. Nevertheless, the positive effects of elevated CO_2 on yield will most probably be affected by the availability of other resources such as water or nutrients. Temperature increases will decrease crop yields of temperature-sensitive crops such as maize, soybean, wheat, and cotton or specialty crops such as almonds, grapes, berries, citrus, or stone fruits at regional and local scale. Additionally crops like rice, which is expected to yield better under increased CO_2, will suffer serious yield losses under high temperatures. Significant crop yields reductions that occur under drought stress for tomato, soybean, maize, cotton are strongly demonstrated. Nevertheless, reviews on C4 photosynthesis response to water stress in interaction with CO_2 concentration revealed that elevated CO_2 concentration lessens the deleterious effect of drought on plant productivity. Weeds with C3 photosynthetic pathway are likely to respond more strongly than C4 types to CO_2 increases through biomass and leaf area increases. The positive response of C3 crops to elevated CO_2 may make C4 weeds less competitive for these crops whereas C3 weeds in C4 or C3 crops, particularly in tropical regions, could become a problem. Temperature increases will mainly affect the distribution of weeds, particularly C4 type, by expanding the geographical range they can be established. This will enhance further yield losses and will affect weed management systems negatively. In addition, the expansion of invasive weed species such as itchgrass, cogongrass and witchweed facilitated by temperature increases will increase the cost for their control. Under water- or nutrient shortage scenarios, an r-strategist with characteristics in the order S-C-R, such as Palmer amaranth, large crabgrass, johnsongrass and spurges will most

probably prevail. It is becoming obvious that selection of cultivars that secure high yields under climate change but also by competing weeds successfully is of major importance. Traits related with a) increased root:shoot ratio, b) vernalization periods, c) maturity, d) regulation of node formation and/or internode distance, e) harvest index variations and f) allelopathy merit further investigation. The cumulative effects of selecting a suitable stress tolerator-competitor cultivar will be reflected in reductions of environmental pollution, lower production costs and sustainable food production. It is therefore imperative to expand research efforts to investigate how crop-weed interference under various abiotic stresses and cropping systems influences cultivar performance and subsequent yield outcome.

Keywords: stress tolerance; climate change; carbon dioxide; drought; temperature; weed competition; competitive ability; cultivar selection; integrated weed management

Table of contents

Abstract 1

1.0 Introduction 5

2.0 Effects of climate change on crops 8

2.1 Effects of elevated CO_2

2.1.1 Effects of elevated CO_2 on crop physiological characteristics 8

2.1.2. Effects of elevated CO_2 on crop yields 9

2.2 Effects of temperature increases

2.2.1 Effects of temperature increases on crop physiological characteristics 9

2.2.2 Effects of temperature increases on crop yield 10

2.3 Effects of water deficit

2.3.1 Effects of water deficit on crop physiological characteristics 11

2.3.2 Effects of water deficit on crop yield 11

2.4 Interactive effects of climate change components on the physiology of crop plants and yield

3.0 Effects of climate change on weeds 13

3.1 Effects of elevated CO_2 13

3.1.1 Direct effects of elevated CO_2 on weeds 13

3.1.2 Indirect effects of elevated CO_2 on weeds 14

3.1.3. Effects of elevated CO_2 on crop-weed interference 15

3.2 Effects of temperature 15

3.2.1 Effects of temperature on weed physiological characteristics 15

3.2.2 Effects of temperature on weed distribution 16

3.3 Effects of water deficit 17

3.3.1 Effects of drought on weed physiological responses 17

3.3.2 Weed adaptation strategies under water deficit and other unfavorable conditions 18

3.4 Interactive effects of climate change components on weed performance and consequences on weed-crop competition 19

4.0 Cultivar selection against weeds and traits that confer competitiveness 21

4.1 Cultivar phenotypic characteristics and weed suppression 22

4.2 Implications for allelopathic properties 23

5.0 Traits for developing an ideotype S-C cultivar 24

5.1 *Cultivars with deep root system* 24

5.2 *Harvest index and dry mater components* 26

5.3 *Late maturing cultivars* 27

5.4 *Nutrients uptake and utilization* 27

5.5 *Heat tolerance-Improvements and expectations* 28

5.6 A *synthesis* 29

7.0. Conclusions 30

References 32

1.0 Introduction

Climate change refers to long term changes in the state of the climate (IPCC, 2014). These changes are identifiable i.e. the mean or the variability of climate change components such as increase of temperature or elevated atmospheric CO_2 levels can be assessed by the application of appropriate analytical and statistical methods (IPCC, 2014). The release of greenhouse gases (carbon dioxide, methane, nitrous oxide) due to various anthropogenic activities is very likely to be one of the major causes of recent climatic change (Glover et al., 2008). Plausible climate change scenarios include higher atmospheric CO_2 concentrations, higher temperatures and changes in precipitation (Adams et al., 1998; Trenberth et al., 2007).

Climate change will impact many aspects of the human and natural world (IPCC 2007), but effects on agricultural production could be of particular significance (Cline, 1992). According to Cline (1992) estimates of annual damages in agriculture due to temperature increase, for example, will be more costly to US economy compared to those in other sectors and activities such as forestry, electricity, water availability or water pollution, air pollution, human mortality and morbidity, leisure activities, migration, human amenities and urban infrastructure. The multifaceted climate alterations necessitates the adaptation of crop plants to tolerate increased heat, extended drought periods (Figure 1a) (Gala Bijl and Fisher, 2011), or increased flooding in tropical places. Additionally, the expected changes in the distribution, abundance, and severity of pests and weeds (Bazzaz and Carlson, 1984; Ziska and Runion 2007; Ziska 2014) will affect cropping systems and pest control methods (Anonymous, 2008). Although climate changes

120 compel agriculture to be adequately productive (Tokatlidis, 2013), its effects on agricultural
121 production can be positive in some agricultural systems and regions and negative in others
122 (Gregory et al., 2005; Obirih-Opareh and Adwoa Onumah, 2014).
123 To counteract the effects of climate change various adaptation strategies have been suggested.
124 These, according to IPCC (2014), are the processes of adjustment to actual or expected climatic
125 changes and its effects. In agricultural production systems adaptations seek to lessen or avoid
126 damages caused by climate changes or exploit beneficial opportunities (IPCC, 2001; Adger et al.,
127 2002). Farmers, throughout history, responded to changes in environment by adopting new crop
128 cultivars and by adjusting their cultural practices (Gala Bijl and Fisher, 2011). At the farm level,
129 these adaptations include alterations in planting and harvest dates, changes in cropping sequence,
130 better management of water for irrigation, optimized use of fertilizers and adoption of various
131 tillage practices (Adams et al., 1998). In addition, studies in Australia showed that crop
132 responses to climate change are strongly cultivar-dependent (Wang et al., 1992). Asfaw and
133 Lipper (2011) predicted that the farmers' primary response to climate change would be to seek
134 and crop cultivars that are most adapted to highly variable, extreme climatic conditions and pest
135 changes brought forth by global warming.
136 Weed interference, in addition to climate change, enhances the risk for further crop yield losses.
137 Despite the advanced technological achievements for weed control, crop yields are suffering
138 great losses due to weed competition (Figure 1b). Overall, weeds caused the greatest potential
139 loss (34%), with animal pests and diseases being, usually, less important (losses of 18 and 16%)
140 (Oerke, 2006). Competitiveness, adaptation, and stress tolerance are the characteristics by which
141 weed species secure their survival in a variety of environmental conditions. Competitiveness,
142 within the context of this paper, pertains to the ability of an organism (weed species in this case)

143 to perform better in acquiring resources in relation to another organism (crop plants) within the
144 same habitat. Adaptation is a change or a process of change by which an organism becomes
145 better suited to a 'new' environment whereas tolerance is the ability of an organism to survive
146 and reproduce under adverse environmental conditions. Weediness, which comprises traits that
147 secure the survival and dispersal of weeds, even under severe environmental conditions, can be
148 described through various morphological, phenological, or physiological characteristics. One of
149 the main components of integrated weed management strategies for farmers is to grow crops able
150 to offset the competitive ability of weeds. The utility of crops with weed-suppressive ability
151 particularly in low input agricultural systems, or in situations when chemical weed control is not
152 possible, can be proved valuable (Gibson et al., 2003; Benaragama et al., 2014). However,
153 selection for weed-suppressive cultivars is difficult because this trait is a manifestation of the
154 joint activity of many genes, controlling many traits. As reports have shown, a combination of
155 characteristics, instead of a single trait, interact for enhanced weed- suppressive ability (Andrews
156 et al., 2015). These traits are related to: a) crop morphological performance at early stages (i.e.
157 rapid emergence, rapid root and shoot growth, early groundcover, early biomass accumulation,
158 rapid leaf area development); b) crop growth characteristics (i.e. height, growth habit, tillering
159 ability, leaf width, maturity date); c) crop physiological performance (i.e. ability for efficient
160 water and nutrients uptake; and d) potential allelopathic properties (Korres and Froud-Williams,
161 2002; Korres, 2005; Mason and Spaner, 2006).
162 Climate change, in combination with an increasing world population, is predicted to escalate the
163 global need for farmland, a resource that is already in high demand (Barrow et al., 2008)
164 dwindling rapidly. The adoption of stress-tolerant cultivars that can withstand adverse climatic
165 changes and produce high yields is an effective strategy against the unprecedented risks of

climate change on crop productivity (Ciais et al, 2005) and the increasing demand for higher food production (Larson, 2013) particularly in low-input farming systems that are common in marginal areas (Darwin and Kennedy, 2000). Furthermore, stress-tolerant cultivars that exhibit attributes of increased suppressive ability against weeds would secure yield production even more either directly by dominating over weeds or indirectly by reducing crop management inputs (Korres and Froud-Willliams, 2002). To our knowledge, information that enables the evaluation of the relative strengths and weaknesses of both crops and weeds under various climate change scenarios is negligible. This paper aims to cover this gap and to discuss the benefits of selecting stress tolerant cultivar as a tool for integrated weed control under various climate change scenarios.

2.0 Effects of climate change on crops

2.1 Effects of elevated CO_2

2.1.1 Effects of elevated CO_2 on crop physiological characteristics

Increasing levels of atmospheric CO_2 due to various anthropogenic activities will directly influence photosynthesis, transpiration, and respiration, the main processes by which elevated CO_2 can be sensed directly by the plants and ecosystems (Drake et al., 1997). C3 and C4 plant types exhibit different responses to CO_2 enrichment. The current amount of CO_2 in the atmosphere is inadequate to saturate the ribulose 1, 5-biphosphate (RuBisCO) enzyme that drives photosynthesis in C3 plants (Taiz and Zeiger, 1991; Chijioke et al., 2011). Therefore future increases in CO_2 concentrations up to 57% by 2050 (Hulme, 1996), or even at higher levels (600-800 ppm) (Schmidhuber and Tubiello, 2007), will most probably favor C3 plant types (Table 1). In contrast C4 type plants are likely to respond less to elevated CO_2 levels as they possess an innate concentrating mechanism that increases CO_2 level at the site of RuBisCO to 2000 ppm.

189 Hence, predicted increases in atmospheric CO_2 concentrations, from a current ambient level of
190 about 370 ppm, are less relevant to the photosynthetic capacity of C4 plants which, most
191 probably, will respond only marginally (Poorter and Navas, 2003). The association of
192 photosynthesis rate and intercellular CO_2 concentration was compared in soybean (C3) and
193 maize (C4). Photosynthesis in soybean was stimulated by 39% under elevated CO_2 concentration
194 but not in maize (Leakey et al., 2009).

195 *2.1.2. Effects of elevated CO_2 on crop yields*

196 Carbon dioxide is fundamental for plant production, and increases of atmospheric CO_2
197 concentrations have the potential to enhance the productivity of agro-ecosystems (Table 1)
198 (Adams et al., 1998). Elevated CO_2 is expected to increase plant yield through root mass and leaf
199 area increases (Table 1) and to alter plant chemical composition, hence the rate of nutrient
200 cycling in soil (Campbell et al, 1997). Increases in marketable yield of cereals, particularly those
201 that exhibit C3 photosynthetic pathway, range between 8-70%, these of row, cash and vegetable
202 crops between 20-144% and these of flowers between 6-35%. The quality of agricultural
203 products may be altered also by elevated CO_2. Nitrogen content, for example, in some non-
204 nitrogen fixing plants grown at elevated CO_2, was found reduced (Ainsworth and Long, 2005;
205 Erbs et al., 2010). These changes could affect the nutritional value, taste, and storage quality of
206 some fruits and vegetables (Chijioke et al., 2011; Vermeulen et al., 2012).

207 **2.2 Effects of temperature increases**

208 *2.2.1 Effects of temperature increases on crop physiological characteristics*

209 Temperature increases result in altered phenology of leaf development, flowering, harvest and
210 fruit production, decreased vernalisation period, and in asynchrony between flowering and
211 pollinators (Baldocchi and Wong 2008). In addition, increased temperatures result in higher

212 respirations rates, shorter seed formation periods, and lesser biomass production; hence, lower
213 yields (Stone and Nicolas, 1995; Adams et al., 1998). Key stages of crop development, seasonal
214 temperature incidents, day-night temperature fluctuations and geographical scale are the major
215 parameters that should be taken under consideration when the effects of temperature on crop
216 yields are evaluated. Only few days of extreme temperatures at the flowering stage can
217 drastically reduce yield in many crops (Wheeler et al. 2000). Pre- and post-anthesis heat
218 incidents at 35 ºC led to significant yield loss of barley, wheat, and triticale (Zheng et al., 2002;
219 Porter and Semenov, 2005; Ugarte et al., 2007). Increases in spring temperatures have been
220 shown to induce earlier spring flowering (Pope et al., 2013), reductions in pollen germination,
221 flowering and ovule size with subsequent fruit yield declines due to smaller, deformed and fewer
222 fruit production in perennial crops (Pope, 2012; DeCeault and Polito, 2008). Each crop species
223 exhibits an optimal temperature for vegetative growth with growth decreasing as temperatures
224 diverge from this optimum. Similarly, there is a range of temperatures within which a plant will
225 set seeds and outside of which the plant will not be able to reproduce. Maize, for instance, will
226 fail to reproduce at temperatures above 32 ºC and soybean above 38 ºC (Figure 2).
227 Consequently, the trend in India toward more production of wheat, rice, and barley, and less
228 production of maize and millets, is likely to accelerate, whereas in the USA, production might
229 shift away from maize into soybean (C3) for forage (Parry, 1990). High temperatures (above 35
230 °C) in combination with high humidity and low wind speed caused a 4 °C-increase in rice
231 panicle temperatures, resulting in floret sterility (Tian et al., 2010).
232 *2.2.2 Effects of temperature increases on crop yield*
233 Crop yields particularly these of temperature-sensitive crops such as maize, soybean, wheat, and
234 cotton (Schlenker and Roberts, 2009), or specialty crops such as almonds, grapes, berries, citrus,

or stone fruits (Lobell and Field, 2011; Lobell et al., 2006) will be decreased with temperature increases at regional and local scale (Bonfils, 2012; Lobell et al., 2006). Night temperature increases resulted in rice and wheat grain yield losses (Lobell et al., 2005; Peng et al., 2004; Mohammad et al., 2009). Thus, even a C3 crop like rice which is expected to yield better under increased CO_2, will suffer serious yield losses under high temperature. Since the majority of global rice is grown in tropical and semi-tropical regions, it is likely that higher temperatures would negatively affect its production in these areas due to an increase in floret sterility that would subsequently decrease yields (Prasad et al. 2006a, b). The detrimental effect of high temperature on rice yield will be exacerbated by increased CO_2 in the atmosphere.

2.3 Effects of water deficit

2.3.1 Effects of water deficit on crop physiological characteristics

Physiological responses of plants to drought stress are complex and vary with plant species and the degree or time of the exposure to drought (Bodner et al, 2015; Evans et al., 1991). Under drought conditions, photosynthesis inhibition occurs because of stomata closure and reductions in the CO_2:O_2 ratio in leaves (Jason et al., 2004).

2.3.2 Effects of water deficit on crop yield

Significant crop yields reductions occur under drought stress through dry weight accumulation reductions in all plant organs and shorter plant life cycles (Blum, 1996). Pace et al. (1999) recorded significantly fewer nodes, lower dry weights of stems and reduction in height and leaf area between water-stressed and well-watered cotton plants (Table 3). In addition, water deficit at flowering may limit the viability of pollen, the receptivity of its stigma, and seed development (Blum, 1996). Reduced yields, especially in rain-fed cropping systems, is the norm under drought conditions (Kramer, 1983), the severity of which may increase due to changing world

climatic trends (Le Houerou, 1996). One possible scenario is that the irrigated wetland rice (13 Mha of cultivated land) in Asia may experience physical water scarcity by 2025, while the irrigated dry-season rice (22 Mha of cultivated land) may suffer economic water scarcity (human, institutional, and financial capital limit access to water even though water in nature is available locally to meet human demands) (Tuong and Bouman, 2003). Deleterious effects of water deficit on crops such as tomato (Ragab et al., 2007), soybean (Sakthivelu et al., 2008; Hamayun et al., 2010), maize (Khodarahmpour, 2011) and many others is well known.

2.4 Interactive effects of climate change components on the physiology of crop plants and yield

In previous sections the effects of climate change components on crop plants were examined individually although environmental changes occur concurrently (Albert et al., 2011) with management practices (Tubiello and Ewert, 2002). For instance, crop yield response to elevated CO_2 levels is relatively greater in rain-fed than in irrigated crops, due to a combination of increased water-use efficiency (Table 4) and root water-uptake capacity (Tubiello and Ewert, 2002). In addition, the projected increases in atmospheric CO_2 concentration will increase crop growth and consequently nitrogen uptake by the crop, thus potentially will increase the need for fertilizer applications if production is to be maximized (Olesen and Bindi, 2002). Elevated CO_2 resulted in a sustained larger N pool in above-ground biomass of grasses during a 5-year study on long-term enhancement of N availability under CO_2 concentration increases, suggesting that more N was taken up each year from the soil under elevated CO_2 (Dijkstra et al., 2008). Also, increased soil moisture under elevated CO_2 supported higher rates of N mineralization, thereby reducing N constraints on plant growth. More of the mineralized N ended up in the above-ground biomass of needle-and-thread [*Hesperostipa comata* (Trin. & Rupr.) Barkworth] (C3) than in

281 blue grama [*Bouteloua gracilis* (Willd. ex Kunth) Lag. ex Griffiths] (C4) under elevated CO_2
282 (Dijkstra et al., 2008). Therefore is possible that C3 species exhibit a higher plant N acquisition
283 and utilization under elevated CO_2 concentrations. Ghannoum (2009) reviewed the C4
284 photosynthesis response to water stress in interaction with CO_2 concentration and reported that
285 elevated CO_2 concentration lessens the deleterious effect of drought on plant productivity. This is
286 due to reduced stomatal conductance, CO_2 assimilation rate, and intercellular CO_2 levels
287 (Ghannoun, 2009; Ripley et al., 2007) therefore, saturating CO_2 concentration keeps the
288 photosynthetic capacity unchanged and limits reductions in plant productivity.

289 **3.0 Effects of climate change on weeds**

290 Compared with crops, weeds have more variable characteristics as they have not been subjected
291 to the same degree of selection for specific favorable traits (e.g. lack of seed dormancy, uniform
292 growth, high yields). Hence, weeds tend to exhibit greater potential capability to adapt to stress
293 than crop plants. The high genetic diversity among weedy plants allow them to achieve a greater
294 competitive fitness against crops as a consequence of climate change (Dukes and Mooney,
295 1999). The major categories under which climate change will affect weed populations include
296 species abundance and richness, geographic range, and phenology (Anonymous, 2013; Curtis
297 and Wang, 1998).

298 **3.1 Effects of elevated CO_2**

299 *3.1.1 Direct effects of elevated CO_2 on weeds*

300 There is an acknowledged consensus regarding the direct impact of increased CO_2 on plant
301 physiology (Ziska, 2004). Many weeds respond positively to elevated CO_2 due to decreased
302 stomatal conductance (Bunce, 1998) and subsequent improvements in water-use efficiency
303 (Patterson et al., 1999; Ziska and Runion, 2006). C3 plant types are likely to respond more

304 strongly than C4 plant types to CO_2 increases (Southworth et al., 2002; Ziska, 2004) (Figure 3)
305 through biomass and leaf area increases (Walthall et al., 2012). Nonetheless, results from various
306 studies indicate significant and wide variations in response to elevated CO_2 due to interactions
307 with temperature, light, water, and nutrients. CO_2 enrichment enhanced the growth and biomass
308 production of annual fescue [*Vulpia myuros* (L.) C.C. Gmel.] (C3 type), Santa Maria feverfew
309 weed (*Parthenium hysterophorus* L.) (C3/C4 intermediate type), and green amaranth
310 (*Amaranthus viridis* L.) (C4 type)(Scott et al., 2014; Naidu and Paroha, 2008). Other direct
311 effects of elevated CO_2 is the production of excess pollen in ragweed (*Ambrosia artemisiifolia*
312 L.) (Wayne et al., 2002) and the accelerated maturity rate in wild oat (*Avena fatua* L.)
313 (Anonymous, 2008).
314 *3.1.2 Indirect effects of elevated CO_2 on weeds*
315 Weed reproductive capacity will most probably be enhanced by increased CO_2 (Patterson et al.,
316 1999; Ziska and Runion, 2006). In case of the green amaranth, a 274 percent increase in flower
317 production under elevated CO_2 (550 ± 30 ppm) in controlled environmental conditions was
318 reported by Naidu and Paroha (2008). Reproductive capacity is linked to resource capture
319 (DeFelice et al., 1988; Benvenuti and Steffani, 1994; Bello et al., 1995) which is related to
320 increased biomass and leaf area (Korres, 2005). Therefore, increases in biomass with elevated
321 CO_2 levels will enhance weed reproductive output as these two traits are positively correlated
322 (Korres and Froud-Williams, 2002; Korres et al., 2015). Hence, increases in reproductive output
323 will result in increases of weed abundance. Disruptions of soil and native plant populations for
324 urban or rural development, emissions that increase atmospheric CO_2 concentrations, and
325 nitrogen deposition to the ground surface which enhance weed growth (Johnson and California
326 Invasive Plant Council, 2013), and roadside activities which lead to the spread of weeds (Korres

et al., 2015) will further enhance weed abundance. In addition, Ziska et al. (2004) observed that elevated CO_2 concentrations increased root biomass of Canada thistle (C3 plant type), suggesting that perennial weeds might be more difficult to control at these higher CO_2 levels.

3.1.3. Effects of elevated CO_2 on crop-weed interference

Some of the world's most troublesome weed species are C4 types and are found in C3 crops (Edwards and Huber, 1981). The positive response of C3 crops to increased CO_2 may make such weeds less competitive (Table 5). In contrast, C3 weeds in C4 or C3 crops, particularly in tropical regions, could become a problem (Table 5), although the final outcome will depend on other climate change components (Morison, 1989). Despite the fact that many weed species exhibiting a C4 photosynthetic pathway show less response to atmospheric CO_2 relative to C3 crops, in most agronomic situations, a mix of both C3 and C4 weeds occurs. As stated earlier increases in CO_2 concentrations will enhance C3 weed growth particularly for those species that reproduce by vegetative means (Ziska and George, 2004; Ziska, 2003). Consequently, the abundance of perennial weeds such as common couch [*Elytrigia repens* (L.) Desv. Ex. Nevski], heartshape pickerelweed [*Monocharia vaginalis* (Burm. F.) Presl], cosmopolitan bulrush [*Scirpus maritimus* L.], hedge bindweed [*Calystegia sepium* (L.) R. Br.], Canada thistle [*Cirsium arvense* (L.) Scop], perennial sowthistle [*Sonchus arvensis* L.], horsenettle [*Solanum carolinense* L.], most of them found in rice or soybean cropping systems, may increase, since elevated CO_2 stimulates greater rhizome and tuber growth (Chandrasena, 2009).

3.2 Effects of temperature

3.2.1 Effects of temperature on weed physiological characteristics

Soil temperature is the primary determinant of seed germination and survival particularly when soil freezes (Zimdahl, 2007). Various responses to temperature fluctuations have been reported

for seed germination of weed species. Common chickweed (*Stellaria media* L.) survives well in cold climates (King, 1966), whereas some of the most troublesome weeds in soybean, maize, and cotton respond to temperature gradients to varying degrees (Ehleringer, 1983). Barnyardgrass (*Echinochloa* spp.) is a weed of warm regions that requires high temperatures for dry matter production and growth (Maun and Bennett, 1986). Similarly prickly sida (*Sida spinosa* L.) needs high temperatures for its development (Anonymous, 2001). The spatial distribution of johnsongrass [*Sorghum halepense* (L.) Pers.] in colder climates is restricted by its rhizome intolerance to temperatures below -3 ^{o}C (Warwick and Black, 1983). Similarly, morningglories are frost intolerant (Halvorson and Guertin, 2003; Zia Ul-Haq et al., 2012) but their germination occurs over a wide range of temperatures (15-35 ^{o}C) (Cole and Coats, 1973-cited in Halvorson and Guertin, 2003) with optimum germination temperature at 24 ^{o}C (Crowley and Buchanan, 1980-cited in Halvorson and Guertin, 2003). In addition, Ziska et al. (2007) reported 88% increase in biomass and 68% increase in leaf area of itchgrass [*Rottboelliia cochinchinensis* (Lour.) W.D. Clayton] in response to a 3 ^{o}C-increase in temperature.

3.2.2 Effects of temperature on weed distribution

The geographical range of many weed species is largely determined by temperature and it has long been recognized that temperature determines successful colonization of new environments by weedy species (Woodward and Williams, 1987). Warming will affect the growth, reproduction and distribution of weeds. Increased temperatures could, for example, alter the latitudinal distinction between Midwest and Midsouth regions within the USA, altering the weed geographical limitations. The greater soybean and maize losses experienced in the Midsouth are associated with a number of very aggressive weed species of tropical or sub-tropical environments such as prickly sida and johnsongrass (Osunsami, 2009; Riar et al., 2013).

373 Obviously, increased temperatures will facilitate the spread of these species into other areas of
374 the Midwest with subsequent effects on soybean and maize production (Walthall et al., 2012).
375 Temperature increases are likely to be particularly important in affecting the relative plant
376 growth of C3 and C4 plants, potentially favouring C4 weeds (Dukes and Mooney, 1999), such as
377 smutgrass (*Sporobolus indicus* L. R. Br.). This again could provide suitable conditions for more
378 robust growth of some species, which are currently limited by low temperatures, whereas the
379 distribution of some tropical and sub-tropical C4 species could shift northwards (Ziska and
380 Runion, 2006; Chandrasena, 2009), thus exposing temperate-zone agriculture to previously
381 unknown aggressive colonizers.
382 In addition, Ziska et al. (2007) stated that an expansion of invasive weed species such as
383 itchgrass, cogongrass [*Imperata cylindrical* (L.) P. Beauv.] and witchweed [*Striga asiatica* (L.)
384 Kuntze] will be facilitated by temperature increases. They also reported an increase in biomass
385 and leaf area of itchgrass by 88 and 68% respectively in response to a 3 ^{o}C-increase. On the
386 contrary, additional warming could restrict the southern range of other cooler-climate invasive
387 weeds such as wild proso millet (*Panicum miliaceum* L.) or Canada thistle (Ziska and Runion,
388 2007).

389 **3.3 Effects of water deficit**

390 *3.3.1 Effects of drought on weed physiological responses*

391 Under more frequent and severe drought stress events due to climate change, the competitive
392 balance would shift in favor of deep-rooted plants (Stratonovitch et al., 2012). Early emerging
393 species, such as the shallow-rooted Sandberg's bluegrass (*Poa sandbergii* Vasey), which uses the
394 resources that are available in the upper soil profile early in the growing season and during
395 periods of light precipitation, will be suppressed (Daudenmire, 1970 cited in Sheley et al., 1996).

In addition, dry soil conditions prolong the longevity of weed seeds due to unfavorable conditions for seed predators (Storrie and Cook 2007) and unfavorable conditions for germination. Weed seeds such as black bindweed (*Polygonum cilinode* Michx.) can last up to seven years in the soil under dry conditions (Storrie and Cook, 2007). A summary of the potential impacts of drought stress on some of the most important Australian weeds are shown in Table 6 where a trend of establishment in higher latitudes is expected (Anonymous, 2008).

3.3.2 Weed adaptation strategies under water deficit and other unfavorable conditions

As reported by Wiese and Vandiner (1970), species with greatest growth under high soil moisture conditions will be the most adversely affected by the combination of competition and water shortage. On the contrary, the more competitive species under semi-drought conditions are likely to be those that produce little growth in moist soils. Based on the competitive exclusion principle, the species that uses a resource more efficiently will eventually, either wholly or partially, displace the other species. This opportunistic behavior characterizes the r-strategists, those with short life cycle and high energy investments into reproduction and dispersability, as opposed to K-strategists (Sheley et al., 1996; Hardin, 1960). Grime (1979) extended the r- and K-classification strategies into stress tolerators (S), competitors (C), ruderals (R) or combinations of the above strategies. Under high stress intensity that can limit plant growth, as in the case of water or nutrient shortage, stress tolerators (S) can perform adequately. Based on the ability of adjacent organisms to exploit the same resource competitors (C) will perform best whereas ruderals (R) can withstand physical damages. Most weeds of annual agricultural systems exhibit ruderal-competitive characteristics, whereas most weeds of rangeland and forest ecosystems exhibit stress tolerance-competitive characteristics. Typically, succession is evolved from ruderal to competitive and finally to stress tolerator species (Korres, 2005). Hence, under water- or

nutrient shortage scenarios, an r-strategist with characteristics in the order S-C-R, will most probably prevail. In a recent weed survey (Korres et al., 2015), the preference of Palmer amaranth, large crabgrass (*Digitaria sanguinalis* L. Scop.), johnsongrass, and spurges (*Euphorbia* spp.) for disturbed habitats was reported. In the same survey, giant ragweed (*Ambrosia trifida* L.), yellow nutsedge (*Cyperus esculentus* L.), barnyardgrass, and hemp sesbania [*Sesbania herbacea* (Mill.) McVaFugh] exhibited a strong preference for moist habitats. Obviously, the former group of weeds is assured of a greater probability for survival under water or nutrient stress conditions in comparison to the latter.

3.4 Interactive effects of climate change components on weed performance and consequences on weed-crop competition

The influence of climate change on simple competitive outcomes will be difficult to predict based simply on a single model, as interactions between the various climate change scenarios are likely to concur and will affect the outcome of the crop-weed competition (Alberto et al., 1996). The growth of a tropical weed is strongly stimulated by relatively small changes in air temperature (Patterson et al., 1984), but the potential synergistic effects of rising CO_2 on these weeds relative to tropical crops is unknown. It is believed that increased CO_2 and temperature can negatively impact plant growth. Scott et al. (2014), for example, reported that increases in both parameters negatively impacted plant growth rates in grassland ecosystems. The effects of elevated CO_2 levels on crops and weeds will alter the weed-crop competitive interactions, sometimes for the benefit of the crop and sometimes for the weeds. Consequently, the control of weeds will also likely be affected by these changes (Patterson, 1995; Coakley et al., 1999). Reduction in transpiration and changes in leaf anatomy and leaf surface characteristics, or greater root to shoot ratio caused by elevated CO_2, could also affect herbicide uptake, thus reducing

herbicide efficiency (Patterson et al., 1999; Olesen and Bindi, 2002; Poorter and Navas, 2003; Dukes et al., 2009). This was confirmed by various studies in which increased CO_2 concentration has affected the efficacy of glyphosate on both C3 and C4 weed photosynthetic types (Ziska et al., 2004; Manea et al., 2011; Ziska et al., 1999) (Table 7). This response to carbon dioxide in combination with the evolution of glyphosate resistance by many weed species (Heap, 2015), will affect weed control schemes significantly. Controlling weeds currently costs the United States, more than $11 billion a year, with the majority spent on herbicides, hence both herbicide use and costs are likely to increase as temperatures and carbon dioxide levels rise (Karl et al., 2009).

Additionally, little attention has been focused on the interactions between nutrient availability or drought with rising CO_2, on weed-crop competition. According to Newton et al. (1996) the proportion of weed biomass increased with elevated CO_2 equally in wet and dry treatments in pasture mixture. In another study, reduced weed competition was observed when tomato (C3 crop) and redroot pigweed (C4 weed) were grown under well-watered conditions, but when drought and high CO_2 occurred synchronously, redroot pigweed performed better (Valerio et al., 2011). Under extreme nutrient limitations, stimulation of biomass with additional CO_2 may be minimal. However, under moderate nutrient limitations, more indicative of agroecosystems, the increase in biomass may be reduced but still occurs (Seneweera et al., 1994). Under a competitive environment between rice (C3 crop type) and barnyardgrass (C4 weed type), the proportion of rice biomass increased relative to barnyardgrass with a 200 ppm increase in atmospheric CO_2, but only when soil nitrogen was adequate. If nitrogen was limited in an enriched CO_2 environment, the competitive ability of rice relative to barnyardgrass was reduced, possibly due to reductions in tiller formation (Zhu et al., 2008). Elevated CO_2 can mitigate some

of the adverse effects of increased temperature and drought and also regulate the adaptive mechanism of black knapweed (*Centaurea nigra* L.) (Qaderi et al., 2013). The effects of drought are likely to vary widely among crops and weeds. In maize, drought has been found to both decrease interference from naturally occurring weed flora dominated by foxtail species (*Setaria* spp.) (McGiffen et al., 1997), and increase the competitive ability of johnsongrass (Leguizamon, 2011). Drought and high temperatures favor the competitive ability of C4 weeds over C3 crops (Fuhrer, 2003), an advantage which will most probably diminish or possibly be reversed under increased CO_2 concentrations (Bazzaz and Carlson, 1984; Carter and Peterson, 1983). Spatial-based effects of temperature increases and prolonged drought periods on weeds have also been anticipated. More particularly, long drought periods interspersed with occasional very wet years will enhance weed invasion because established vegetation, both native and crops, will be weakened, leaving some areas open to invasion (Chandrasena, 2009). In general, wetter and milder winters are likely to increase the survival of some winter annual weeds, whereas warmer summers and longer growing seasons may permit thermophile summer annuals to grow in regions further north (Peters et al., 2014). Alterations in temperature and nutrients supply can reduce photosynthetic rate of Palmer amaranth. The combination of temperature between 36-46 ^{o}C with resource supply constraints may restrict the potential distribution range of Palmer amaranth (Ehleringer, 1983; Ward et al., 2013).

4.0 Cultivar selection against weeds and traits that confer competitiveness

Crop ability to suppress weeds can be considered in two ways, namely a) an ability to tolerate weed competition which can be measured by the ability of the crop to maintain high yields under weedy conditions, and b) the ability of the crop to suppress the growth of weeds, usually determined by comparing different biological characteristics in mixtures with that in pure stands,

known as weed suppression ability or competitive ability (Callaway, 1992; Korres, 2004 Andrews et al., 2015). However, there is a confusion between cultivar tolerance to weed competition and cultivar weed suppressive ability (Olesen et al., 2004). Furthermore, crop tolerance to weed competition varies widely over seasons and locations (Cousens and Mokhtari, 1998; Olesen et al., 2004). Thus, weed suppression criterion has been emphasized here for the selection of suitable cultivars against weeds under various climate change scenarios.

4.1 Cultivar phenotypic characteristics and weed suppression

Unlike breeding for diseases and pest resistance, little research has been done on breeding crop cultivars which are more competitive to weeds. Certain crop cultivars are known to be better competitors with weeds than others (Callaway, 1992). For example, white bean (*Phaseolus vulgaris* L.) cultivars differ in their ability to compete with weeds (Malik et al., 1993). Certain tomato cultivars (*Lycopersicon esculentum* L.) have considerable tolerance to dodder (*Cuscuta* spp.), a severe parasitic weed in many parts of the world (Goldwasser et al., 2001). Cultivars of small grain cereals with certain characteristics such as short stature, earlier maturity, better winter hardiness or early season growth have shown differential competitive abilities when grown in mixtures compared to monocultures (Juskiw et al., 2000). As stated by various authors breeding crop cultivars with an enhanced ability to suppress weeds would be a sustainable contribution to improved weed management in many crops (Didon and Bostrom, 2003; Lemerle et al., 2001; Paolini et al., 1998; Vollmann et al., 2010). Therefore, cultivar selection with traits that enhance its ability to suppress weeds such as these mentioned above could be explored under various climate change scenarios. Additionally, the belowground traits such as root length density, root elongation rate, total root length, and root spatial distribution are important factors for attributing competition effect (Gealy et al. 2013a; Fargione & Tilman, 2006; Stevanato et al.,

2011). The greater ability to extract water from dry soil may affect or even determine the competitive ability of a cultivar (Song et al, 2010). Reports have shown that under weed competition the root:shoot ratio of the crop and weeds was reduced (Kasperbauer and Karlen, 1994; Thomas and Alison, 1975; Stone et al, 1998), particularly of the less competitive species, although soil water content was not a limiting factor (Thomas and Alison, 1975; Rajcan and Swanton, 2001). However, as stated by Rajcan and Swanton (2001), competition for water should be viewed as an outcome of the interaction between both soil-plant-atmosphere and the crop-weed systems, rather than simply as a shortage of available water.

4.2 Implications for allelopathic properties

Weed suppression can vary with management factors such as planting method, seeding density, flood depth, and nitrogen fertilization whereas in some cases, activated charcoal has reduced the inhibition of weeds in soils, implicating allelopathic activity as a possible contributing factor (Kong et al., 2008; Kong et al., 2011). Rondo, for example, a rice cultivar grown in a commercial organic rice production operation in Texas, USA that combines a high yield potential and a weed suppression ability is considered as a potential cultivar with allelopathic properties (Gealy and Yan 2012). Bertholdsson (2010) bred spring wheat for improved allelopathic potential by conventional breeding. The material used originated from a cross between a Swedish cultivar with low allelopathic activity and a Tunisian cultivar with high allelopathic activity.

Therefore, research efforts have focused on combining allelopathic activity with other weed-suppressive traits in small grains such a rice (Figure 3). Breeders in Asia showed that allelopathic traits in rice can be quantitatively inherited (Chen et al., 2008) and weed-suppressive cultivars have now been developed in that region (Kong et al., 2011; Ma et al., 2006; Pheng et al., 2009a, 2009b). Similar progress has been reported in the USA (Gealy et al., 2013b). Breeding efforts

with other small grains in Europe, using a dual screening approach of seedling bioassays for allelopathic potential coupled with field evaluations for general weed suppression, have resulted in germplasm with improved weed suppression or tolerance (Bertholdsson 2005, 2007, 2010). It has been reported that early season crop biomass and allelopathic potential were key traits for improved weed suppression by the crop (Bertholdsson, 2011; Bertholdsson et al., 2012). Worthington and Reberg-Horton (2013) have reviewed important breeding issues for small grains associated with optimization of weed-competitive ability and allelopathic traits. Rice traits such as rapid seedling growth, leaf area, and tiller production, and high yield potential have improved weed suppression and minimized crop yield loss (Gealy and Moldenhauer, 2012; Gealy and Yan, 2012; Gibson et al., 2003; Pérez de Vida et al., 2006). Zhao et al. (2006) successfully selected cultivars for weed-suppressive traits such as yield, early vigor, and height under weed-free conditions to identify weed- competitive cultivars.

5.0 Traits for developing an ideotype S-C cultivar

The use of tolerant cultivars to a wide range of climatic fluctuations as adaptive tool is widely spread (Matthews et al., 1994). In Australia, for example, the use of late maturing cultivars secures high yield outcomes that could be otherwise affected by inconsistent climatic conditions (Connor and Wang, 1993). A similar strategy, at crop level, is used in Canada and China, where the diversification of crops counteracts the climatic fluctuations (Hulme et al., 1992; Cohen et al., 1992).

5.1 *Cultivars with deep root system*

Adapting a cultivar with a deep root system, particularly in areas which experiencing prolonged dry periods can be a useful tool (Bodner et al., 2015). Newly introduced wheat cultivars can better exploit water and nutrients (Korres et al., 2008) mainly due to their greater ability to

maintain water uptake and consequently to survive longer in dry soils (Song et al., 2009). Sorghum, for example, seems an attractive option for dry lands where crops frequently encounter drought stress compared to maize. Sorghum has deep root system, high root density, cuticle and epicuticular deposition in leaves, and efficient stomata function under water stress (Assefa et al., 2010; Starggenborg et al., 2008; Schittenhelm and Schroetterm, 2014). Traits related to competitiveness for water and nutrients that could affect the weed suppressive ability of the crop include root density, root length, water uptake rate and root surface area (Aarssen, 1989; Callaway, 1992; Mohler, 2001). In the long-term, breeding drought-tolerant cultivars might be advantageous for weed suppression as well as a means to cope with climatic changes in areas with prolonged summer dry periods (Bodner et al., 2015). Acquiring and utilizing water and nutrients more adequately compared to weeds due to their extensive root system, for example, will enable to crop cultivars to maintain growth even under drought conditions. Cultivars with high early vigor and earlier maturity can be used as an effective adaptation strategy for areas with semi-arid continental climates in temperate zones where more frequent generative droughts are forecasted (Gouache et al., 2012; Bodner et al., 2015). Genetic manipulation using molecular breeding has resulted in commercialization of drought-resistant crops such as the maize-DroughtGrade^{TM} (Monsanto, St. Louis, USA) that is already used extensively in the US (Waltz, 2014). Differences in resistance to drought are known to exist within genotypes of plant species (Grzesiak et al., 2012) e.g. in, wheat (Winter et al., 1988; Paknejad et al., 2007), rapeseed (Richards & Thurling, 1978), oat (Larsson and Gorny, 1988), and triticale (Royo et al., 2000; Grzesiak et al., 2012). Nevertheless, drought tolerance does not necessarily provide competitive advantages to the crop. As reported by Cerqueira et al. (2013), two drought-tolerant upland rice cultivars were affected by the competition of shrubby false buttonweed (*Spermacoce verticillata*

580 L.) regardless of water conditions (presence and absence). In addition, as reported by Chauhan
581 and Abugho (2013), rain-fed rice plants under weed competition with spiny amaranth
582 (*Amaranthus spinosus* L.) and Chinese sprangletop [*Leptochloa chinensis* (L.) Nees] (C4 types)
583 did not survive under limited water conditions. On the contrary both weed species, survived and
584 produced a significant number of tillers and leaves.

585 5.2 *Harvest index and dry mater components*

586 To promote adaptation to high temperatures, plant breeders have suggested phenotypic traits
587 related to heat tolerance during flowering, high harvest index, small leaves, and reduced leaf area
588 per unit of ground area (Walthall et al., 2012). Differences between winter wheat cultivars in
589 harvest index at high temperatures imply that heat-tolerant cultivars maintain higher grain
590 development, compared to more temperature-sensitive cultivars (Wardlaw and Moncur, 1995).
591 Lower harvest indices are an indication of injudicious investment of assimilates, a result of
592 favoring biomass production over commercial yield (Hay and Walker, 1992). Therefore,
593 genotypes with high harvest indices are expected to be weak competitors because of the relative
594 fewer resources allocated for stem and leaf expansion (Kawano and Jennings, 1983), traits that
595 confer competitiveness. Mann (1980) stated that it might be possible to obtain improvements in
596 harvest index and therefore yield, suggesting further reductions in straw length and maintenance
597 of above ground biomass. Korres (2000) investigating winter wheat cultivar characteristics for
598 increased competitive ability, found a negative relationship between number of leaf area/m^2 and
599 infertile tillers/m^2. Questions that merit further thought are related to the manipulation of leaf
600 area and infertile tiller production. If the production of infertile tillers could be manipulated,
601 would this result in leaf area investments? Would increases in leaf area, hence interception of
602 photosynthetic active radiation, in response to increased day length as the crop enters

603 reproductive development cause higher yield production and enhance competitive ability? Would
604 leaf area duration be affected and what would be the consequences for grain yield?
605 However, specific leaf area, a characteristic which is positively correlated with relative growth
606 rate, is usually reduced by elevated CO_2 thereby counteracting the positive response of
607 photosynthesis (Bruhn et al., 2001).
608 5.3 *Late maturing cultivars*
609 Late-maturing soybean cultivars (group IV) depressed weed seed production and seed weight of
610 both pitted morningglory (*Ipomoea lacunosa* L.) and hemp sesbania [*Sesbania exaltata* (Raf.)
611 Rydb. ex A.W. Hill] presumably through increased crop competitiveness (Bennet and Shaw,
612 2000) due to their ability to maintain vegetative growth longer (Nordby et al., 2002).
613 Nevertheless, Rosenzweig and Tubiello (2007) suggested that under warmer climates, crops
614 would tend to mature faster, resulting in less time available for carbohydrate accumulation and
615 grain production. Responses to specific adaptation strategies for given cropping systems can still
616 vary considerably, as a function of location and climate change scenario. Adapting longer-
617 maturing cultivars, in a winter cereal production system requires enough precipitation over an
618 extended growing season to sustain grain filling. If both warmer and drier conditions prevail,
619 such an adaptation strategy is not applicable. On the contrary, the adaptation of fast growing
620 species (i.e. those with high sink strength, hence positive response of photosynthesis) has the
621 advantage of better competition for resources thus faster adaptation to a changed climate.
622 5.4 *Nutrients uptake and utilization*
623 Nutrient utilization, mainly nitrogen, is an important factor for cultivar selection as an adaptive
624 strategy, but also as a crop competitiveness tools under various climate change scenarios. There
625 is a general agreement that crop cultivars, particularly of cereals, can differ in their

responsiveness to nitrogen (Gent and Kiyomoto, 1998; Duan et al., 2007; Benin et al., 2012) possibly due to greater sink capacity, hence better nitrogen utilization or more extensive root systems (Lupton et al., 1974; Foulkes et al., 1994). Crop biomass is a component of two processes namely the amount of accumulated intercepted radiation and radiation use efficiency (Monteith, 1977; Gallagher and Biscoe, 1978). Foulkes et al. (1994) stated that maximum growth depends on the acquisition of sufficient nitrogen to form a canopy of sufficient size to intercept the majority of the incident radiation when adequate moisture to balance evaporation from the canopy is provided. One of the main traits conferring resistance to drought in winter wheat is the flowering date (Foulkes et al., 1997). More particularly, cultivars with early flowering are less prone to drought effects due to shorter life cycle they exhibit. Susceptible cultivars to dry conditions, especially towards the end to the growing season, uptake and utilize lower nitrogen. Hence, cultivars with efficient N uptake and utilization that exhibit drought resistance characteristics can be used for weed suppression and also as adaptive tools in less fertile or dry soils.

5.5 *Heat tolerance-Improvements and expectations*

Improvements of heat-stress tolerant germplasm lines have resulted in the development of the Hoveyzeh rice cultivar from Khuzestan delta in south Iran which attains spikelet fertility at average day temperatures of 45 ºC (Jennings et al., 1979). Despite the impressive achievements by plant breeding programs, efforts to generate heat-tolerant crops have not been very successful. This is mainly because abiotic stress-tolerance in plants is quantitatively inherited and it is found to be controlled by multiple genes/quantitative trait loci (Blum et al., 1988). Advances in agricultural biotechnology have been successful in developing heat-tolerance transgenically under controlled conditions (Grover et al., 2013).

5.6 A *synthesis*

Breeding objectives should be re-orientated towards a selection of traditional × modern crop characteristics that will result in increased weed suppressive ability (Dingkuhn et al., 2010), an ability to thrive in harsh environments and high yielding potential (Jones et al., 1997; Johnson et al., 1998). Hybrids of *Oryza glaberrima* × *Oryza sativa* share common parental characteristics such as weed competitiveness, ability to grow under stressful conditions without jeopardizing their yield (Jones et al., 1997; Johnson et al., 1998).

Priority should be focused on crop traits suitable for climate change scenarios for several reasons. This is true considering the detrimental effects of increased temperature or extended drought periods on crop yields, for example, in combination with the enhanced plasticity and adaptation ability the weed species respond to various environment changes. If an appropriate trait for climate change adaptation favors the weed suppressive ability of the crop plant then its selection should be prioritized. The following Table (Table 8), in an attempt to facilitate the selection process, summarizes the major responses of both crop plants and weeds under various scenarios of climate change.

As mentioned earlier, increased temperatures will reduce vernalization (i.e. the promotion of flowering in response to a prolonged exposure to low temperatures) requirements for both crops and weeds, particularly grasses. This in turn will shorten the vegetative period due to early reproductive induction (Chauvel et al., 2002), at the vegetative points, of the apex (Chouard, 1960; Chauvel et al., 2002) which will result in biomass reductions for both crop plants and weeds and consequent yield reductions. As it was stated earlier increases in biomass production or its components e.g. leaf area, tillers, stem weight etc. are positively related to increased

671 competitiveness for both crop plants and weeds as in the case of cereals, particularly winter
672 wheat, and blackgrass [*Alopecurus myosuroides* Huds.] (Chauvel et al., 2002).

673 • Cultivars that retain appropriate vernalization periods under increased temperatures, hence
674 maintaining vegetative growth stages, can preserve yield production but also to exhibit
675 suppressive ability against weeds.

676 • The development of tolerant cultivars to drought with increased root:shoot ratio will result in
677 enhanced water and nutrient uptake, unaffected growth rates and biomass production, hence
678 improved weed suppressing ability.

679 • Traits related with the maturity of cultivars is another option that merits further consideration
680 for developing cultivars tolerant to drought and enhanced suppressive ability against weeds.

681 • Traits associated with the regulation of node formation and/or internode distance, particularly
682 under drought stress conditions, can be used for developing high yielding and competitive
683 cultivars against weeds.

684 • Traits or plant attributes related with harvest index variations such as these of infertile tillers
685 and leaf area as mentioned above merit further investigation since they can influence both
686 yield production through increased utilization of resources (i.e. PAR) and weed suppressive
687 ability (e.g. shading).

688 • Cultivars that exhibit allelophathic attributes should be prioritized in breeding programs.

689 • **7.0. Conclusions**

690 Climate change is predicted to affect agricultural production in many ways. Climate change is
691 likely to affect the growth of both crops and weeds, sometimes benefiting the crop sometimes the
692 weeds. Crop yield in many areas will decrease due to increased temperatures or extended
693 drought periods whereas weed competition, despite the technological advances, will increase

694 further crop yield reductions. A dual adaptive approach is needed not only to counteract the
695 negative effects of climate change but also to enhance crop competitiveness against weeds. As it
696 has been shown in this paper cultivar selection serves this adaptive approach adequately.
697 Cultivars with C3 photosynthetic pathway are more suitable for adaptation to elevated CO_2 but
698 also to compete with weeds, particularly those with C4 photosynthetic pathway. In addition,
699 cultivars with mechanisms to resist drought through increases in root:shoor ratio will gain a
700 significant advantage under dry conditions in marginal areas. The potential of these cultivars for
701 weed suppression will more likely enhance, due to their ability to acquire water and nutrients
702 effectively. However, increased temperatures, accompanied by extended drought periods, favour
703 the selection of cultivars with longer maturity period which have also proved to be highly
704 competitive by maintaining longer vegetative growth. Cultivars with allelopathic abilities should
705 be used in integrated weed management systems since they have shown great potential for high
706 yield production but also increased weed suppressing ability. This paper investigates the
707 complex interactions between crops and weeds under various climate change scenarios aiming to
708 facilitate decision -making processes towards sustainable crop production systems. Developing
709 cultivars to tolerate climate changes such as drought, temperature increases or nutrient shortage
710 can reduce fertilizer and irrigation inputs considerably. The incorporation of cultivars with
711 enhanced weed suppression ability into the system can reduce herbicide inputs substantially
712 (Callaway, 1992; Gealy et al., 2014; Gealy et al., 2003; Korres et al., 2008; Travlos, 2012). This
713 is even more demanding considering the increase of weed herbicide resistance evolution (Heap,
714 2015). The cumulative effects from selecting a suitable S-C cultivar will be reflected in
715 reductions of environmental pollution, lower production costs and sustainable food production. It
716 is therefore imperative to expand research efforts to investigate how crop-weed interference

under various abiotic stresses and cropping systems influences cultivar performance and subsequent yield outcome. This information could be incorporated into breeding programs for improving cultivars performance under abiotic (climate change) and biotic (weed competition) stresses without compromising final yield.

References

Aarssen LW (1989) Competitive ability and species coexistence: A 'plant's eye' view. Oikos 56, 386-401. doi: 10.2307/3565625

Adams RM, Hurd BH, Lenhart S, Leary N (1998) Effects of global climate change on agriculture: an interpretative review. Clim Res 11, 19-30. doi:10.3354/cr011019

Adger WN, Kelly PM, Winkels A, Huy LQ, Locke C (2002) Migration, remittances, livelihood and trajectories and social resilience in Vietnam. Ambio 31, 358-366. doi: 10.1579/0044-7447-31.4.358

Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air CO_2 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO_2. New Phytol 165:2, 351-372. doi: 10.1111/j.1469-8137.2004.01224.x

Albert KR, Ro-Poulsen H, Mikkelsen TN, Michelsen A, van der Linden L, Beier C (2011) Interactive effects of elevated CO_2, warming, and drought on photosynthesis of *Deschampsia flexuosa* in a temperate heath ecosystem. J Exp. Bot. 62:4253-4266. doi: 10.1093/jxb/err133

Alberto A, Ziska L, Cervancia C, Manalo P (1996) The influence of increasing carbon dioxide and temperature on competitive interactions between a C3 crop, rice (*Oryza sativa*) and a C4 weed *(Echinochloa glabrescens*). Funct Plant Biol 23:6, 795-802. doi: 10.107/PP9960795

Andrews IKS, Storkey J, Sparkes DL (2015) A review of the potential for competitive cereal cultivars as a tool in integrated weed management. Weed Res 55, 239-248. doi: 10.1111/wre.12137

Anonymous (2001) *Sida spinosa*. European and Mediterranean Plant Protection Organization. 02/9188, Point 7.8.

Anonymous (2008) Climate change impacts on pest animals and weeds. Communicating Climate Change. Module 13. Australian Government, Department of Agriculture, Fisheries and Forestry. Bureau of Meteorology.

Anonymous (2013) Climate change consortium for specialty crops: impacts and strategies for resilience. California Dept. of Food and Agriculture, 64 p.

Asfaw S, Lipper L (2011) Economics of PGRFA management for adaptation to climate change: A review of selected literature. Background study paper No 60. Commission on Genetic Resources for Food and Agriculture. Agricultural Economic Development Division (ESA), FAO, Rome, Italy.

Assefa Y, Staggenborg SA, Prasad VPV (2010) Grain sorghum water requirement and responses to drought stress: A review. Crop Manage 9:1. doi:10.1094/CM-2010-1109-01-RV

Baldocchi D, Wong S (2008) Accumulated winter chill is decreasing in the fruit growing regions of California. Clim Change 87:1, 153-166. doi: 10.1007/s10584-007-9367-8

Barrow JR, Lucero ME, Reyes-Vera I, Havstad KM(2008) Do symbiotic microbes have a role in plant evolution, performance and response to stress? Commun. Integr. Biol. 1, 69-73. doi:10.4161/cib.1.1.6238

Bazzaz FA, Carlson MR (1984) The response of plants to elevated CO_2. I. Competition among the assemblage of annuals at two levels of soil moisture. Oecologia 62, 196-198

Bello IA, Owen MDK, Hatterman-Valenti HM (1995) Effect of shade on velvetleaf (*Abutilon theophrasti*) growth, seed production, and dormancy. Weed Technol 9, 452-455

Benaragama D, Rossnagel DB, Shirtliffe SJ (2014) Breeding for Competitive and High-Yielding Crop Cultivars. Crop Sci 54, 1015-1025. doi: 10.1614/WS-D-10-00121.1

Benin G, Bornhofen E, Beche E, Pagliosa ES, da Silva CL, Pinnow C (2012) Agronomic performance of wheat cultivars in response to nitrogen fertilization levels. Acta Scientiarum. Agronomy 34:3, 275-283. doi: 10.4025

Benvenuti M, Steffani A (1994) Effects of shade on reproduction and some morphological characteristics of *Abutilon theophrasti* Medicus, *Datura stramonium* L. and *Sorghum halepense* L. Pers. Weed Res 34, 283-288. doi: 10.1111/j.1365-3180.1994.tb01996.x

Bertholdsson NO (2010) Breeding spring wheat for improved allelopathic potential. Weed Res 50:49-57. Doi: 10.1111/j.1365-3180.2009.00754.x

Bertholdsson NO (2005) Early vigour and Allelopathy-two useful traits for enhanced barley and wheat competitiveness against weeds. Weed Res 45:94-102. Doi: 10.1111/j.1365-3180.2004.00442.x

Bertholdsson NO (2011) Use of multivariate statistics to separate allelopathic and competitive factors influencing weed suppression ability in winter wheat. Weed Res 51:273–283. Doi: 10.1111/j.1365-3180.2011.00844.x

Bertholdsson NO (2007) Varietal variation in allelopathic activity in wheat and barley and possibilities for use in plant breeding. Allelopathy J 19:1. ISSN: 0971-4693.

Bertholdsson NO, Andersson SC, Merker A (2012) Allelopathic potential of *Triticum* spp., *Secale* spp. and *Triticosecale* spp. and use of chromosome substitutions and translocations to improve weed suppression ability in winter wheat. Plant Breeding 131:75-80. Doi: 10.1111/j.1439-0523.2011.01895.x

Blum A (1996) Crop responses to drought and the interpretation of adaptation. Plant Growth Regul 20, 135-148. doi: 10.1007/BF00024010

Bodner G, Nakhforoosh, and Kaul HP (2015) Management of crop water under drought: a review. Agron Sustain Dev 35, 401-442. doi: 10.1007/s13593-015-0283-4

Bruhn D, Mikkelsen TN, Pilegaard K, Gavito ME, Saxe H (2001) Climate change in a plant ecophysiological perspective. In Jorgensen AMK, Fenger J, Halsnaes K (eds) Climate change research. Danish contributions. Danish Meteorological Institute/Gads Forlag, Copenhagen, pp. 167-190.

Bunce JA (1998) Effects of humidity on short-term responses of stomatal conductance to an increase in carbon dioxide concentration. Plant, Cell and Environ 21, 115-120. doi: 10.1046/j.1365-3040.1998.00253.x

Brunce JA, Ziska LH (2000) Crop ecosystems responses to climatic change. Crop/weed interactions. In Reddy K. R. and Hodges H. F (eds) Climate change and global crop productivity. Cab International, Wallingford, New York, pp. 333-348. ISBN: 978-0851994390

Callaway MB (1992) A compendium of crop varietal tolerance to weeds. Am. J. Alternative Agr. 7, 169-180. doi: http://dx.doi.org/10.1017/S088918930000477X

Campbell BD, Stafford Smith DM, McKeon GM (1997) Elevated CO_2 and water supply interactions in grasslands: A pastures and rangelands management perspective. Global Change Biology 3, 177-187. doi: 10.1046/j.1365-2486.1997.00095.x

Carter DR, Peterson KM (1983) Effects of CO_2 enriched atmosphere on the growth and competitive interaction of a C3 and C4 grass. Oecologia 58, 188-193. doi: 10.1007/BF00399215

Cerqueira FB, Erasmo EAL, Silva JIC, Nunes TV, Carvalho GP, Silva AA (2013) Competition between drought-tolerant upland rice cultivars and weeds under water stress condition. Planta Daninha 31:2, 291-302. doi: http://dx.doi.org/10.1590/S0100-83582013000200006

Chandrasena N (2009) How will weed management change under climate change? Some perspectives. J Crop Weed 5:2, 95-105.

Chauvel B, Munier-Jolain NM, Grandgirard D, Gueritaine G (2002) Effect of vernalization on the development and growth of Alopecurus myosuroides. Weed Res 42:166-175

Chen XH, Hu F, Kong CH (2008) Varietal improvement in rice allelopathy. Allelopathy J 22:379-384.

Chijioke OB, Haile M, Waschkeit C (2011) Implication of Climate Change on Crop Yield and Food Accessibility in Sub Saharan Africa. Interdisciplinary Term Paper, University of Bonn, Germany. Global Climate Adaptation Partnership (GCAP) (2012). Projects and Clients. http://www.climateadaptation.cc/projects-clients [Accessed September 18, 2015]

Chouard P (1960) Vernalization and its relations to dormancy. Annu. Rev. Plant. Physiol. 11:191-238

Ciais P, Reichstein M, Viovy N, Granier A, Ogee J, Allard V, Aubinet M, Buchmann N, Bernhofer C, Carrara A, Chevallier F, De Noblet N, Friend AD, Friedlingstein P, Grunwald T, Heinesch B, Keronen P, Knohl A, Krinner G, Loustau D, Manca G, Matteucci G, Miglietta F, Ourcival JM, Papale D, Pilegaard K, Rambal S, Seufert G, Soussana JF, Sanz MJ, Schulze ED, Vesala T, Valentini R (2005). Europe-wide reduction in primary productivity caused by the heat and drought in 2003. Nature 437, 529-533. doi:10.1038/nature03972

Cline WR (1992) The economics of global warming. Washington, DC: Peterson Institute for International Economics. ISBN: 978-0-88132-132-6

Coakley SM, Scherm H, Chakraborty S (1999) Climate change and plant disease management. Annu Rev Phytopathol 37, 399-426. doi: 10.1146/annurev.phyto.37.1.399

Cohen S, Wheaton E, Masterton J (1992) Impacts of Climatic Change Scenarios in the Prairie Provinces: A Case Study from Canada. SRC Publication No. E-2900-4-D-92, Saskatchewan Research Council, Saskatoon, Canada, 157 p.

Connor DJ, Wang YP (1993) Climatic change and the Australian wheat crop. In Proceedings of the Third Symposium on the Impact of Climatic Change on Agricultural Production in the Pacific Rim, Taipei, Taiwan, ROC, pp. 29-47.

Cousens RD, Mokhtari S (1998) Seasonal and site variability in the tolerance of wheat cultivars to interference from *Lolium rigidum*. Weed Res 38, 301-307. doi: 10.1046/j.1365-3180.1998.00097.x

Curtis PS, Wang XA (1998) Meta-analysis of elevated CO2 effects on woody plant mass, form, and physiology. Oecologia 113, 299-313. doi: 10.1023/A:1020305006949

Darwin R, Kennedy D (2000) Economic effects of CO_2 fertilization of crops: transforming changes in yield into changes in supply. Environ Mod Assess 5, 157-168. doi:10.1023/A:1019013712133

Daudenmire R (1970) Steppe vegetation of Washington. Washingtom Agric. Exp. Stm. Bull. No. 62.

DeCeault MT, Polito VS (2008) High temperatures during bloom can inhibit pollen germination and tube growth, and adversely affect fruit set in the *Prunus Domestica* cultivars "improved French" and "Muir Beauty". In Proceedings of IX International Symposium on Plum and Prune Genetics, Breeding and Pomology, March 16-19, 2008, Dipartimento Colture Arboree, University of Palermo, Italy, 874 p.

DeFelice MS, Witt WW, Barrett M (1988) Velvetleaf (*Abutilon theophrasti*) growth and development in conventional and no-till corn (Zea mays). Weed Sci 36, 609-615.

Didon UME, Bostrom U (2003) Growth and development of six barley (*Hordeum vulgare* ssp. *vulgare* L.) cultivars in response to a model weed (*Sinapis alba* L.). J Agron Crop Sci 189, 409-417. doi: 10.1046/j.0931-2250.2003.00065.x

Dijkstra FA, Pendall E, Mosier AR, King JY, Milchunas DG, Morgan JA (2008) Long-term enhancement of N availability and plant growth under elevated CO_2 in a semi-arid grassland. Func Ecol. 22:975-982 .doi: 10.1111/j.1365-2435.2008.01398.x

Dingkuhn M, Singh BB, Clerget B, Chantereau J, Sultan B (2010) Past, Present and Future Criteria to Breed Crops for Water-Limited Environments in West Africa. In Proceedings of the 4th International Crop Science Congress "New directions for a diverse planet", 26 September - 1 October 2004, Brisbane, Australia.

Drake BG, Gonzalez-Meler MA, Long SP (1997) More efficient plants: A consequence of rising atmospheric CO_2? Annu. Rev. Plant Physiol. Plant Mol. Biol. 1997. 48, 609-639. doi: 10.1146/annurev.arplant.48.1.609

Duan YH, Zhang YL, Ye LT, Fan XR, Xu GH, Shen QR (2007) Responses of rice cultivars with different nitrogen use efficiency to partial nitrate nutrition. Ann Bot 99, 1153-1160. doi: 10.1093/aob/mcm051

Dukes JS, Mooney HA (1999) Does global change increase the success of biological invaders? Trends Ecol Evol 14:4, 135-139

Dukes JS, Pontius J, Orwig D, Garnas JR, Rodgers VL, Brazee N, Cooke B, Theoharides KA, Stange EE, Harrington R, Ehrenfeld J, Gurevitch J, Lerdau M, Stinson K, Wick R, Ayres M (2009) Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: what can we predict? Can J For Res 39:231-248. doi: 10.1139/X08-171

Edwards GE, Huber S (1981) The C4 Pathway. In M.D. Hatch and N.K. Boardman (eds) The Biochemistry of Plants. Academic Press, New York, pp. 238-281. ISBN: 012675408X

Ehleringer J (1983) Ecophysiology of *Amaranthus palmeri*, a Sonoran Desert summer annual. Oecologia 57, 107-112. doi: 10.1007/BF00379568

939 Erbs M, Manderscheid R, Jansen G, Seddig S, Pacholski A, Weigel HJ (2010) Effects of free-air
940 CO_2 enrichment and nitrogen supply on grain quality parameters and elemental composition of
941 wheat and barley grown in crop rotation. Agric Ecosyst Environ 136, 59-68. doi:
942 10.1016/j.agee.2009.11.009
943
944 Evans RO, Skagss RW, Sneed RE (1991) Stress day index models to predict corn and soybean
945 relative yield under high water table conditions. Trans ASAE 5, 1997-2005. doi: 0001-2351 /91
946 /3405-1997
947
948 Fargione J, Tilman D (2006) Plant species traits and capacity for resource reduction predict yield
949 and abundance under competition in nitrogen-limited grassland. Funct Ecol 20, 533-540. doi:
950 10.1111/j.1365-2435.2006.01116.x
951
952 Foulkes MJ, Scott PK, Sylvester-Bradley R (1997) Optimising winter wheat varietal selection on
953 drought-prone soil types. In Optimising cereal inputs: Its Scientific Basis. Asp Appl Biol 50, 61-
954 77.
955
956 Foulkes MJ, Scott RK, Sylvester-Bradley R, Clare RW, Evans EJ, Frost DL, Kettlewell PS,
957 Ramsbottom JE, White E (1994) Suitabilities of UK winter wheat (*Triticum aestivum* L.)
958 varieties to soil and husbandry conditions. Plant Varieties Seeds 7, 161-181.
959
960 Fuhrer J (2003) Agroecosystem responses to combinations of elevated CO_2, ozone, and global
961 climate change. Agric Ecosyst Environ 97, 1-20. doi: 10.1016/S0167-8809(03)00125-7
962
963 Gala Bijl C, Fisher M (2011) Crop adaptation to climate change. CSA News Magazine July
964 2011, 5-9
965
966 Gealy DR, Anders M, Watkins B, Duke S (2014) Crop performance and weed suppression by
967 weed-suppressive rice cultivars in furrow- and flood-irrigated systems under reduced herbicide
968 inputs. Weed Sci 62:303-320. doi: 10.1614/WS-D-13-00104.1.
969
970 Gealy DR, Wailes EJ, Estorninos LE Jr, Chavez RSC (2003) Rice cultivar differences in
971 suppression of barnyardgrass (*Echinochloa crus-galli*) and economics of reduced propanil
972 rates. Weed Sci 51:601–609. doi: 10.1614/0043-1745(2003)051[0601:RCDISO]2.0.CO;2
973
974 Gealy DR, Moldenhauer KAK (2012) Use of 13C isotope discrimination analysis to quantify
975 distribution of barnyardgrass and rice roots in a four-year study of weed suppressive rice. Weed
976 Sci 60:133-142. doi: 10.1614/WS-D-10-00145.1
977
978 Gealy DR, Moldenhauer KA, Duke S (2013a) Root distribution and potential interactions
979 between allelopathic rice, sprangletop (*Leptochloa* spp.), and barnyardgrass (*Echinochloa crus-*
980 *galli*) based on 13C isotope discrimination analysis. J Chem Ecol 39:186-203. doi:
981 10.1007/s10886-013-0246-7
982

Gealy DR, Moldenhauer KA, Jia MH (2013b) Field performance of STG06L-35-061, a new genetic resource developed from crosses between weed-suppressive *indica* rice and commercial southern U.S. long-grains. Plant Soil 370:277-293. doi: 10.1007/s11104-013-1587-2

Gealy DR Yan W (2012) Weed suppression potential of 'Rondo' and other *indica* rice germplasm lines. Weed Technol. 26: 517-524. doi: http://dx.doi.org/10.1614/WT-D-11-00141.1

Gent MPN, Kiyomoto RK (1998) Physiological and Agronomic consequences of Rht genes in wheat. In Basra AS (ed) Crop Science: Recent Advances, pp 27-46.

Gibson KD, Fischer AJ, Foin TC, Hill JE (2003) Crop traits related to weed suppression in water-seeded rice (Oryza sativa L.). Weed Sci 51:87-93. doi: http://dx.doi.org/10.1614/0043-1745(2003)051[0087:CTRTWS]2.0.CO;2

Ghannoum O (2009) C4 photosynthesis and water stress. Ann. Bot. 103:635-644. doi: 10.1093/aob/mcn093

Glover J, Johnson H, Lizzio J, Wesley V, Hattersley P, Knight C (2008) Australia's crops and pastures in a changing climate - can biotechnology help? Australian Government Bureau of Rural Sciences, Canberra.

Goldwasser Y, Lanini WT, Wrobel RL (2001) Tolerance of tomato varieties to Lespedeza dodder. Weed Sci 49, 520-523. doi: http://dx.doi.org/10.1614/0043-1745(2001)049[0520:TOTVTL]2.0.CO;2

Gouache D, Le Bris X, Bogard M, Deudon O, Page C, Gate P (2012) Evaluating agronomic adaptation options to increasing heat stress under climate change during wheat grain filling in France. Eur J Agron 39, 62-70. doi: http://dx.doi.org/10.1016/j.eja.2012.01.009

Gregory PJ, Ingram JS, Brklacich M (2005) Climate change and food security. Phil. Trans. R. Soc. B 360, 2139-2148. doi: 10.1098/rstb.2005.174

Grime JP (1979) Evidence for the strategies in plants and relevance to ecological and evolutionary theory. Am Natur 111, 1169-1192

Grover A, Mittal D, Negi M, Lavania D (2013) Generating high temperature tolerant transgenic plants: Achievements and challenges. Plant Sci 205-206, 38-47. doi: http://dx.doi.org/10.1016/j.plantsci.2013.01.005

Grzesiak MT, Marcinska I, Janowiak F, Rzepka A, Hura T (2012) The relationship between seedling growth and grain yield under drought conditions in and triticale genotypes. Acta Physiol Plantarum 34, 1757-1764. doi: 10.1007/s11738-012-0973-3

Halvorson WL, Guertin P (2003) Factsheet for *Ipomoea purpurea* (L.) Roth. USGS weeds in the west project: status of introduced plants in Southern Arizona parks. U.S. Geological Survey, Southwest Biological Science Center and University of Arizona.

Hamayun M, Sohn EY, Khan SA, Shinwari ZK, Khan AL, Lee IJ (2010) Silicon alleviates the adverse effects of salinity and drought stress on growth and endogenous plant growth hormones of soybean (*Glycine max* L.). Pak J Bot 42:1713-1722. doi: 10.1007/s00344-012-9274-8

Hardin G (1960) The Competitive Exclusion Principle. Science 131:3409, 1292-1297

Hay JL, Walker AJ (1992) An introduction to the physiology of crop yield. Longman Scientific and Technical and John Wiley and sons Publications, pp. 163-168. ISBN: 978-0582408081

Heap I (2015) The International Survey of Herbicide Resistant Weeds. Online Internet. www.weedscience.org [Accessed, November 12, 2015].

Hulme M (1996) Global warming. Prog Phys Geogr 20, 216-223

Hulme M, Wigley T, Jiang T, Zhao Z, Wang F, Ding Y, Leemans R, Markham A (1992) Climate change due to the greenhouse effect and its implications for China. CRU/WWF/SMA, World Wide Fund for Nature, Gland, Switzerland.

IPCC (2014). Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL (eds.) Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1132 p.

IPCC (2007) Climate change 2007: the physical science basis. Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds) Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, UK/New York, NY: Cambridge University Press.

IPCC (2001) Impacts, Adaptation and Vulnerability. Contribution of the Working Group II to the Third Assessment Report on the Intergovernmental Panel on Climate Change. UK: Cambridge University Press. ISBN: 978-0521015004

Jennings PR, Coffman WR, Kauffman HE (1979) Rice improvement. IRRI, Los Banos, Philippines, 186 p.

Johnson DE, Dingkuhn M, Jones MP, Mahamane MC (1998) The influence of rice plant type on the effect of weed competition on *O. sativa* and *O. glaberrima*. Weed Res 38:207-216. doi: 10.1046/j.1365-3180.1998.00092.x

Johnson D, California Invasive Plant Council (2013) Invasive plants, climate change and agriculture, presented at the California Department of Food and Agriculture Climate Change Adaptation Consortium, March 20, American Canyon, CA

Jones MP, Dingkuhn M, Aluko GK, Semon M (1997) Interspecific *O. sativa* L.× *O. glaberrima* Steud. progenies in upland rice improvement. Euphytica 92, 237-246. doi: 10.1023/A:1002969932224

Juskiw PE, Helm JH, Salmon DF (2000) Competitive ability in mixtures of small grain cereals. Crop Sci 40, 159-164. doi: 10.2135/cropsci2000.401159x

Karl TR, Melillo JM, Peterson TC (eds.) (2009) Global climate change impacts in the United States. A State of Knowledge Report from the U.S. Global Change Research Program. Cambridge University Press, New York, USA, 196 p. ISBN: 978-0-521-14407-0

Kasperbauer MJ, Karlen DL (1994) Plant spacing and reflected far-red light effects on phytochrome-regulated photosynthate allocation in corn seedlings. Crop Sci 34, 1564-1569

Kawano K, Jennings PR (1983) Tropical crop breeding-achievements and challenges. In IRRI, Potential productivity of field crops under different environments, Los Banos Laguna, Phillipines

Khodarahmpour Z (2011) Effect of drought stress induced by polyethylene glycol (PEG) on germination indices in corn (*Zea mays* L.) hybrids. Afr J Biotech 10, 18222-18227

King LJ (1966) Weeds of the World-Biology and Control. Interscience Pub., Inc., NY, 270 p. ISBN: 112596779X

Kong CH, Hu F, Wang P, Wu JL (2008) Effect of allelopathic rice varieties combined with cultural management options on paddy field weeds. Pest Manag Sci 64:276-282. doi: 10.1002/ps.1521.

Kong CH, Chen XH, Hu F, Zhang SZ (2011) Breeding of commercially acceptable allelopathic rice cultivars in China. Pest Manag Sci 67:1100-1106. doi: 10.1002/ps.2154

Korres NE (2000) The effects of seed rate and varietal selection for weed suppression and herbicide sensitivity in winter wheat (*Triticum aestivum* L.). PhD Thesis, Reading University, Reading University, UK.

Korres NE (2005) Encyclopaedic dictionary of weed science. Theory and digest, Intercept and Lavoisier (pubs). Andover, Paris, 695 p. ISBN: 1-898298-99-8.

Korres NE, Froud-Williams RJ (2002) Effects of winter wheat cultivars and seed rate on the biological characteristics of naturally occurring weed flora. Weed Res 42, 417-428. doi: 10.1046/j.1365-3180.2002.00302.x

Korres NE, Froud-Williams RJ (2004) The interrelationships of winter wheat cultivars, crop density and competition of naturally occurring weed flora. Biol Agric Hortic 22:1, 1-20. doi: 10.1080/01448765.2004.9754984

Korres NE, Froud-Williams RJ, Chachalis D, Pavli O, Skaracis GN (2008) Yielding ability and competitiveness of wheat cultivars against weeds, 4th EPSO Conference: Plants for Life, Toulon (Cote d' Azur), France, 22-26 June 2008, pp. 52.

Korres NE, Norsworthy JK, Bagavathiannan MV, Mauromoustakos A (2015) Distribution of arable weed populations along eastern Arkansas Mississippi Delta roadsides: Factors affecting weed occurrence. Weed Technol. doi: http://dx.doi.org/10.1614/WT-D-14-00152.1

Korres NE, Norsworthy JK (2015) Influence of Palmer amaranth interrow distance and emergence date on seed production in wide-row and drill-seeded soybean. In Proceedings of the Weed Science Society of America, Annual Meeting, Lexington, Kentucky, USA, February 9-12, 2015

Kramer PJ (1983) Water deficits and plant growth. In: P.J. Kramer (ed.). Water relations of plants. Academic Press, New York, pp 342-389. ISBN: 978-0124250406

Larson C (2013) Losing arable land, China faces stark choice: adapt or go hungry. Science 339, 644-645. doi:10.1126/science.339. 6120.644

Larsson S, Gorny AG (1988) Grain yield and drought resistance indices of oat cultivars in field rain shelter and laboratory experiments. J Agron Crop Sci 161, 277-286. doi: 10.1111/j.1439-037X.1988.tb00668.x

Le Houerou HN (1996) Climate changes, drought and desertification. J Arid Environ 34, 133-185. doi: http://dx.doi.org/10.1006/jare.1996.0099

Leakey ADB, Uribelarrea M, Ainsworth EA, Naidu SL, Rogers A, Ort DR, Long SP (2006) Photosynthesis, productivity, and yield of are not affected by open-air elevation of CO_2 concentration in the absence of drought. Plant Physiol 140:2, 779-790. doi: 10.1104/pp.104.900185

Leakey ADB, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR (2009) Elevated CO_2 effects on plant carbon, nitrogen and water relations: six important lessons from FACE. J. Exp. Bot. 60:2859-2876. doi: 10.1093/jxb/erp096

Leguizamon ES, Yanniccari ME, Guiamet JJ, Acciaresi HA (2011) Growth, gas exchange and competitive ability of *Sorghum halepense* populations under different soil water availability. Can J Plant Sci 91, 1011-1025. doi: 10.4141/cjps10202

Lemerle D, Verbeek B, Orchard B (2001) Ranking the ability of wheat varieties to compete with Lolium rigidum. Weed Res 41, 197-209. doi: 10.1046/j.1365-3180.2001.00232.x

Lobell DB, Ortiz-Monasterio JI, Asner GP, Matson PA, Naylor RL (2005) Analysis of wheat yield and climatic trends in Mexico. Field Crops Res 94, 250-256. doi: http://dx.doi.org/10.1016/j.fcr.2005.01.007

Lobell DB, Field CB (2011) California perennial crops in a changing climate. Clim Change 109:1, 317-333. doi: 10.1007/s10584-011-0303-6

Lobell DB, Field CB, Cahill KN, Bonfils C (2006) Impacts of future climate change on California perennial crop yields. Model projections with climate and crop uncertainties. Agric For Meteorol 141:2-4, 208-218. doi: http://dx.doi.org/10.1016/j.agrformet.2006.10.006

Lupton FGH, Oliver RH, Ellis FB, Barnes BT, Howse KR, Welbank PJ, Taylor PJ (1974) Root and shoot growth of semi-dwarf and taller winter wheats. Ann Appl Biol 77, 129-144. doi: 10.1111/j.1744-7348.1974.tb06881.x

Ma HJ, Shin DH, Lee IJ, Koh JC, Park SK, Kim KU (2006) Allelopathic K21 selected as promising allelopathic rice. Weed Biol Manag 6:189-196. doi: 10.1111/j.1445-6664.2006.00219.x

Malik VS, Swanton CJ, Michaels TE (1993). Interaction of white bean (*Phaseolus vulgaris* L.) cultivars, row spacing, and seeding density with annual weeds. Weed Sci 41, 62-68

Manea A, Leishman MR, Downey PO (2011) Exotic C4 grasses have increased tolerance to glyphosate under elevated carbon dioxide. Weed Sci 59, 28-36. doi: http://dx.doi.org/10.1614/WS-D-10-00080.1

Mann GC (1980) Variety development. Proceedings of the 16 th NIAB Crop Conference, 7-15.

Mason HE, Spaner D (2006) Competitive ability of wheat in conventional and organic management systems: A review of the literature. Can J Plant Sci 86, 333-343. doi: http://pubs.aic.ca/doi/abs/10.4141/P05-051

Matthews RB, Kropff MJ, Bachelet D, van Laar HH (1994) The impact of global climate change on rice production in Asia: a simulation study. Report No. ERL-COR-821, U.S. Environmental Protection Agency, Environmental Research Laboratory, Corvallis, OR.

Maun MA, Bennett SCH (1986) The biology of Canadian weeds. 77. *Echinochloa crus-galli* (L.) Beauv. Can J Plant Sci 66, 739-759. doi: http://pubs.aic.ca/doi/abs/10.4141/cjps86-093

McGiffen ME Jr, Forcella F, Lindstrom MJ, Reicosky DC (1997) Covariance of cropping systems and foxtail density as predictors of weed interference. Weed Sci 45, 388-396. ISSN: 00431745

Mohler CL (2001) Enhancing the competitive ability of crops. In Liebman M, Mohler CL, Staver CP, (eds). Ecological Management of Agricultural Weeds, Cambride Univeristy Press: Cambridge, UK, pp. 231-269. ISBN: 1139427245, 9781139427241

Monteith JL (1977) Climate and efficiency of crop production in Britain. Philos Trans R Soc Lond, B 281, 277-294. doi: 10.1098/rstb.1977.0140

Morison JIL (1989) Plant growth in increased atmospheric CO_2 in Fantechi R, Ghazi A (eds). Carbon Dioxide and Other Greenhouse Gases: Climatic and Associated Impacts. Kluwer Academic Publishers, Dordrecht, Boston, Landon, pp. 228-244. ISBN: 0-7923-0191-9

Naidu VSGR, Paroha S (2008) Growth and biomass partitioning in two weed species *Parthenium hysterophorus* (C_3) and *Amaranthus viridis* (C_4) under elevated CO_2. Eco Env Cons 14:4, 9-12

Newton PCD, Clark H, Bell CC, Glasgow EM (1996) Interaction of soil moisture and elevated CO_2 on the above-ground growth rate, root length density and gas exchange of turves from temperate pasture. J Exp Bot 47:6, 771-779. doi: 10.1093/jxb/47.6.771

Nordby DE, Alderks DL, Nafziger ED (2002) Competitiveness with weeds of soybean cultivars with different maturity and canopy width characteristics. Weed Tech 21:1082-1088

Obirih-Opareh N, Adwoa Onumah J (2014) Climate Change Impact Pathways on Agricultural Productivity in Africa: A Review. J Environ Earth Sci 4:4, 115-121. ISSN: 2224-3216

Oerke EC (2006) Crop losses to pests. J Agric Sci 144: 31-43. doi: http://dx.doi.org/10.1017/S0021859605005708

Olesen J. E. and Bindi M. (2002). Consequences of climate change for European agricultural productivity, land use and policy. Eur J Agron 16, 239-262. doi: 10.1016/S1161-0301(02)00004-7

Olesen JE, Hansen PK, Berntsen J, Christensen S (2004) Simulation of above-ground suppression of competing species and competition tolerance in winter wheat varieties. Field Crops Res 89, 263-280. doi: 10.1016/j.fcr.2004.02.005

Osunsami S (2009) Killer Pigweeds Threaten Crops in the South. http://abcnews.go.com/WN/pig-weed-threatensagriculture - industry-overtaking-fields-crops/story?id8766404 [Accessed May 5, 2015]

Pace PF, Crale HT, El-Halawany SHM, Cothren JT, Senseman SA (1999) Drought induced changes in shoot and root growth of young cotton plants. J Cotton Sci 3:183-187. doi: http://www.journal.cotton.org/journal/1999-03/4/upload/jcs03-183.pdf

Paknejad F, Nasri M, Moghadam HRT, Zahedi H, Alahmadi MF (2007) Effects of drought stress on chlorophyll fluorescence parameters, chlorophyll content and grain yield of wheat cultivars. J Biol Sci 7, 841-847. doi: http://en.journals.sid.ir/ViewPaper.aspx?ID=103989

Paolini R, Del Puglia S, Principi M, Barcellona O, Riccardi E (1998) Competition between safflower and weeds as influenced by crop genotype and sowing time. Weed Res 38, 247-255. doi: 10.1046/j.1365-3180.1998.00096.x

Parry ML (1990) Climate change and world agriculture. London: Earthscan Publications. doi: 10.1080/00139157.1991.9931405

Patterson DT, Westbrook JK, Joyce RJV, Lingren PD, Rogasik J (1999) Weeds, insects and diseases. Clim Change 43, 711-727. doi: 10.1023/A:1005549400875

Patterson DT (1995) Weeds in a changing climate. Weed Sci 43, 685-701. ISSN: 00431745

Patterson DT, Flint EP, Beyers JL (1984) Effects of CO_2 enrichment on competition between a C4 weed and a C3 crop. Weed Sci 32:1, 101-105. ISSN: 00431745

Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X, Centeno GS, Khush GS, Cassman KG (2004) Rice yields decline with higher night temperature from global warming. PNAS 101:27, 9971-9975. doi: 10.1073/pnas.0403720101

Peters K, Breitsameter L, Gerowitt B (2014) Impact of climate change on weeds in agriculture: a review. Agron Sustain Dev 34:707-721. doi: 10.1007/s13593-014-0245-2

Pheng S, Olofsdotter M, Jahn G, Nesbitt H, Adkins S (2009a) Potential allelopathic rice lines for weed management in Cambodian rice production. Weed Biol Manag 9:259–266. doi: 10.1111/j.1445-6664.2009.00349.x

Pheng S, Olofsdotter M, Jahn G, Adkins S (2009b) Allelopathic potential of Cambodian rice lines under field conditions. Weed Biol Manag 9:267-275. doi: 10.1111/j.1445-6664.2009.00350.x

Poorter H, Navas ML (2003) Plant growth and competition at elevated CO_2: On winners, losers and functional groups. New Phytol 157:2, 175-198. doi: 10.1046/j.1469-8137.2003.00680.x

Pope KS, DoseV, Da Silva D, Brown PH, Leslie CA, DeJong TM (2013) Detecting nonlinear response of spring phenology to climate change by Bayesian analysis. Glob Chang Biol 19:5, 1518-1525. doi: 10.1111/gcb.12130

Pope KS (2012) Climate Change Adaptation: Temperate Perennial Crops, presented at the California Department of Food and Agriculture Climate Change Adaptation Consortium, November 28, Modesto, CA

Porter JR, Semenov MA (2005) Crop responses to climatic variation. Philos Trans R Soc B: Biol Sci 360:2021-2035. doi: 10.1098/rstb.2005.1752

Prasad PVV, Boote KJ, Allen LH Jr (2006a) Adverse high temperature effects on pollen viability, seed-set, seed yield and harvest index of grain-sorghum (*Sorghum bicolor* L. moench) are more severe at elevated carbon dioxide due to higher tissue temperatures. Agric For Meteorol, 139:3-4, 237-251. doi: 10.1016/j.agrformet.2006.07.003

Prasad PV, Boote KJ, Allen LH, Sheehy JE, Thomas JM (2006b) Species, ecotype and cultivar differences in spikelet fertility and harvest index of rice in response to high temperature stress. Field Crops Res 95:398-411. doi: 10.1016/j.fcr.2005.04.008

Qaderi MM, Lynch AL, Godin VJ (2013) Single and interactive effects of temperature, carbon dioxide, and watering regime on the invasive weed black knapweed (*Centaurea nigra*) Ecoscience 20:4, 328-338. doi: http://dx.doi.org/10.2980/20-4-3631

Ragab AR, Abdel-Raheem AT, Kasem ZA, Omar FD, Samera AM (2007) Evaluation of R1 tomato somaclone plants selected under poly ethylene glycol (PEG) treatments. Afr Crop Sci Soc 8, 2017-2025. doi: http://www.acss.ws/upload/xml/research/468.pdf

Rajcan I, Swanton CJ (2001) Understanding -weed competition: resource competition, light quality and the whole plant. Field Crops Res 71, 139-150. doi: 10.1016/S0378-4290(01)00159-9

Riar DS, Norworthy JK, Steckel LE, Stephenson DO, Eubank TW, Scott RC (2013) Assessment of weed management practices and problem weeds in the midsouth United States-soybean: a consultant's perspective. Weed Technol 27:612-622. doi: http://dx.doi.org/10.1614/WT-D-12-00167.1

Richards RA, Thurling N (1978) Variation between and within species of rapeseed (*Brasica campestris* and *B napus*) in response to drought stress. I. Sensitivity at different stages of development. Aust J Agric Res 29, 469-477. doi: 10.1071/AR9780469

Ripley BS, Gilbert ME, Ibrahim DG, Osborne CP (2007) Drought constraints on C4 photosynthesis: Stomatal and metabolic limitations in C3 and C4 subspecies of *Alloteropsis semialata*. J. Exp. Bot. 58:1351-1363. doi: 10.1093/jxb/erl302

Rosenzweig C, Tubiello FN (2007) Adaptation and mitigation strategies in agriculture: an analysis of potential synergies. Mitig Adapt Strat Glob Change 12, 855-873. doi: 10.1007/s11027-007-9103-8

Royo C, Abaza M, Bianco R, Moral LFG (2000) Triticale grain growth and morphometry as affected by drought stress, late sowing and simulated drought stress. Aust J Physiol *27*, 1051-1059. doi: 10.1071/PP99113

Sakthivelu G, Devi MKA, Giridhar P, Rajasekaran T, Ravishankar GA, Nedev T, Kosturkova G (2008) Drought induced alterations in growth, osmotic potential and in vitro regeneration of soybean cultivars. Genet Appl Plant Physiol 34, 103-112. doi: http://www.bio21.bas.bg/ipp/gapbfiles/v-34_pisa-08/08_pisa_1-2_103-112.pdf

Schittenhelm S, Schroetterm S (2014) Comparison of drought tolerance of , sweet sorghum and sorghum-sudangrass hybrids. J Agron Crop Sci 200, 46-53. doi: 10.1111/jac.12039

Schmidhuber J, Tubiello FN (2007) Global food security under climate change. Proc. Natl. Acad. Sci. U.S.A (PNAS) 104:50, 19703-19708. doi: 10.1073/pnas.0701976104

Scott JK, Murphy H, Kriticos DJ, Webber BL, Ota N, Loechel B (2014) Weeds and climate change: Supporting weed management adaptation. CSIRO, Australia.

Seneweera S, Milham P, Conroy J (1994) Influence of elevated CO_2 and phosphorus nutrition on the growth and yield of a short-duration rice *(Oryza sativa* L. cv. Jarrah). Aust J Plant Physiol 21, 281-292. doi: 10.1071/PP9940281

Sheley RL, Svejcar TJ, Maxwell BD (1996) A theoretical framework for developing successional weed management strategies on rangeland. Weed Technol 10, 766-773. doi: ISSN: 0890037X

Song L, Li FM, Fan XW, Xiong YC, Wang WQ, Wu XB, Turner NC (2009) Soil water availability and plant competition affect the yield of spring wheat. Eur J Agron 31, 51-60. doi: 10.1016/j.eja.2009.03.003

Song L, Zhang DW, Li FM, Fan XW, Ma Q, Turner NC (2010) Soil water availability alters the inter- and intra-cultivar competition of three spring wheat cultivars bred in different eras. J Agron Crop Sci, 196, pp. 323-335. doi: 10.1016/j.eja.2009.03.003

Southworth J, Pfeifer RA, Habeck M, Randolph JC, Doering OC, Johnston JJ, Rao DG (2002) Changes in soybean yields in the midwestern United States as a result of future changes in climate, climate variability, and CO_2 fertilization. Clim Chang 53:4, 447-475. doi: 10.1023/A:1015266425630

Starggenborg SA, Dhuyvetter KC, Gordon BW (2008) Grain sorghum and corn comparisons: yield, economic, and environmental responses. Agron J 100, 1600-1604. doi: 10.2134/agronj2008.0129

Stevanato P, Trebbi D, Bertaggia M, Colombo M, Broccanello C, Concheri G, Saccomani M (2011). Root traits and competitiveness against weeds in sugar beet. Int Sugar J 113, 497-501. doi: http://www.researchgate.net/profile/Piergiorgio_Stevanato/publication/236848489_Root_traits_and_competitiveness_against_weeds_in_sugar_beet/links/0deec519e734789524000000.pdf [Accessed October 12, 2015]

Stone MJ, Cralle HT, Chandler JM, Bovey RW, Carson KH (1998) Above- and below-ground interference of wheat (*Triticum aestivum*) by Italian ryegrass (*Lolium multiflorum*). Weed Sci 46, 438-441. ISSN: 00431745

Stone PJ, Nicolas ME (1995) Effect of timing of heat stress during grain filling on two wheat varieties differing in heat tolerance. I. Grain Growth. Aust J Plant Physiol 22, 927-93. doi: 10.1071/PP9950927

Storrie A, Cook T (2007) What impact does drought have on weeds? Primefact 430, State of New South Wales through NSW Department of Primary Industries. Job number 7322, pp. 1-3.

Stratonovitch P, Storkey J, Semenov AA (2012) A process-based approach to modelling impacts of climate change on the damage niche of an agricultural weed. Glob Chang Biol 18, 2071-2080. doi: 10.1111/j.1365-2486.2012.02650.x

Streck NA (2005) Climate change and agroecosystems: the effect of elevated atmospheric CO_2 and temperature on crop growth, development, and yield. Ciencia Rural, Santa Maria 35:3, 730-740. doi: http://dx.doi.org/10.1590/S0103-84782005000300041

Taiz L, Zeiger E (1991) Plant Physiology. Benjamin/Cummings series, Benjamin-Cummings Pub Co., Lewiston, NY, pp. 559

Thomas PEL, Allison JCS (1975) Competition between and *Rottboellia exaltata*. J Agric Sci 84, 305-312

Tian X, Matsui T, Li S, Yoshimoto M, Kobayasi K, Hasegawa T (2010) Heat-induced floret sterility of hybrid rice (*Oryza sativa*) cultivars under humid and low wind conditions in the field of Jianghan Basin, China. Plant Prod Sci 13:3, 243-251. doi: http://doi.org/10.1626/pps.13.243

Travlos IS (2012) Reduced herbicide rates for an effective weed control in competitive wheat cultivars. Int J Plant Prod 6, 1-13. ISSN: 1735-6814

Trenberth KE, Jones PD, Ambenje P, Bojariu R, Easterling D, Klein Tank A, Parker D, Rahimzadeh F, Renwick JA, Rusticucci M, Soden B, Zhai P (2007) Observations: Surface and Atmospheric Climate Change. In Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds) (2007) Climate Change: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Tubiello FN, Ewert F (2002) Simulating the effects of elevated CO_2 on crop growth and yield: approaches and applications for climate change. Eur J Agron 18:1-2, 57-74. doi: 10.1016/S1161-0301(02)00097-7

Tuong TP, Bouman BAM (2003) Rice Production in Water-Scarce Environments, In Kijne JW, Barker R, Molden D. (eds.) Water Productivity in Agriculture: Limits and Opportunities for Improvements, CABI Publishing pp. 53-67. ISBN: 0851996698, 9780851996691

Ugarte C, Calderini DF, Slafer GA (2007) Grain weight and grain number responsiveness to preanthesis temperature in wheat, barley and triticale. Field Crops Res 100:240-248. doi: 10.1016/j.fcr.2006.07.010

Valerio MM, Tomecek B, Lovelli S, Ziska LH (2011) Quantifying the effect of drought on carbon dioxide-induced changes in competition between a C3 crop (tomato) and a C4 weed (*Amaranthus retroflexus*). Weed Res 51, 591-600. doi: 10.1111/j.1365-3180.2011.00874.x

Vermeulen SJ, Campbell BM, Ingram JSI (2012) Climate change and food systems. Annu Rev Environ Resour 37:195-222. doi: 10.1146/annurev-environ-020411-130608

Vollmann J, Wagentristl H, Hartl W (2010) The effects of simulated weed pressure on early maturity soybeans. Eur J Agron 32, 243-248. doi: 10.1016/j.eja.2010.01.001

Walthall CL, Hatfield J, Backlund P, Lengnick L, Marshall E, Walsh M, Adkins S, Aillery M, Ainsworth EA, Ammann C, Anderson CJ, Bartomeus I, Baumgard LH, Booker F, Bradley B, Blumenthal DM, Bunce J, Burkey K, Dabney SM, Delgado JA, Dukes J, Funk A, Garrett K, Glenn M, Grantz DA, Goodrich D, Hu S, Izaurralde RC, Jones RAC, Kim SH, Leaky ADB, Lewers K, Mader TL, McClung A, Morgan J, Muth DJ, Nearing M, Oosterhuis DM, Ort D, Parmesan C, Pettigrew WT, Polley W, Rader R, Rice C, Rivington M, Rosskopf E, Salas WA, Sollenberger LE, Srygley R, Stockle C, Takle ES, Timlin D, White JW, Winfree R, Wright-Morton L, Ziska LH (2012) Climate Change and Agriculture in the United States: Effects and Adaptation. USDA Technical Bulletin 1935. Washington, DC, pp. 186.

Waltz E (2014) Beating the heat. Nat Biotechnol 32, 610-613. doi: 10.1038/nbt.2948

Wang YP, Handoko Jr, Rimmington GM (1992) Sensitivity of wheat growth to increased air temperature for different scenarios of ambient CO_2 concentration and rainfall in Victoria, Australia-a simulation study. Clim Res 2, 131-149. doi: 10.3354/cr002131

Ward SM, Webster TM, Steckel LE (2013) Palmer Amaranth (*Amaranthus palmeri*): A Review. Weed Technol 27:12-27. doi: http://dx.doi.org/10.1614/WT-D-12-00113.1

Wardlaw IF, Moncur L (1995) The response of wheat to high temperature following anthesis. I. The rate and duration of kernel filling. J Plant Physiol 22, 391-397. doi: 10.1071/PP9950391

Warwick SI, Black LD (1983) The biology of Canadian weeds. 61. *Sorghum halepense* (L.) Pers. Can J Plant Sci 63, 997-1014

Wayne P, Foster S, Connolly J, Bazzaz F, Epstein P (2002) Production of allergenic pollen by ragweed (*Ambrosia artemisiifolia* L.) is increased in CO_2-enriched atmospheres. Ann Allerg Asthma Immunol 8, 279-282. doi: 10.1016/S1081-1206(10)62009-1

Wheeler TR, Craufurd PQ, Ellis RH, Porter JR, Prasad PVV (2000) Temperature variability and the yield of annual crops. Agric Ecosyst Environ 82, 159-167. doi: 10.1016/S0167-8809(00)00224-3

Wiese AF, Vandiver CW (1970) Soil moisture effects on competitive ability of weeds. Weed Sci 18, 518-519. ISSN: 0043-1745

Woodward FI (1988) Temperature and the distribution of plant species, in Long SP, Woodward FI (eds). Plants and temperature. Published for the Society for Experimental Biology by the Company of Biologists. University of Cambridge, Cambridge, UK

Woodward FI, Williams BG (1987) Climate and plant distribution at global and local scales. Plant Ecol 69:1, 189-197. doi: 10.1007/978-94-009-4061-1_19

Worthington M, Reberg-Horton SC (2013) Breeding cereal crops for enhanced weed suppression: Optimizing allelopathy and competitive ability. J Chem Ecol 39:2. doi: 10.1007/s10886-013-0247-6

Zhao DL, Atlin GN, Bastiaans L, Spiertz JHJ (2006) Developing selection protocols for weed competitiveness in aerobic rice. Field Crop Res 97:272–285. doi:10.1016/j.fcr.2005.10.008

Zheng S, Nakamoto H, Yoshikawa K, Furuya T, Fukuyama M (2002) Influences of high night temperature on flowering and pod setting in soybean. Plant Prod Sci 5:3, 215-218. doi: http://doi.org/10.1626/pps.5.215

Zhu C, Zeng Q, Ziska LH, Zhu J, Xie Z, Liu G (2008) Effect of nitrogen supply on carbon dioxide-induced changes in competition between rice and barnyardgrass (*Echinochloa crus-galli*). Weed Sci 56:1, 66-71. doi: http://dx.doi.org/10.1614/WS-07-088.1

Zia-Ul-Haq M, Riaz M, De Feo V (2012) *Ipomea hederacea* Jacq.: A Medicinal Herb with Promising Health Benefits. Molecules 17, 13132-13145. doi: 10.3390/molecules171113132

Zimdahl RL (2007) Fundamentals of weed science. Academic Press, Burlington, San Diego, pp. 655. ISBN: 0123978181

Ziska LH (2003) Evaluation of the growth response of six invasive species to past, present and future carbon dioxide concentrations. J Exp Bot 54, 395-404. doi: 10.1093/jxb/erg027

Ziska LH (2004) Rising carbon dioxide and weed ecology. In Inderjit (ed). Weed biology and management, Kluwer Academic Press, Dordrecht, 159-176. ISBN: 978-90-481-6493-6

Ziska LH (2014) Climate, CO_2 and invasive weed management, in Ziska LH, Dukes JS (eds) Invasive species and global climate change. CABI Invasives Series, Boston, Wallingford, pp. 293-305. ISBN: 9781780641645

Ziska LH, Bunce JA (2007) Predicting the impact of changing CO_2 on crop yields: Some thoughts on food. New Phytol 175:4, 607-618. doi: 10.1111/j.1469-8137.2007.02180.x

Ziska LH, George K (2004) Rising carbon dioxide and invasive, noxious plants: potential threats and consequences. Water Resour Rev 16, 427-446

Ziska LH, Teasdale JR, Bunce JA (1999) Future atmospheric carbon dioxide may increase tolerance to glyphosate. Weed Sci 47, 608-615. ISSN: 00431745

Ziska LH (2014) Increasing minimum daily temperatures are associated with enhanced pesticide use in cultivated soybean along a latitudinal gradient in the Mid-Western United States. PLoS ONE 9(6): e98516. doi:10.1371/journal.pone.0098516.

Ziska LH, Runion GB (2007) Future Weed, Pest and Disease Problems for Plants. In: Newton PCD, Carran A, Edwards GR, Niklaus PA, (eds). Agroecosystems in a Changing Climate. CRC Press, Boston, MA, 262-279. ISBN: 9780849320880

Ziska LH, Faulkner S, Lydon J (2004) Changes in Biomass and Root:shoot Ratio of Field-grown Canada Thistle (*Cirsium Arvense*), a Noxious, Invasive Weed, with Elevated CO_2: Implications for Control with Glyphosate. Weed Sci 52:4, 584-588. doi: http://dx.doi.org/10.1614/WS-03-161R

Ziska LH, Runion GB (2006) Future weed, pest and disease problems for plants. In: Newton P, Carman A, Edwards G, Niklaus P (eds.) Agroecosystems in a Changing Climate. CRC. New York. Chapter 11, pp. 262-287. ISBN:

Table 1. Response of C3 and C4 weeds and crops to doubled atmospheric CO_2 levels in relation to biomass and leaf area production for both crop plants and weed species with C3 and C4 photosynthetic pathway

C3 species	Biomass	Leaf area	C4 species	Biomass	Leaf area
		Range of response (× growth at ambient CO_2 concentrations)			
Abutilon theophrastii	1-1.52	0.87-1.17	*Amaranthus retroflexus*	0.9-1.41	0.94-1.25
Bromus mollis	1.37	1.04	Andropogon virginicus	0.8-1.17	0.88-1.29
Bromus tectorum	1.54	1.46	*Cyperus rotundus*	1.02	0.92
Cassia obtusifolia	1.4-1.6	1.1-1.34	*Digitaria ciliaris*	1.06-1.6	1.04-1.66
Chenopodium album	1-1.6	1.22	*Echinochloa crus-galli*	0.95-1.6	0.98-1.77
Datura stramonium	1.7-2.72	1.46	*Eleusine indica*	1.02-1.2	0.95-1.77
Elytrigia repens	1.64	1.3	*Paspalum plicatum*	1.08	1.02
Phalaris aquatic	1.43	1.31	*Rottboellia cochinchinensis*	1.21	1.13
Plantago lanceolata	1-1.33	1.33	*Setaria faberii*	0.93-1.35	1-1.4
Rumex crispus	1.18	0.96	*Sorghum halepense*	0.56-1.1	0.99-1.3
		Range of response (% increase)			
Triticum aestivum	17-31		*Zea mays*	3.7-9	
Hordeum vulgare	30		*Sorghum bicolor*	9	
Glysine max	39				
Gossypium hirsutum	84				
Ipomoea batatas	59-111				

Adopted from Chandrasena (2009); Patterson (1985); Streck (2005).

Table 2. Effects of doubling CO_2 concentration on marketable yield* of major cereal, row, cash, vegetable crops and flowers.

Crop	Marketable yield (% increase)
Maize**	3.7-29
Sorghum**	6
Wheat**	8-35
Barley**	70
Rice**	25
Soybean***	22-45
Tobacco***	42
Potato***	51
Tomato***	20-26
Lettuce***	35-44
Cucumber***	30
Sunflower***	144
Chrysanthemum****	6
Cyclamen****	35
Rose****	8-27

Adopted from Streck (2005).

*Values shown in this Table were obtained by the compilation and analysis of the results of more than 770 reports about the effects of CO_2 enrichment on the economic yield of 24 agricultural crops and 14 other species; **cereals crops, *** row, cash and vegetables; ****flowers

Table 3. Plant height, stem and leaf dry weight, leaf area, and node number in drought-stressed and well-watered control cotton plants at the end of the drought (49 days after planting).

	Treatment	
Plant part	Drought*	Control
Plant height	20.0	27.9
Stem dry weight (g)	1.13	1.39
Leaf dry weight (g)	1.41	2.16
Leaf area (cm^2)	56	153
Node number	7.8	9.4

The drought treatment was imposed by withholding water for 13 d. *Means in a row are significantly different at the 0.05 probability level (based on Pace et al., 1999).

Table 4. Seasonal water use efficiency (g DM/kg water) under various water regimes and ambient and double CO_2 concentrations in various crop species

	Ambient CO_2	Double CO_2	Ratio
Sorghum	3.08	4.13	1.34
Wheat (well watered)	5.1	6.3	1.23
Wheat (water shortage)	6.2	8.9	1.43
Wheat	2.62	3.45	1.31
Wheat (well watered)	1.58	2.14	1.35*
Wheat (water shortage)	1.27	1.86	1.46*
Faba beans	4.91	7.82	1.59
Water hyacinth	1.4	2.6	1.85

Adopted from Morison 1993; *grain only

Table 5. Response of crop and weed species grown under competition as a function of high CO_2 concentration

C4 weed vs. C3 crops	High CO_2 favours	Environment
Sorghum halepense vs. *Festuca pratensis*	Crop	Greenhouse
Sorghum halepense vs. *Glysine max*	Crop	Growth chamber
Amaranthus retroflexus vs. Glysine max	Crop	Field
Echinochloa glabrescens vs *Oryza sativa*	Crop	Greenhouse
Paspalum dilatatum vs. various grasses	Crop	Growth chamber
Various grasses vs. *Medicago sativa*	Crop	Field
C3 weed vs. C3 crops		
Chenopodium album vs. *Beta vulgaris*	Crop	Growth chamber
Taraxacum officinale vs. *Medicago sativa*	Weed	Field
Plantago lanceolata vs. pasture	Weed	Growth chamber
Taraxacum and *Plantago* vs. pasture	Weed	Field
Cirsium arvensis vs. *Glysine max*	Weed	Field
Chenopodium album vs. *Glysine max*	Weed	Field
C4 weed vs. C4 crop		
Amaranthus retroflexus vs. *Shorghum bicolor*	Weed	Field

C3 weeds vs. C4 crops		
Xanthium strumarium vs. *Sorghum bicolor*	Weed	Greenhouse
Abutilon theophrasti vs. *Sorghum bicolor*	Weed	Field

Based on Bunce and Ziska (2000), Walthall et al. (2012).

Table 6. Potential effects of drought on Australian agricultural weeds

Weed	Impact
Blackberry (*Rubus fruticosus* L.)	Expected to retreat to higher altitudes due to its sensitivity to higher temperatures and drought
Chilean needle grass [*Nassella neesiana* (Trin. & Rupr.) Barkworth)]	Expected to increase its range because its increased invasiveness ability (long-lived, seed dispersed by wind and water) and drought tolerance
Gorse (*Ulex europaeus* L.)	Establishment into high-rainfall zones due to its sensitivity to drought
Lantana (*Lantana camara* L.)	Establishment into high-rainfall zones

Adopted from Anonymous (2008)

Table 7. Effects of increased CO_2 concentration on glyphosate efficacy for various weed species with different photosynthetic pathways

Common name	Latin name	P/S pathway*	Efficacy change
Canada thistle	*Cirsium arvense* (L.) Scop	C3	Reduced
Dallisgrass	*Paspalum dilatatum* Poir.	C4	Reduced
Lambsquarters	*Chenopodium album* L.	C3	Reduced
Lovegrass	*Eragrostis curvula* (Schrad.) Nees	C4	Reduced
Quackgrass	*Elytrigia repens* (L.) Gould	C3	Reduced
Redroot pigweed	*Amaranthus retroflexus* L.	C4	None
Rhodes grass	*Chloris gayana* Kunth	C4	Reduced
Smut grass	*Sporobolus indicus* (L.) R. Br.	C4	None

Adopted from Ziska, 2014; *Photosynthetic pathway

1752 **Table 8**. Response of crop plants and weeds under elevated CO_2, increased temperature and

1753 prolonged drought periods

			Climate change component		
	Plant response	Result	CO_2*	Temperature	Drought
Crop plants	Root mass	Root:shoot ratio	+		
	Leaf area	Interception of PAR**	+		
	Leaf development	Leaf area		-	
	Flowering	Vegetative stage		-	
	Harvesting	Yield		-	-
	Fruit production	Yield		-	
	Vernalization	Vegetative stage		-	
	Stomata conductance	Rate of photosynthesis		-	-
	Stomata closure	WUE		+	+
	CO_2:O_2	Rate of photosynthesis			-
	Respiration rate	Biomass production		+	
	Seed formation period	Yield		-	
	Biomass production	Yield	+	-	-
	Node number	Biomass, height			-
Weeds	Stomata closure	WUE	+		
	Maturity rate	Vegetative stage	+		
	Root biomass	Root:shoot ratio	+		
	Distribution			+	
	Vernalization	Vegetative stage		-	

	Biomass		+	
	Seed germination***	Distribution	+	
	Rhizomes***	Distribution	+	
	Seed longevity			+

*Elevated CO_2 favors, in most cases, C3 plant types; **photosynthetically active radiation; ***seed germination and rhizomes production, for most weed species, are affected negatively by low temperatures as it is mentioned in the text. Therefore, it is assumed that under relatively elevated temperatures will be affected positively. + and – signs indicate a positive or negative effect respectively.

Figure captions

Figure 1. A water stressed cotton field (a) (with permission from D. M. Oosterhuis) and heavily infested cotton field by Palmer amaranth (b) (with permission from J. K. Norsworthy)

Figure 2. Vegetative and reproductive response of maize and soybean to temperature increases (based on Karl et al., 2009).

Figure 3. Response of CO_2 assimilation in C3 vs. C4 plants to increases in CO_2 concentration (based on Taiz and Zeiger, 1991).

Figure 4. Rice weed suppression plots at Stuttgart, Arkansas, USA in which the superior competitiveness of cultivars STG06L-35-061 and PI312777 compared with Katy and Lemont is shown. A “light” infestation of barnyardgrass can be observed in the former compared to later plots. No herbicide was used to control grass weeds (with permission from D. R. Gealy, USDA-ARS)".



1832



1833

1834



1835

1836



1837

1838